

ALKAPTONURIA



A 3-in-1 Medical Reference

A Bibliography and Dictionary
for Physicians, Patients,
and Genome Researchers

TO INTERNET REFERENCES

ALKAPTONURIA

A BIBLIOGRAPHY AND
DICTIONARY

FOR PHYSICIANS, PATIENTS,
AND GENOME RESEARCHERS



JAMES N. PARKER, M.D.
AND PHILIP M. PARKER, PH.D., EDITORS

ICON Health Publications
ICON Group International, Inc.
7404 Trade Street
San Diego, CA 92121 USA

Copyright ©2007 by ICON Group International, Inc.

Copyright ©2007 by ICON Group International, Inc. All rights reserved. This book is protected by copyright. No part of it may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without written permission from the publisher.

Printed in the United States of America.

Last digit indicates print number: 10 9 8 7 6 4 5 3 2 1

Publisher, Health Care: Philip Parker, Ph.D.
Editor(s): James Parker, M.D., Philip Parker, Ph.D.

Publisher's note: The ideas, procedures, and suggestions contained in this book are not intended for the diagnosis or treatment of a health problem. As new medical or scientific information becomes available from academic and clinical research, recommended treatments and drug therapies may undergo changes. The authors, editors, and publisher have attempted to make the information in this book up to date and accurate in accord with accepted standards at the time of publication. The authors, editors, and publisher are not responsible for errors or omissions or for consequences from application of the book, and make no warranty, expressed or implied, in regard to the contents of this book. Any practice described in this book should be applied by the reader in accordance with professional standards of care used in regard to the unique circumstances that may apply in each situation. The reader is advised to always check product information (package inserts) for changes and new information regarding dosage and contraindications before prescribing any drug or pharmacological product. Caution is especially urged when using new or infrequently ordered drugs, herbal remedies, vitamins and supplements, alternative therapies, complementary therapies and medicines, and integrative medical treatments.

Cataloging-in-Publication Data

Parker, James N., 1961-
Parker, Philip M., 1960-

Alkaptonuria: A Bibliography and Dictionary for Physicians, Patients, and Genome Researchers/ James N. Parker and Philip M. Parker, editors

p. cm.

Includes bibliographical references, glossary, and index.

ISBN: 0-497-11321-X

1. Alkaptonuria-Popular works. I. Title.

Disclaimer

This publication is not intended to be used for the diagnosis or treatment of a health problem. It is sold with the understanding that the publisher, editors, and authors are not engaging in the rendering of medical, psychological, financial, legal, or other professional services.

References to any entity, product, service, or source of information that may be contained in this publication should not be considered an endorsement, either direct or implied, by the publisher, editors, or authors. ICON Group International, Inc., the editors, and the authors are not responsible for the content of any Web pages or publications referenced in this publication.

Copyright Notice

If a physician wishes to copy limited passages from this book for patient use, this right is automatically granted without written permission from ICON Group International, Inc. (ICON Group). However, all of ICON Group publications have copyrights. With exception to the above, copying our publications in whole or in part, for whatever reason, is a violation of copyright laws and can lead to penalties and fines. Should you want to copy tables, graphs, or other materials, please contact us to request permission (E-mail: iconedit@san.rr.com). ICON Group often grants permission for very limited reproduction of our publications for internal use, press releases, and academic research. Such reproduction requires confirmed permission from ICON Group International, Inc. **The disclaimer above must accompany all reproductions, in whole or in part, of this book.**

Acknowledgements

The collective knowledge generated from academic and applied research summarized in various references has been critical in the creation of this book which is best viewed as a comprehensive compilation and collection of information prepared by various official agencies which produce publications on alkaptonuria. Books in this series draw from various agencies and institutions associated with the United States Department of Health and Human Services, and in particular, the Office of the Secretary of Health and Human Services (OS), the Administration for Children and Families (ACF), the Administration on Aging (AOA), the Agency for Healthcare Research and Quality (AHRQ), the Agency for Toxic Substances and Disease Registry (ATSDR), the Centers for Disease Control and Prevention (CDC), the Food and Drug Administration (FDA), the Healthcare Financing Administration (HCFA), the Health Resources and Services Administration (HRSA), the Indian Health Service (IHS), the institutions of the National Institutes of Health (NIH), the Program Support Center (PSC), and the Substance Abuse and Mental Health Services Administration (SAMHSA). In addition to these sources, information gathered from the National Library of Medicine, the United States Patent Office, the European Union, and their related organizations has been invaluable in the creation of this book. Some of the work represented was financially supported by the Research and Development Committee at INSEAD. This support is gratefully acknowledged. Finally, special thanks are owed to Tiffany Freeman for her excellent editorial support.

About the Editors

James N. Parker, M.D.

Dr. James N. Parker received his Bachelor of Science degree in Psychobiology from the University of California, Riverside and his M.D. from the University of California, San Diego. In addition to authoring numerous research publications, he has lectured at various academic institutions. Dr. Parker is the medical editor for health books by ICON Health Publications.

Philip M. Parker, Ph.D.

Philip M. Parker is the Chaired Professor of Management Science at INSEAD (Fontainebleau, France and Singapore). Dr. Parker has also been Professor at the University of California, San Diego and has taught courses at Harvard University, the Hong Kong University of Science and Technology, the Massachusetts Institute of Technology, Stanford University, and UCLA. Dr. Parker is the associate editor for ICON Health Publications.

About ICON Health Publications

To discover more about ICON Health Publications, simply check with your preferred online booksellers, including Barnes&Noble.com and Amazon.com which currently carry all of our titles. Or, feel free to contact us directly for bulk purchases or institutional discounts:

ICON Group International, Inc.
7404 Trade Street
San Diego, CA 92121 USA
Fax: 858-635-9414
Web site: www.icongrouponline.com/health

Table of Contents

FORWARD	1
CHAPTER 1. STUDIES ON ALKAPTONURIA.....	3
<i>Overview</i>	3
<i>Genetics Home Reference</i>	3
<i>What Is Alkaptonuria?</i>	3
<i>How Common Is Alkaptonuria?</i>	4
<i>What Genes Are Related to Alkaptonuria?</i>	4
<i>How Do People Inherit Alkaptonuria?</i>	4
<i>Where Can I Find Additional Information about Alkaptonuria?</i>	4
<i>References</i>	5
<i>What Is the Official Name of the HGD Gene?</i>	6
<i>What Is the Normal Function of the HGD Gene?</i>	6
<i>What Conditions Are Related to the HGD Gene?</i>	6
<i>Where Is the HGD Gene Located?</i>	6
<i>References</i>	7
<i>Federally Funded Research on Alkaptonuria</i>	8
<i>The National Library of Medicine: PubMed</i>	9
CHAPTER 2. ALTERNATIVE MEDICINE AND ALKAPTONURIA	31
<i>Overview</i>	31
<i>National Center for Complementary and Alternative Medicine</i>	31
<i>Additional Web Resources</i>	31
<i>General References</i>	32
APPENDIX A. HELP ME UNDERSTAND GENETICS	34
<i>Overview</i>	34
<i>The Basics: Genes and How They Work</i>	34
<i>Genetic Mutations and Health</i>	45
<i>Inheriting Genetic Conditions</i>	51
<i>Genetic Consultation</i>	59
<i>Genetic Testing</i>	61
<i>Gene Therapy</i>	67
<i>The Human Genome Project and Genomic Research</i>	70
APPENDIX B. PHYSICIAN RESOURCES.....	73
<i>Overview</i>	73
<i>NIH Guidelines</i>	73
<i>NIH Databases</i>	74
<i>Other Commercial Databases</i>	77
APPENDIX C. PATIENT RESOURCES	78
<i>Overview</i>	78
<i>Patient Guideline Sources</i>	78
<i>Finding Associations</i>	80
<i>Resources for Patients and Families</i>	80
ONLINE GLOSSARIES	82
<i>Online Dictionary Directories</i>	83
ALKAPTONURIA DICTIONARY	84
INDEX.....	110

FORWARD

In March 2001, the National Institutes of Health issued the following warning: "The number of Web sites offering health-related resources grows every day. Many sites provide valuable information, while others may have information that is unreliable or misleading."¹ Furthermore, because of the rapid increase in Internet-based information, many hours can be wasted searching, selecting, and printing. Since only the smallest fraction of information dealing with alkaptonuria is indexed in search engines, such as **www.google.com** or others, a non-systematic approach to Internet research can be not only time consuming, but also incomplete. This book was created for medical professionals, students, and members of the general public who want to know as much as possible about alkaptonuria, using the most advanced research tools available and spending the least amount of time doing so.

In addition to offering a structured and comprehensive bibliography, the pages that follow will tell you where and how to find reliable information covering virtually all topics related to alkaptonuria, from the essentials to the most advanced areas of research. Special attention has been paid to present the genetic basis and pattern of inheritance of alkaptonuria. Public, academic, government, and peer-reviewed research studies are emphasized. Various abstracts are reproduced to give you some of the latest official information available to date on alkaptonuria. Abundant guidance is given on how to obtain free-of-charge primary research results via the Internet. **While this book focuses on the field of medicine, when some sources provide access to non-medical information relating to alkaptonuria, these are noted in the text.**

E-book and electronic versions of this book are fully interactive with each of the Internet sites mentioned (clicking on a hyperlink automatically opens your browser to the site indicated). If you are using the hard copy version of this book, you can access a cited Web site by typing the provided Web address directly into your Internet browser. You may find it useful to refer to synonyms or related terms when accessing these Internet databases. **NOTE:** At the time of publication, the Web addresses were functional. However, some links may fail due to URL address changes, which is a common occurrence on the Internet.

For readers unfamiliar with the Internet, detailed instructions are offered on how to access electronic resources. For readers unfamiliar with medical terminology, a comprehensive glossary is provided. We hope these resources will prove useful to the widest possible audience seeking information on alkaptonuria.

The Editors

¹ From the NIH, National Cancer Institute (NCI): <http://www.cancer.gov/>.

CHAPTER 1. STUDIES ON ALKAPTONURIA

Overview

In this chapter, we will show you how to locate peer-reviewed references and studies on alkaptonuria. For those interested in basic information about alkaptonuria, we begin with a condition summary published by the National Library of Medicine.

Genetics Home Reference

Genetics Home Reference (GHR) is the National Library of Medicine's Web site for consumer information about genetic conditions and the genes or chromosomes responsible for those conditions. Here you can find a condition summary on alkaptonuria that describes the major features of the condition, provides information about the condition's genetic basis, and explains its pattern of inheritance. In addition, a summary of the gene or chromosome related to alkaptonuria is provided.²

The Genetics Home Reference has recently published the following summary for alkaptonuria:

What Is Alkaptonuria?³

Alkaptonuria is an inherited condition that causes urine to turn black when exposed to air. Ochronosis, a buildup of dark pigment in connective tissues such as cartilage and skin, is also characteristic of the disorder. People with alkaptonuria typically develop arthritis in adulthood, particularly in the spine and large joints.

² This section has been adapted from the National Library of Medicine: <http://ghr.nlm.nih.gov/>.

³ Adapted from the Genetics Home Reference of the National Library of Medicine: <http://ghr.nlm.nih.gov/condition=alkaptonuria>.

How Common Is Alkaptonuria?

The condition is rare, affecting 1 in 250,000 to 1 million people worldwide. Alkaptonuria is more common in certain areas of Slovakia (where it has an incidence of about 1 in 19,000 people) and in the Dominican Republic.

What Genes Are Related to Alkaptonuria?

Mutations in the **HGD** (<http://ghr.nlm.nih.gov/gene=hgd>) gene cause alkaptonuria.

The HGD gene makes an enzyme called homogentisate oxidase, which helps break down the amino acids phenylalanine and tyrosine (important building blocks of proteins) into smaller molecules. Mutations in the HGD gene prevent the enzyme from playing its role in this process. As a result, a substance called homogentisic acid builds up in the body. Excess homogentisic acid is deposited in connective tissues, which causes cartilage and skin to darken and leads to arthritis. Homogentisic acid is also present in urine, which makes the urine turn dark when exposed to air.

How Do People Inherit Alkaptonuria?

This condition is inherited in an autosomal recessive pattern, which means two copies of the gene in each cell are altered. Most often, the parents of an individual with an autosomal recessive disorder each carry one copy of the altered gene but do not show signs and symptoms of the disorder.

Where Can I Find Additional Information about Alkaptonuria?

You may find the following resources about alkaptonuria helpful. These materials are written for the general public.

MedlinePlus - Health Information

- Encyclopedia: Alkaptonuria:
<http://www.nlm.nih.gov/medlineplus/ency/article/001200.htm>
- Health Topic: Metabolic Disorders:
<http://www.nlm.nih.gov/medlineplus/metabolicdisorders.html>

Educational Resources - Information Pages

- Madisons Foundation:
<http://www.madisonsfoundation.org/content/3/1/display.asp?did=378>
- Medical College of Wisconsin:
<http://healthlink.mcw.edu/article/921733488.html>
- Orphanet:
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=GB&Expert=56

Patient Support - for Patients and Families

- Children Living with Inherited Metabolic Diseases (CLIMB) (UK):
<http://www.climb.org.uk>
- National Organization for Rare Disorders:
http://www.rarediseases.org/search/rdbdetail_abstract.html?disname=Alkaptonuria

Professional Resources

You may also be interested in these resources, which are designed for healthcare professionals and researchers.

- Gene Reviews - Clinical summary:
<http://www.genetests.org/query?dz=alkap>
- Gene Tests - DNA tests ordered by healthcare professionals:
<http://www.genetests.org/query?testid=22213>
- ClinicalTrials.gov - Linking patients to medical research:
<http://clinicaltrials.gov/search/condition=%22alkaptonuria%22?recruiting=false>
- PubMed - Recent literature:
<http://ghr.nlm.nih.gov/condition=alkaptonuria/show/PubMed;jsessionid=A7671B80A5FD213E742DB28C0835FB84>
- OMIM - Genetic disorder catalog:
<http://www.ncbi.nlm.nih.gov/entrez/dispomim.cgi?id=203500>

References

These sources were used to develop the Genetics Home Reference condition summary on alkaptonuria.

- Gene Review
- Lubics A, Schneider I, Sebok B, Havass Z. Extensive bluish gray skin pigmentation and severe arthropathy. Endogenous ochronosis (alkaptonuria). Arch Dermatol. 2000 Apr;136(4):548-9, 551-2. Review. No abstract available. PubMed citation
- Phornphutkul C, Introne WJ, Perry MB, Bernardini I, Murphey MD, Fitzpatrick DL, Anderson PD, Huizing M, Anikster Y, Gerber LH, Gahl WA. Natural history of alkaptonuria. N Engl J Med. 2002 Dec 26;347(26):2111-21. PubMed citation
- Scriver, Charles R; The metabolic & molecular bases of inherited disease; 8th ed.; New York : McGraw-Hill, c2001. p2109-2123. NLM Catalog

A summary of the gene related to alkaptonuria is provided below:

What Is the Official Name of the HGD Gene?⁴

The official name of this gene is “homogentisate 1,2-dioxygenase (homogentisate oxidase).”

HGD is the gene's official symbol. The HGD gene is also known by other names, listed below.

What Is the Normal Function of the HGD Gene?

Homogentisate oxidase, the protein made by the HGD gene, is active chiefly in the liver and kidneys. It is one of a series of enzymes needed to break down two molecular building blocks, the amino acids phenylalanine and tyrosine, when they are no longer needed or are present in excess. These two amino acids play a role in making proteins, certain hormones, pigments, and brain chemicals called neurotransmitters.

Homogentisate oxidase is responsible for one specific step in the breakdown of phenylalanine and tyrosine. Previous steps convert the two amino acids into a molecule called homogentisic acid. Homogentisate oxidase adds two oxygen atoms to homogentisic acid, converting it to another molecule called maleylacetoacetate. Other enzymes break down maleylacetoacetate into smaller molecules that are later used for energy or to make other products that can be used by the body.

What Conditions Are Related to the HGD Gene?

Alkaptonuria - Caused by Mutations in the HGD Gene


More than 40 mutations in the HGD gene have been identified in people with alkaptonuria. Many of these mutations change single amino acids in the homogentisate oxidase protein. For example, a substitution of the amino acid valine for methionine at position 368 (also written as Met368Val) is the most common HGD mutation in European populations. Mutations in the HGD gene probably inactivate the enzyme by changing its structure. Without active homogentisate oxidase, homogentisic acid builds up in the body. It is deposited in connective tissues such as cartilage and skin, which causes them to darken and leads to arthritis. Homogentisic acid is also present in urine, causing the urine to turn black when exposed to air.

Where Is the HGD Gene Located?

Cytogenetic Location: 3q21-q23

Molecular Location on chromosome 3: base pairs 121,822,762 to 121,889,491

⁴ Adapted from the Genetics Home Reference of the National Library of Medicine:
<http://ghr.nlm.nih.gov/gene=hgd;jsessionid=A7671B80A5FD213E742DB28C0835FB84>.



The HGD gene is located on the long (q) arm of chromosome 3 between positions 21 and 23.

More precisely, the HGD gene is located from base pair 121,822,762 to base pair 121,889,491 on chromosome 3.

References

These sources were used to develop the Genetics Home Reference gene summary on the HGD gene.

- Amaya AA, Brzezinski KT, Farrington N, Moran GR. Kinetic analysis of human homogentisate 1,2-dioxygenase. *Arch Biochem Biophys*. 2004 Jan 1;421(1):135-42. PubMed citation
- Goicoechea De Jorge E, Lorda I, Gallardo ME, Perez B, Perez De Ferran C, Mendoza H, Rodriguez De Cordoba S. Alkaptonuria in the Dominican Republic: identification of the founder AKU mutation and further evidence of mutation hot spots in the HGO gene. *J Med Genet*. 2002 Jul;39(7):E40. No abstract available. PubMed citation
- Rodriguez JM, Timm DE, Titus GP, Beltran-Valero De Bernabe D, Criado O, Mueller HA, Rodriguez De Cordoba S, Penalva MA. Structural and functional analysis of mutations in alkaptonuria. *Hum Mol Genet*. 2000 Sep 22;9(15):2341-50. PubMed citation
- Scriver, Charles R; *The metabolic & molecular bases of inherited disease*; 8th ed.; New York : McGraw-Hill, c2001. p2113-2120. NLM Catalog
- Srsen S, Muller CR, Fregin A, Srsnova K. Alkaptonuria in Slovakia: thirty-two years of research on phenotype and genotype. *Mol Genet Metab*. 2002 Apr;75(4):353-9. Review. PubMed citation
- Zatkova A, Chmelikova A, Polakova H, Ferakova E, Kadasi L. Rapid detection methods for five HGO gene mutations causing alkaptonuria. *Clin Genet*. 2003 Feb;63(2):145-9. PubMed citation

Federally Funded Research on Alkaptonuria

The U.S. Government supports a variety of research studies relating to alkaptonuria. These studies are tracked by the Office of Extramural Research at the National Institutes of Health.⁵

CRISP (Computerized Retrieval of Information on Scientific Projects)

CRISP is a searchable database of federally funded biomedical research projects conducted at universities, hospitals, and other institutions. Search the CRISP Web site at http://crisp.cit.nih.gov/crisp/crisp_query.generate_screen. You will have the option to perform targeted searches by various criteria, including geography, date, and topics related to alkaptonuria.

For most of the studies, the agencies reporting into CRISP provide summaries or abstracts. As opposed to clinical trial research using patients, many federally funded studies use animals or simulated models to explore alkaptonuria. The following is typical of the type of information found when searching the CRISP database for alkaptonuria:

- **Project Title: PATHOPHYSIOLOGY AND TREATMENT OF DISORDERS IN THE DISTAL**

Principal Investigator & Institution: Grompe, Markus; Professor; Molecular and Medical Genetics; Oregon Health & Science University Portland, or 972393098

Timing: Fiscal Year 2004; Project Start 01-DEC-1993; Project End 14-FEB-2005

Summary: (Copied from Applicant Abstract): Homogentisic acid dioxygenase (HGD), maleylacetoacetate isomerase (MAI) and fumarylacetoacetate hydrolase (FAH) sequentially catalyze the terminal degradation of tyrosine into Krebs cycle intermediates. Deficiency of HGD causes the human disorder **alkaptonuria**, (AKU) deficiency of FAH causes tyrosinemia type I (HT1) and the phenotype of MAI deficiency in man is unknown. HT1 is characterized by severe liver dysfunction, renal tubular damage and finally liver cancer. Previously we have generated a murine model of HT 1 and shown that the drug NTBC can prevent the neonatal lethality and liver dysfunction of FAH deficient mice. We also demonstrated a strong selective growth advantage of FAH expressing cells in mutant liver. This in vivo selection resulted in a near complete repopulation of diseased liver with healthy tissue and raised the hope that human HT1 may be curable by gene therapy. To broaden our understanding of the pathophysiology and potential therapy of HT1 we have now also established murine models of HGD and MAI deficiency. This application pertains to the use of murine models of enzyme deficiencies in tyrosine catabolism to explore the biology of novel strategies for liver and kidney gene therapy, to better understand the etiology of hepatic cancer in HT1 and dissect the pathophysiology of the diseases in this pathway.

Website: http://crisp.cit.nih.gov/crisp/Crisp_Query.Generate_Screen

⁵ Healthcare projects are funded by the National Institutes of Health (NIH), Substance Abuse and Mental Health Services (SAMHSA), Health Resources and Services Administration (HRSA), Food and Drug Administration (FDA), Centers for Disease Control and Prevention (CDCP), Agency for Healthcare Research and Quality (AHRQ), and Office of Assistant Secretary of Health (OASH).

- **Project Title: SPECTROSCOPIC STUDIES OF MONONUCLEAR NON-HEME FE ENZYMES**

Principal Investigator & Institution: Solomon, Edward I.; Professor; Chemistry; Stanford University 1215 Welch Road, Mod B Stanford, Ca 943055402

Timing: Fiscal Year 2004; Project Start 01-JUL-1988; Project End 30-JUN-2007

Summary: (provided by applicant): Mononuclear non-heme iron enzymes are involved in a wide range of reactions with oxygen including dioxygenation, hydroxylation, desaturation, 4-electron oxidation and H-atom abstraction. These enzymes are involved in the biosynthesis of antibiotics (penicillins, cephalosporins, vancomycin) and lactamase inhibitors (clavulanic acid), associated with genetic diseases (phenylketonuria, **alkaptonuria**, tyrosinemia, Refsum's disease), involved in the biosynthesis of leukotrienes and lipoxins (lipoxygenase (LO)), important in the bioremediation of pollutants including PCBs, and in the treatment of certain cancers (Bleomycin). These enzymes have been much more difficult to study than heme systems as they do not have the dominant spectroscopic features of the porphyrin and mostly activate dioxygen from a high-spin ferrous site which is a non-Kramers ion and generally not EPR active. New methods have been developed for the study of this wide class of enzymes including variable-temperature variable-field magnetic circular dichroism (VTVH MCD) of non-Kramers ions, VTVH MCD/zero field splitting/density functional theory of Kramers ions to define electronic structure of oxygen intermediates, and L-edge X-ray absorption spectroscopy (XAS) to define the covalencies of the metal orbitals. Application of these methods has thus far determined a general mechanistic strategy utilized by many members of this class of enzymes, defined the nature of the highly covalent Tyr-Fe III bond in the intradiol dioxygenases, and steps in the mechanism of H-atom abstraction by LO, determined the electronic structure of activated bleomycin (involved in H-atom abstraction from DNA) and shown its reactivity to be fundamentally different from that of heme systems, and developed insight into its reaction coordinate. These studies are now focused on: 1) Extending the VTVH MCD studies to define structure/function correlations for a wide range of members of this class of enzymes and their mutants related to genetic diseases; 2) understanding O₂ reaction coordinates of the ferrous sites with different co-factors to generate reactive oxygen intermediates; 3) trapping and determining the geometric and electronic structures of these oxygen intermediates and their differences in reactivity (hydroxylation, desaturation, etc.); and 4) utilizing the L-edge XAS method to define electronic structure differences between heme and non-heme iron sites which correlate with differences in O₂ activation.

Website: http://crisp.cit.nih.gov/crisp/Crisp_Query.Generate_Screen

The National Library of Medicine: PubMed

One of the quickest and most comprehensive ways to find academic studies in both English and other languages is to use PubMed, maintained by the National Library of Medicine.⁶ The advantage of PubMed over previously mentioned sources is that it covers a greater number of domestic and foreign references. It is also free to use. If the publisher has a Web site that offers full text of its journals, PubMed will provide links to that site, as well as to

⁶ PubMed was developed by the National Center for Biotechnology Information (NCBI) at the National Library of Medicine (NLM) at the National Institutes of Health (NIH). The PubMed database was developed in conjunction with publishers of biomedical literature as a search tool for accessing literature citations and linking to full-text journal articles at Web sites of participating publishers. Publishers that participate in PubMed supply NLM with their citations electronically prior to or at the time of publication.

sites offering other related data. User registration, a subscription fee, or some other type of fee may be required to access the full text of articles in some journals.

To generate your own bibliography of studies dealing with alkaptonuria, simply go to the PubMed Web site at <http://www.ncbi.nlm.nih.gov/pubmed>. Type **alkaptonuria** (or synonyms) into the search box, and click **Go**. The following is the type of output you can expect from PubMed for alkaptonuria (hyperlinks lead to article summaries):

- **A 46-year-old man with a 4-year history of diffuse brownish black pigmentation. Endogenous ochronosis (alkaptonuria).**
 Author(s): Micali G, Di Stefano AG, Nasca MR, Musumeci ML.
 Source: Archives of Dermatology.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9449917&query_hl=13&itool=pubmed_docsum

- **A case of alkaptonuria with root canal stenosis.**
 Author(s): Koh KB, Low EH, Ch'ng SL, Zakiah I.
 Source: Singapore Med J.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=8009267&query_hl=13&itool=pubmed_docsum

- **A case of alkaptonuria, ochronosis and ochronotic arthropathy.**
 Author(s): Entwisle BR, Muirden KD, Adam WR, Parkin JD.
 Source: The Medical Journal of Australia.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5803867&query_hl=13&itool=pubmed_docsum

- **A family with alkaptonuria showing quasidominant inheritance.**
 Author(s): de Silva AP, Seneviratne SL, Gunatilake SB, Fonseka M, Jayasekera R, de Silva HJ.
 Source: Ceylon Med J.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10675999&query_hl=13&itool=pubmed_docsum

- **A fungal perspective on human inborn errors of metabolism: alkaptonuria and beyond.**
 Author(s): Penalva MA.
 Source: Fungal Genetics and Biology : Fg & B.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=11567547&query_hl=13&itool=pubmed_docsum

- **A novel point mutation associated with alkaptonuria.**
 Author(s): Higashino K, Liu W, Ohkawa T, Yamamoto T, Fukui K, Ohno M, Imanishi H, Iwasaki A, Amuro Y, Hada T.
 Source: Clinical Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9630082&query_hl=13&itool=pubmed_docsum

- **aku, a mutation of the mouse homologous to human alkaptonuria, maps to chromosome 16.**
 Author(s): Montagutelli X, Lalouette A, Coude M, Kamoun P, Forest M, Guenet JL.
 Source: Genomics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=8188247&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria (a case report).**
 Author(s): Sachdeva JR, Gupta MM, Jindal VK.
 Source: Journal of Postgraduate Medicine.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=3795131&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria and aortic stenosis.**
 Author(s): Vlay SC, Hartman AR, Culliford AT.
 Source: Annals of Internal Medicine.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=3946992&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria and Gilbert's syndrome. Report of two affected siblings and hepatic ultrastructure in one sibling.**
 Author(s): Brown NK, Smuckler EA.
 Source: The American Journal of Medicine.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5420562&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria and lumbar disc herniation. Report of three cases.**
 Author(s): Farzannia A, Shokouhi G, Hadidchi S.
 Source: Journal of Neurosurgery.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12546396&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria and ochronosis in three siblings. Ascorbic acid treatment monitored by urinary HGA excretion.**
 Author(s): Forslind K, Wollheim FA, Akesson B, Rydholm U.
 Source: Clin Exp Rheumatol.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=3180550&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria and ochronosis.**
 Author(s): Blumsohn D.
 Source: South African Medical Journal. Suid-Afrikaanse Tydskrif Vir Geneeskunde.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=594852&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria and ochronosis. A survey and 5 cases.**
Author(s): Christensen K, Manthorpe R.
Source: Human Heredity.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=6862457&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria and ochronosis: case report and review.**
Author(s): Albers SE, Brozena SJ, Glass LF, Fenske NA.
Source: Journal of the American Academy of Dermatology.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=1401313&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria and ochronotic arthropathy.**
Author(s): Mam MK, Sethi TS.
Source: J Indian Med Assoc.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=3794358&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria and photography: a patient's urine tells the story.**
Author(s): Maxwell D.
Source: Cmaj : Canadian Medical Association Journal = Journal De L'association Medicale Canadienne.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=15824403&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria and renal failure: a case report and review of the literature.**
Author(s): Venkateshan VS, Chandra B, Graziano V, Steinlauf P, Marquet E, Irmieri V, Needle MA.
Source: Modern Pathology : an Official Journal of the United States and Canadian Academy of Pathology, Inc.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=1495952&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria and renal failure: a case report.**
Author(s): Kazancioglu R, Taylan I, Aksak F, Durak H, Kumbasar B, Yenigun M, Sar F.
Source: Journal of Nephrology.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=15365968&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria caused by compound heterozygote mutations.**
Author(s): Elcioglu NH, Aytug AF, Muller CR, Gurbuz O, Ergun T, Kotiloglu E, Elcioglu M.
Source: Genet Couns.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12872815&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria detected by neuroblastoma screening system.**
 Author(s): Sakura N, Kato Y, Hamakawa M, Yamaguchi S.
 Source: European Journal of Pediatrics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=1396899&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria diagnosed by flexible bronchoscopy.**
 Author(s): Parambil JG, Daniels CE, Zehr KJ, Utz JP.
 Source: Chest.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=16304331&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria in a Chinese baby.**
 Author(s): Wang TR, Hwu WL.
 Source: Journal of Inherited Metabolic Disease.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=2515375&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria in an Ethiopian.**
 Author(s): Lester FT, Tereschenko OI.
 Source: East Afr Med J.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=747995&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria in an orangutan (*Pongo pygmaeus*).**
 Author(s): Keeling ME, McClure HM, Kibler RF.
 Source: American Journal of Physical Anthropology.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=4689767&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria in Italy: polymorphic haplotype background, mutational profile, and description of four novel mutations in the homogentisate 1,2-dioxygenase gene.**
 Author(s): Porfirio B, Chiarelli I, Graziano C, Mannoni A, Morrone A, Zammarchi E, De Bernabe DB, De Cordoba SR.
 Source: Journal of Medical Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10819641&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria in siblings.**
 Author(s): Reddi YR, Rao MN, Laxman S.
 Source: Indian J Pediatr.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=4784732&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria in Slovakia: thirty-two years of research on phenotype and genotype.**
Author(s): Srsen S, Muller CR, Fregin A, Srsnova K.
Source: Molecular Genetics and Metabolism.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12051967&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria in the Dominican Republic: identification of the founder AKU mutation and further evidence of mutation hot spots in the HGO gene.**
Author(s): Goicoechea De Jorge E, Lorda I, Gallardo ME, Perez B, Perez De Ferran C, Mendoza H, Rodriguez De Cordoba S.
Source: Journal of Medical Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12114497&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria in the Trencin District of Czechoslovakia.**
Author(s): Srsen S, Cisarik F, Pasztor L, Harmecko L.
Source: American Journal of Medical Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=263435&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria presenting as lumbar disc prolapse: case report and review of literature.**
Author(s): Reddy DR, Prasad VS.
Source: Spinal Cord : the Official Journal of the International Medical Society of Paraplegia.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9670391&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria with atypical joint involvement.**
Author(s): Alrehaily A, Pope JE.
Source: The Journal of Rheumatology.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=11833486&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria with extensive ochronosis.**
Author(s): Wyre HW Jr.
Source: Archives of Dermatology.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=434871&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria with lumbar disc herniation: a report of two cases.**
Author(s): Emel E, Karagoz F, Aydin IH, Hacisalihoglu S, Seyithanoglu MH.
Source: Spine.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10954648&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria with nephrocalcinosis.**
 Author(s): Goldberg BH, Penso JS, Stern LM, Bergstein JM.
 Source: The Journal of Pediatrics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=1245970&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria with ochronotic nephropathy and multiple joint replacement for ochronotic arthropathy.**
 Author(s): Jagose JT, Bailey RR, Rothwell AG.
 Source: N Z Med J.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9236810&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria with ochronotic spondylosis.**
 Author(s): Rao SB, Rao KS, Dinakar I.
 Source: J Indian Med Assoc.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5414387&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria, ochronosis and ochronotic arthritis.**
 Author(s): Molony J, Kelly DJ.
 Source: J Ir Med Assoc.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5411095&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria, ochronosis, and low-back pain. A case report.**
 Author(s): McCollum DE, Odom GL.
 Source: The Journal of Bone and Joint Surgery. American Volume.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=4220888&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria, ochronosis, and ochronotic arthropathy.**
 Author(s): Mannoni A, Selvi E, Lorenzini S, Giorgi M, Airo P, Cammelli D, Andreotti L, Marcolongo R, Porfirio B.
 Source: Seminars in Arthritis and Rheumatism.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=14978662&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria.**
 Author(s): Lorenzini S, Mannoni A, Selvi E.
 Source: The New England Journal of Medicine.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12672874&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria.**
Author(s): Carlesimo M, Bonaccorsi P, Tamburrano G, Carboni I, Parisi A, Calvier S.
Source: Dermatology (Basel, Switzerland).
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10449966&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria.**
Author(s): Singh BN, Mehrotra S, Bhargava B, Agarwal S.
Source: J Assoc Physicians India.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=2632541&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria.**
Author(s): Brenton DP, Krywawych S.
Source: Clin Rheum Dis.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=3078781&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria.**
Author(s): Biswas TK, Nayek G, Chatterjee A, Bhattacharjee D, Halder R, Roy DN.
Source: J Indian Med Assoc.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=4078344&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria.**
Author(s): Jayasinghe KS, Wijesiriwardena BC, Sivaramana V, Sheriff MH, Dharmadasa K.
Source: Ceylon Med J.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=6534535&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria.**
Author(s): Kotwal MR, Rinchhen CZ.
Source: J Indian Med Assoc.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=6863950&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria.**
Author(s): Srsen S.
Source: Johns Hopkins Med J.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=513428&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria.**
Author(s): Sussman T.
Source: Proc R Soc Med.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5770593&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria. (A case report).**
 Author(s): Madan MS.
 Source: Indian Pediatrics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5478380&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria. A case report.**
 Author(s): Adams EJ, Ward JR.
 Source: Rocky Mt Med J.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=1114286&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria. A case report.**
 Author(s): Sainani GS, Balsara AB.
 Source: Indian Journal of Medical Sciences.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5576749&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria. A report of three cases from one family tree.**
 Author(s): Kulkarni PS, Shah KA, Venugopal AR, Shah SS.
 Source: Journal of Postgraduate Medicine.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=4792968&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria. Report of a case in a neonate.**
 Author(s): Saxena H, Srivastava JR, Goswami P.
 Source: Indian J Pediatr.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5375503&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria. Report of two cases in siblings.**
 Author(s): Verma IC.
 Source: Indian J Pediatr.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5719646&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria: (a case report).**
 Author(s): Meundi DB, Bhandari SG.
 Source: Indian Pediatrics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=700885&query_hl=13&itool=pubmed_docsum
- **Alkaptonuria: a case report.**
 Author(s): Odabas AR, Karakuzu A, Selcuk Y, Erdem T, Cetinkaya R.
 Source: The Journal of Dermatology.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=11349468&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria: early detection.**
 Author(s): Khadagawat R, Teckchandani R, Garg P, Arya A, Choudhary B.
 Source: Indian Pediatrics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=7875895&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria: from humans to moulds and back.**
 Author(s): Scazzocchio C.
 Source: Trends in Genetics : Tig.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9097720&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria: prevalence among Bedouins related by consanguinity.**
 Author(s): Modilevsky T, Manor E, Marmor A.
 Source: Monogr Hum Genet.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=732828&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria: report of six cases.**
 Author(s): Nora JR, Nora GE, Fitzgerald M.
 Source: Imj Ill Med J.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=4383995&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria: such a long journey.**
 Author(s): Scriver CR.
 Source: Nature Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=8782808&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria: tracked down to chromosome 3.**
 Author(s): McKusick VA.
 Source: Genomics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=8188236&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria--a rare presentation.**
 Author(s): Sharma JK, Dhawan SN, Singh V, Bhandari VW.
 Source: J Assoc Physicians India.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=3182707&query_hl=13&itool=pubmed_docsum

- **Alkaptonuria--a series of seven cases.**
 Author(s): Parikh A, Khubchandani RP, Bharucha BA, Kumta NB, Pandya MB, Naik G.
 Source: J Assoc Physicians India.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=3074083&query_hl=13&itool=pubmed_docsum

- **Allelic heterogeneity of alkaptonuria in Central Europe.**
 Author(s): Muller CR, Fregin A, Srsen S, Srsnova K, Halliger-Keller B, Felbor U, Seemanova E, Kress W.
 Source: European Journal of Human Genetics : Ejhg.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10482952&query_hl=13&itool=pubmed_docsum
- **An autopsy case of alkaptonuria with ochronosis.**
 Author(s): Nishimori I, Ito J, Hakuno T.
 Source: Acta Pathol Jpn.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5537432&query_hl=13&itool=pubmed_docsum
- **Analysis for homogentisic acid by NMR spectrometry, to aid diagnosis of alkaptonuria.**
 Author(s): Yamaguchi S, Koda N, Yamamoto H.
 Source: Clinical Chemistry.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=2758664&query_hl=13&itool=pubmed_docsum
- **Analysis of alkaptonuria (AKU) mutations and polymorphisms reveals that the CCC sequence motif is a mutational hot spot in the homogentisate 1,2 dioxygenase gene (HGO).**
 Author(s): Beltran-Valero de Bernabe D, Jimenez FJ, Aquaron R, Rodriguez de Cordoba S.
 Source: American Journal of Human Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10205262&query_hl=13&itool=pubmed_docsum
- **Analysis of alkaptonuria incidence in one region of Northwest Slovakia: a preliminary report.**
 Author(s): Neuwirth A, Srsen S, Petricek J, Mruskovic.
 Source: Birth Defects Orig Artic Ser.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=4462635&query_hl=13&itool=pubmed_docsum
- **Answer please. Alkaptonuria and ochronosis.**
 Author(s): Bosacco SJ, Fye M, Finn CA.
 Source: Orthopedics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=2020639&query_hl=13&itool=pubmed_docsum
- **Anterior megalophthalmos associated with lamellar cataract on alkaptonuria.**
 Author(s): Rao VA.
 Source: Indian J Ophthalmol.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=7141594&query_hl=13&itool=pubmed_docsum

- **Aortic stenosis and coronary artery disease caused by alkaptonuria, a rare genetic metabolic syndrome.**
Author(s): Vavuranakis M, Triantafillidi H, Stefanadis C, Toutouzas P.
Source: *Cardiology*.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10085494&query_hl=13&itool=pubmed_docsum
- **Aortic valve regurgitation in alkaptonuria.**
Author(s): Yoshikai M, Murayama J, Yamada N.
Source: *J Heart Valve Dis*.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=15473492&query_hl=13&itool=pubmed_docsum
- **Aortic valve stenosis in alkaptonuria. Images in cardiovascular medicine.**
Author(s): Hangaishi M, Taguchi J, Ikari Y, Ohno M, Kurokawa K, Kotsuka Y, Furuse A.
Source: *Circulation*.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9736602&query_hl=13&itool=pubmed_docsum
- **Are we ready to try to cure alkaptonuria?**
Author(s): La Du BN Jr.
Source: *American Journal of Human Genetics*.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9529368&query_hl=13&itool=pubmed_docsum
- **Ascorbic acid and alkaptonuria.**
Author(s): Kamoun P, Coude M, Forest M, Montagutelli X, Guenet JL.
Source: *European Journal of Pediatrics*.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=1537362&query_hl=13&itool=pubmed_docsum
- **Bilateral hip arthroplasty in alkaptonuria.**
Author(s): Warren NP, Coombs RR.
Source: *Journal of the Royal College of Surgeons of Edinburgh*.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=2355378&query_hl=13&itool=pubmed_docsum
- **Black-copper reduction reaction simulating alkaptonuria. Occurrence after intravenous urography.**
Author(s): Lee S, Schoen I.
Source: *The New England Journal of Medicine*.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5943272&query_hl=13&itool=pubmed_docsum

- **Bluish patches on the ears and axillae with dark urine: ochronosis and alkaptonuria.**
 Author(s): Turiansky GW, Levin SW.
 Source: International Journal of Dermatology.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=11554996&query_hl=13&itool=pubmed_docsum
- **Characterization of the pigment from homogentisic acid and urine and tissue from an alkaptonuria patient.**
 Author(s): Menon IA, Persad SD, Haberman HF, Basu PK, Norfray JF, Felix CC, Kalyanaraman B.
 Source: Biochemistry and Cell Biology = Biochimie Et Biologie Cellulaire.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=1647170&query_hl=13&itool=pubmed_docsum
- **Cloning of the homogentisate 1,2-dioxygenase gene, the key enzyme of alkaptonuria in mouse.**
 Author(s): Schmidt SR, Gehrig A, Koehler MR, Schmid M, Muller CR, Kress W.
 Source: Mammalian Genome : Official Journal of the International Mammalian Genome Society.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9069115&query_hl=13&itool=pubmed_docsum
- **Dark pigmentation of ear cerumen in alkaptonuria.**
 Author(s): Srsen S.
 Source: Lancet.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=79943&query_hl=13&itool=pubmed_docsum
- **Detection of alkaptonuria and mapping of its occurrence in Slovakia.**
 Author(s): Srsen S, Koska L, Kapralik I, Sitaj S, Srsnova K.
 Source: Czech Med.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=535504&query_hl=13&itool=pubmed_docsum
- **Diagnosis of alkaptonuria by NMR urinalysis: rapid qualitative and quantitative analysis of homogentisic acid.**
 Author(s): Yamaguchi S, Koda N, Ohashi T.
 Source: The Tohoku Journal of Experimental Medicine.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=3810644&query_hl=13&itool=pubmed_docsum
- **Diagnosis of alkaptonuria in children.**
 Author(s): Srsen S, Srsnova K.
 Source: Padiatr Padol.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=460920&query_hl=13&itool=pubmed_docsum

- **Discoloration of the teeth in alkaptonuria (ochronosis) and parkinsonism.**
 Author(s): Siekert RG, Gibilisco JA.
 Source: Oral Surg Oral Med Oral Pathol.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5262835&query_hl=13&itool=pubmed_docsum

- **Dura mater involvement in ochronosis (alkaptonuria).**
 Author(s): Liu W, Prayson RA.
 Source: Archives of Pathology & Laboratory Medicine.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=11419988&query_hl=13&itool=pubmed_docsum

- **Early detection of alkaptonuria.**
 Author(s): Verma SB.
 Source: Indian J Dermatol Venereol Leprol.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=16394410&query_hl=13&itool=pubmed_docsum

- **Effects of ascorbic acid and low-protein diet in alkaptonuria.**
 Author(s): Mayatepek E, Kallas K, Anninos A, Muller E.
 Source: European Journal of Pediatrics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9809834&query_hl=13&itool=pubmed_docsum

- **Effects of ascorbic acid in alkaptonuria: alterations in benzoquinone acetic acid and an ontogenic effect in infancy.**
 Author(s): Wolff JA, Barshop B, Nyhan WL, Leslie J, Seegmiller JE, Gruber H, Garst M, Winter S, Michals K, Matalon R.
 Source: Pediatric Research.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=2771520&query_hl=13&itool=pubmed_docsum

- **Exacerbation of the ochronosis of alkaptonuria due to renal insufficiency and improvement after renal transplantation.**
 Author(s): Introne WJ, Phornphutkul C, Bernardini I, McLaughlin K, Fitzpatrick D, Gahl WA.
 Source: Molecular Genetics and Metabolism.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12359141&query_hl=13&itool=pubmed_docsum

- **Extensive bluish gray skin pigmentation and severe arthropathy. Endogenous ochronosis (alkaptonuria).**
 Author(s): Lubics A, Schneider I, Sebok B, Havass Z.
 Source: Archives of Dermatology.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10768658&query_hl=13&itool=pubmed_docsum

- **Glycosuria in alkaptonuria—a dilemma.**
 Author(s): Kelley WN, Fox IH, Feldman JM, Wyngaarden JB.
 Source: Lancet.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=4188122&query_hl=13&itool=pubmed_docsum

- **High frequency of alkaptonuria in Slovakia: evidence for the appearance of multiple mutations in HGO involving different mutational hot spots.**
 Author(s): Zatkova A, de Bernabe DB, Polakova H, Zvarik M, Ferakova E, Bosak V, Ferak V, Kadasi L, de Cordoba SR.
 Source: American Journal of Human Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=11017803&query_hl=13&itool=pubmed_docsum

- **HLA antigens and alkaptonuria.**
 Author(s): Pourel GJ, Raffoux C, Faure G, Netter P, Streiff F.
 Source: J Rheumatol Suppl.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=266609&query_hl=13&itool=pubmed_docsum

- **Homozygosity mapping of the gene for alkaptonuria to chromosome 3q2.**
 Author(s): Pollak MR, Chou YH, Cerda JJ, Steinmann B, La Du BN, Seidman JG, Seidman CE.
 Source: Nature Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=8252048&query_hl=13&itool=pubmed_docsum

- **Imaging in alkaptonuria.**
 Author(s): Misson N, Van Ruyssevelt C, Mullier J.
 Source: J Belge Radiol.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=8767836&query_hl=13&itool=pubmed_docsum

- **Increased urolithiasis in patients with alkaptonuria in childhood.**
 Author(s): Zibolen M, Srsnova K, Srsen S.
 Source: Clinical Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10945668&query_hl=13&itool=pubmed_docsum

- **Interference in urine oxalate assay (sigma diagnostics oxalate oxidase method) from homogentisic acid in alkaptonuria.**
 Author(s): Biggs PA, Middleton JE, Welch RP.
 Source: Clinical Chemistry.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=3731474&query_hl=13&itool=pubmed_docsum

- **Intermittent alkaptonuria.**
Author(s): Amaresan MS, Bhupati D, Rao PV.
Source: J Indian Med Assoc.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5506301&query_hl=13&itool=pubmed_docsum
- **Minocycline-induced hyperpigmentation masquerading as alkaptonuria in individuals with joint pain.**
Author(s): Suwannarat P, Phornphutkul C, Bernardini I, Turner M, Gahl WA.
Source: Arthritis and Rheumatism.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=15529343&query_hl=13&itool=pubmed_docsum
- **Molecular analyses of the HGO gene mutations in Turkish alkaptonuria patients suggest that the R58fs mutation originated from central Asia and was spread throughout Europe and Anatolia by human migrations.**
Author(s): Uyguner O, Goicoechea de Jorge E, Cefle A, Baykal T, Kayserili H, Cefle K, Demirkol M, Yuksel-Apak M, Rodriguez de Cordoba S, Wollnik B.
Source: Journal of Inherited Metabolic Disease.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12872836&query_hl=13&itool=pubmed_docsum
- **Molecular defects in alkaptonuria.**
Author(s): Gehrig A, Schmidt SR, Muller CR, Srsen S, Srsnova K, Kress W.
Source: Cytogenetics and Cell Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9154114&query_hl=13&itool=pubmed_docsum
- **Molecular diagnosis of alkaptonuria mutation by analysis of homogentisate 1,2-dioxygenase mRNA from urine and blood.**
Author(s): Ramos SM, Hernandez M, Roces A, Larruga JM, Gonzalez P, Gonzalez AM, Pinto FM, Cabrera VM.
Source: American Journal of Medical Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9674916&query_hl=13&itool=pubmed_docsum
- **Mutation and polymorphism analysis of the human homogentisate 1, 2-dioxygenase gene in alkaptonuria patients.**
Author(s): Beltran-Valero de Bernabe D, Granadino B, Chiarelli I, Porfirio B, Mayatepek E, Aquaron R, Moore MM, Festen JJ, Sanmarti R, Penalva MA, de Cordoba SR.
Source: American Journal of Human Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9529363&query_hl=13&itool=pubmed_docsum

- **Mutational analysis of the HGO gene in Finnish alkaptonuria patients.**
 Author(s): Beltran-Valero de Bernabe D, Peterson P, Luopajarvi K, Matintalo P, Alho A, Konttinen Y, Krohn K, Rodriguez de Cordoba S, Ranki A.
 Source: Journal of Medical Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10594001&query_hl=13&itool=pubmed_docsum
- **Natural history of alkaptonuria.**
 Author(s): Phornphutkul C, Introne WJ, Perry MB, Bernardini I, Murphey MD, Fitzpatrick DL, Anderson PD, Huizing M, Anikster Y, Gerber LH, Gahl WA.
 Source: The New England Journal of Medicine.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12501223&query_hl=13&itool=pubmed_docsum
- **Neonatal severe primary hyperparathyroidism and alkaptonuria in a boy born to related parents with familial hypocalciuric hypercalcemia.**
 Author(s): Steinmann B, Gnehm HE, Rao VH, Kind HP, Prader A.
 Source: Helv Paediatr Acta.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=6543841&query_hl=13&itool=pubmed_docsum
- **Novel mutations in the homogentisate-1,2-dioxygenase gene identified in Slovak patients with alkaptonuria.**
 Author(s): Zatkova A, Polakova H, Micutkova L, Zvarik M, Bosak V, Ferakova E, Matussek J, Ferak V, Kadasi L.
 Source: Journal of Medical Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10970188&query_hl=13&itool=pubmed_docsum
- **NTBC and alkaptonuria.**
 Author(s): Anikster Y, Nyhan WL, Gahl WA.
 Source: American Journal of Human Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9718357&query_hl=13&itool=pubmed_docsum
- **Ochronosis and alkaptonuria. Case report and literature review.**
 Author(s): Kneebone TS, Selner AJ.
 Source: Journal of the American Podiatric Medical Association.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=7473092&query_hl=13&itool=pubmed_docsum
- **Ochronosis and alkaptonuria: report of a new case with calcified aortic valve stenosis.**
 Author(s): Dereymaeker L, Van Parijs G, Bayart M, Daenen W, Lauwerijns J, De Geest H.
 Source: Acta Cardiol.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=2316305&query_hl=13&itool=pubmed_docsum

- **Ochronotic arthropathy: disappearance of alkaptonuria after liver transplantation for hepatitis B-related cirrhosis.**
 Author(s): Kobak AC, Oder G, Kobak S, Argin M, Inal V.
 Source: J Clin Rheumatol.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=16371803&query_hl=13&itool=pubmed_docsum
- **Ocular ochronosis from alkaptonuria.**
 Author(s): Carlson DM, Helgeson MK, Hiatt JA.
 Source: J Am Optom Assoc.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=1813514&query_hl=13&itool=pubmed_docsum
- **Ocular ochronosis in alkaptonuria patients carrying mutations in the homogentisate 1,2-dioxygenase gene.**
 Author(s): Felbor U, Mutsch Y, Grehn F, Muller CR, Kress W.
 Source: The British Journal of Ophthalmology.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10340975&query_hl=13&itool=pubmed_docsum
- **Ocular Ochronosis with Alkaptonuria. Report of a case.**
 Author(s): Himmel S, Adelstein N.
 Source: Ohio State Med J.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=6042337&query_hl=13&itool=pubmed_docsum
- **Picture of the month. Denouement and discussion: alkaptonuria.**
 Author(s): Nitu M, Ancona K, Coleman C, Tunnessen WW Jr.
 Source: Archives of Pediatrics & Adolescent Medicine.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=11529810&query_hl=13&itool=pubmed_docsum
- **Pink napkins--presenting feature in a case of alkaptonuria.**
 Author(s): Sutherland DA, Nicol AD, Williams AJ.
 Source: Journal of Inherited Metabolic Disease.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=6434825&query_hl=13&itool=pubmed_docsum
- **'Pseudohypouricosuria' in alcaptonuria: homogentisic acid interference in the measurement of urinary uric acid with the uricase-peroxidase reaction.**
 Author(s): Moriwaki Y, Yamamoto T, Nasako Y, Ohata H, Takahashi S, Tsutsumi Z, Yamakita J, Higashino K.
 Source: Annals of Clinical Biochemistry.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10456213&query_hl=14&itool=pubmed_docsum

- **Rapid detection methods for five HGO gene mutations causing alkaptonuria.**
 Author(s): Zatkova A, Chmelikova A, Polakova H, Ferakova E, Kadasi L.
 Source: Clinical Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12630963&query_hl=13&itool=pubmed_docsum
- **Reversal of clinical symptoms and radiographic abnormalities with protein restriction and ascorbic acid in alkaptonuria.**
 Author(s): Morava E, Kosztolanyi G, Engelke UF, Wevers RA.
 Source: Annals of Clinical Biochemistry.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12542920&query_hl=13&itool=pubmed_docsum
- **Screening for alkaptonuria in the newborn in Slovakia.**
 Author(s): Srsen S, Varga F.
 Source: Lancet.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=79941&query_hl=13&itool=pubmed_docsum
- **Screening infants for neuroblastoma: discovery of alkaptonuria in one case.**
 Author(s): Mathieu P, Prevel A, David L, Bergeron C, Bory C, Chauvin F, Dia S, Frappaz D, Greffe J, Mathieu M, Philip T.
 Source: Clinica Chimica Acta; International Journal of Clinical Chemistry.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9293385&query_hl=13&itool=pubmed_docsum
- **Sequence analysis of the homogentisate 1,2 dioxygenase gene in a family affected by alkaptonuria.**
 Author(s): Walter K, Gaa A, Schaefer HE.
 Source: Journal of Medical Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10465119&query_hl=13&itool=pubmed_docsum
- **Simple colour reaction for alkaptonuria.**
 Author(s): Valmikinathan K, Verghese N.
 Source: Journal of Clinical Pathology.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5909703&query_hl=13&itool=pubmed_docsum
- **Simultaneous occurrence of alkaptonuria and of Klinefelter syndrome in one patient.**
 Author(s): Srsen S.
 Source: Padiatr Padol.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=859724&query_hl=13&itool=pubmed_docsum

- **Sir Archibald Garrod and alkaptonuria -'story of metabolic genetics'.**
 Author(s): Prasad C, Galbraith PA.
 Source: Clinical Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=16098007&query_hl=13&itool=pubmed_docsum

- **Spontaneous tendon ruptures in alkaptonuria.**
 Author(s): Logani V, Eachempati KK, Malhotra R, Bhan S.
 Source: The Journal of Bone and Joint Surgery. British Volume.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=15446547&query_hl=13&itool=pubmed_docsum

- **Spontaneous tendon ruptures in alkaptonuria.**
 Author(s): Manoj Kumar RV, Rajasekaran S.
 Source: The Journal of Bone and Joint Surgery. British Volume.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12931812&query_hl=13&itool=pubmed_docsum

- **Structural and functional analysis of mutations in alkaptonuria.**
 Author(s): Rodriguez JM, Timm DE, Titus GP, Beltran-Valero De Bernabe D, Criado O, Mueller HA, Rodriguez De Cordoba S, Penalva MA.
 Source: Human Molecular Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=11001939&query_hl=13&itool=pubmed_docsum

- **Tetralogy of Fallot and alkaptonuria in successive generations.**
 Author(s): Haiya PP, Balaji NK, Sukumar IP, Cherian G.
 Source: Indian Pediatrics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=7399662&query_hl=13&itool=pubmed_docsum

- **The black aorta: alkaptonuria diagnosed during coronary artery bypass.**
 Author(s): Paull DE, Picone AC, Baisden CE.
 Source: Southern Medical Journal.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=1948246&query_hl=13&itool=pubmed_docsum

- **The eye in alkaptonuria.**
 Author(s): Wirtschafter JD.
 Source: Birth Defects Orig Artic Ser.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=782595&query_hl=13&itool=pubmed_docsum

- **The human gene for alkaptonuria (AKU) maps to chromosome 3q.**
 Author(s): Janocha S, Wolz W, Srsen S, Srsnova K, Montagutelli X, Guenet JL, Grimm T, Kress W, Muller CR.
 Source: Genomics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=8188241&query_hl=13&itool=pubmed_docsum
- **The incidence of alkaptonuria: a study in chemical individuality. 1902.**
 Author(s): Garrod AE.
 Source: Molecular Medicine (Cambridge, Mass.).
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=8784780&query_hl=13&itool=pubmed_docsum
- **The investigation of aromatic acids in phenylketonuria, alkaptonuria and tyrosinosis using gas-liquid chromatography.**
 Author(s): Hill A, Hoag GN, Zaleski WA.
 Source: Clinica Chimica Acta; International Journal of Clinical Chemistry.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=5022110&query_hl=13&itool=pubmed_docsum
- **The Lancet. The incidence of alkaptonuria: a study in chemical individuality.**
 Author(s): Garrod AE.
 Source: Nutrition Reviews.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=1091883&query_hl=13&itool=pubmed_docsum
- **The molecular basis of alkaptonuria.**
 Author(s): Fernandez-Canon JM, Granadino B, Beltran-Valero de Bernabe D, Renedo M, Fernandez-Ruiz E, Penalva MA, Rodriguez de Cordoba S.
 Source: Nature Genetics.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=8782815&query_hl=13&itool=pubmed_docsum
- **The success of dietary protein restriction in alkaptonuria patients is age-dependent.**
 Author(s): de Haas V, Carbasius Weber EC, de Klerk JB, Bakker HD, Smit GP, Huijbers WA, Duran M, Poll-The BT.
 Source: Journal of Inherited Metabolic Disease.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=9870204&query_hl=13&itool=pubmed_docsum
- **Tyrosine metabolism in cirrhosis: acquired alkaptonuria.**
 Author(s): Nordlinger BM, Fulenwider JT, Faraj BA, Bethel RA, Rudman D.
 Source: Surg Forum.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=401217&query_hl=13&itool=pubmed_docsum

- **Use of nitisinone in patients with alkaptonuria.**

Author(s): Suwannarat P, O'Brien K, Perry MB, Sebring N, Bernardini I, Kaiser-Kupfer MI, Rubin BI, Tsilou E, Gerber LH, Gahl WA.

Source: Metabolism: Clinical and Experimental.

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=15931605&query_hl=13&itool=pubmed_docsum

CHAPTER 2. ALTERNATIVE MEDICINE AND ALKAPTONURIA

Overview

In this chapter, we will begin by introducing you to official information sources on complementary and alternative medicine (CAM) relating to alkaptonuria. At the conclusion of this chapter, we will provide additional sources.

National Center for Complementary and Alternative Medicine

The National Center for Complementary and Alternative Medicine (NCCAM) of the National Institutes of Health (<http://nccam.nih.gov/>) has created a link to the National Library of Medicine's databases to facilitate research for articles that specifically relate to alkaptonuria and complementary medicine. To search the database, go to the following Web site: <http://www.nlm.nih.gov/nccam/camonpubmed.html>. Select **CAM on PubMed**. Enter **alkaptonuria** (or synonyms) into the search box. Click **Go**. The following references provide information on particular aspects of complementary and alternative medicine that are related to alkaptonuria:

- **Homogentisic acid autoxidation and oxygen radical generation: implications for the etiology of alkaptonuric arthritis.**
Author(s): Martin JP Jr, Batkoff B.
Source: Free Radical Biology & Medicine.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=3121448&query_hl=1&itool=pubmed_docsum

Additional Web Resources

A number of additional Web sites offer encyclopedic information covering CAM and related topics. The following is a representative sample:

- Alternative Medicine Foundation, Inc.: <http://www.herbmed.org/>
- AOL: <http://health.aol.com/healthyliving/althealth>

- Chinese Medicine: <http://www.newcenturynutrition.com/>
- drkoop.com[®]: <http://www.drkoop.com/naturalmedicine.html>
- Family Village: http://www.familyvillage.wisc.edu/med_altn.htm
- Google: <http://directory.google.com/Top/Health/Alternative/>
- Healthnotes: <http://www.healthnotes.com/>
- Open Directory Project: <http://dmoz.org/Health/Alternative/>
- Yahoo.com: http://dir.yahoo.com/Health/Alternative_Medicine/

General References

A good place to find general background information on CAM is the National Library of Medicine. It has prepared within the MEDLINEplus system an information topic page dedicated to complementary and alternative medicine. To access this page, go to the MEDLINEplus site at <http://www.nlm.nih.gov/medlineplus/alternativemedicine.html>. This Web site provides a general overview of various topics and can lead to a number of general sources.

APPENDICES

APPENDIX A. HELP ME UNDERSTAND GENETICS

Overview

This appendix presents basic information about genetics in clear language and provides links to online resources.⁷

The Basics: Genes and How They Work

This section gives you information on the basics of cells, DNA, genes, chromosomes, and proteins.

What Is a Cell?

Cells are the basic building blocks of all living things. The human body is composed of trillions of cells. They provide structure for the body, take in nutrients from food, convert those nutrients into energy, and carry out specialized functions. Cells also contain the body's hereditary material and can make copies of themselves.

Cells have many parts, each with a different function. Some of these parts, called organelles, are specialized structures that perform certain tasks within the cell. Human cells contain the following major parts, listed in alphabetical order:

- **Cytoplasm:** The cytoplasm is fluid inside the cell that surrounds the organelles.
- **Endoplasmic reticulum (ER):** This organelle helps process molecules created by the cell and transport them to their specific destinations either inside or outside the cell.
- **Golgi apparatus:** The golgi apparatus packages molecules processed by the endoplasmic reticulum to be transported out of the cell.
- **Lysosomes and peroxisomes:** These organelles are the recycling center of the cell. They digest foreign bacteria that invade the cell, rid the cell of toxic substances, and recycle worn-out cell components.

⁷ This appendix is an excerpt from the National Library of Medicine's handbook, *Help Me Understand Genetics*. For the full text of the *Help Me Understand Genetics* handbook, see <http://ghr.nlm.nih.gov/handbook>.

- **Mitochondria:** Mitochondria are complex organelles that convert energy from food into a form that the cell can use. They have their own genetic material, separate from the DNA in the nucleus, and can make copies of themselves.
- **Nucleus:** The nucleus serves as the cell's command center, sending directions to the cell to grow, mature, divide, or die. It also houses DNA (deoxyribonucleic acid), the cell's hereditary material. The nucleus is surrounded by a membrane called the nuclear envelope, which protects the DNA and separates the nucleus from the rest of the cell.
- **Plasma membrane:** The plasma membrane is the outer lining of the cell. It separates the cell from its environment and allows materials to enter and leave the cell.
- **Ribosomes:** Ribosomes are organelles that process the cell's genetic instructions to create proteins. These organelles can float freely in the cytoplasm or be connected to the endoplasmic reticulum.


What Is DNA?

DNA, or deoxyribonucleic acid, is the hereditary material in humans and almost all other organisms. Nearly every cell in a person's body has the same DNA. Most DNA is located in the cell nucleus (where it is called nuclear DNA), but a small amount of DNA can also be found in the mitochondria (where it is called mitochondrial DNA or mtDNA).

The information in DNA is stored as a code made up of four chemical bases: adenine (A), guanine (G), cytosine (C), and thymine (T). Human DNA consists of about 3 billion bases, and more than 99 percent of those bases are the same in all people. The order, or sequence, of these bases determines the information available for building and maintaining an organism, similar to the way in which letters of the alphabet appear in a certain order to form words and sentences.

DNA bases pair up with each other, A with T and C with G, to form units called base pairs. Each base is also attached to a sugar molecule and a phosphate molecule. Together, a base, sugar, and phosphate are called a nucleotide. Nucleotides are arranged in two long strands that form a spiral called a double helix. The structure of the double helix is somewhat like a ladder, with the base pairs forming the ladder's rungs and the sugar and phosphate molecules forming the vertical sidepieces of the ladder.

An important property of DNA is that it can replicate, or make copies of itself. Each strand of DNA in the double helix can serve as a pattern for duplicating the sequence of bases. This is critical when cells divide because each new cell needs to have an exact copy of the DNA present in the old cell.



U.S. National Library of Medicine

DNA is a double helix formed by base pairs attached to a sugar-phosphate backbone.

What Is Mitochondrial DNA?

Although most DNA is packaged in chromosomes within the nucleus, mitochondria also have a small amount of their own DNA. This genetic material is known as mitochondrial DNA or mtDNA.

Mitochondria are structures within cells that convert the energy from food into a form that cells can use. Each cell contains hundreds to thousands of mitochondria, which are located in the fluid that surrounds the nucleus (the cytoplasm).

Mitochondria produce energy through a process called oxidative phosphorylation. This process uses oxygen and simple sugars to create adenosine triphosphate (ATP), the cell's main energy source. A set of enzyme complexes, designated as complexes I-V, carry out oxidative phosphorylation within mitochondria.

In addition to energy production, mitochondria play a role in several other cellular activities. For example, mitochondria help regulate the self-destruction of cells (apoptosis). They are also necessary for the production of substances such as cholesterol and heme (a component of hemoglobin, the molecule that carries oxygen in the blood).


Mitochondrial DNA contains 37 genes, all of which are essential for normal mitochondrial function. Thirteen of these genes provide instructions for making enzymes involved in oxidative phosphorylation. The remaining genes provide instructions for making molecules called transfer RNAs (tRNAs) and ribosomal RNAs (rRNAs), which are chemical cousins of

DNA. These types of RNA help assemble protein building blocks (amino acids) into functioning proteins.

What Is a Gene?

A gene is the basic physical and functional unit of heredity. Genes, which are made up of DNA, act as instructions to make molecules called proteins. In humans, genes vary in size from a few hundred DNA bases to more than 2 million bases. The Human Genome Project has estimated that humans have between 20,000 and 25,000 genes.

Every person has two copies of each gene, one inherited from each parent. Most genes are the same in all people, but a small number of genes (less than 1 percent of the total) are slightly different between people. Alleles are forms of the same gene with small differences in their sequence of DNA bases. These small differences contribute to each person's unique physical features.




Genes are made up of DNA. Each chromosome contains many genes.

What Is a Chromosome?

In the nucleus of each cell, the DNA molecule is packaged into thread-like structures called chromosomes. Each chromosome is made up of DNA tightly coiled many times around proteins called histones that support its structure.

Chromosomes are not visible in the cell's nucleus—not even under a microscope—when the cell is not dividing. However, the DNA that makes up chromosomes becomes more tightly packed during cell division and is then visible under a microscope. Most of what researchers know about chromosomes was learned by observing chromosomes during cell division.


Each chromosome has a constriction point called the centromere, which divides the chromosome into two sections, or "arms." The short arm of the chromosome is labeled the "p arm." The long arm of the chromosome is labeled the "q arm." The location of the centromere on each chromosome gives the chromosome its characteristic shape, and can be used to help describe the location of specific genes.



DNA and histone proteins are packaged into structures called chromosomes.

How Many Chromosomes Do People Have?

In humans, each cell normally contains 23 pairs of chromosomes, for a total of 46. Twenty-two of these pairs, called autosomes, look the same in both males and females. The 23rd pair, the sex chromosomes, differ between males and females. Females have two copies of the X chromosome, while males have one X and one Y chromosome.



U.S. National Library of Medicine

The 22 autosomes are numbered by size.

The other two chromosomes, X and Y, are the sex chromosomes.

This picture of the human chromosomes lined up in pairs is called a karyotype.

How Do Geneticists Indicate the Location of a Gene?

Geneticists use maps to describe the location of a particular gene on a chromosome. One type of map uses the cytogenetic location to describe a gene's position. The cytogenetic location is based on a distinctive pattern of bands created when chromosomes are stained with certain chemicals. Another type of map uses the molecular location, a precise description of a gene's position on a chromosome. The molecular location is based on the sequence of DNA building blocks (base pairs) that make up the chromosome.

Cytogenetic Location

Geneticists use a standardized way of describing a gene's cytogenetic location. In most cases, the location describes the position of a particular band on a stained chromosome:

17q12

It can also be written as a range of bands, if less is known about the exact location:


17q12-q21

The combination of numbers and letters provide a gene's "address" on a chromosome. This address is made up of several parts:

- The chromosome on which the gene can be found. The first number or letter used to describe a gene's location represents the chromosome. Chromosomes 1 through 22 (the autosomes) are designated by their chromosome number. The sex chromosomes are designated by X or Y.

- The arm of the chromosome. Each chromosome is divided into two sections (arms) based on the location of a narrowing (constriction) called the centromere. By convention, the shorter arm is called p, and the longer arm is called q. The chromosome arm is the second part of the gene's address. For example, 5q is the long arm of chromosome 5, and Xp is the short arm of the X chromosome.
- The position of the gene on the p or q arm. The position of a gene is based on a distinctive pattern of light and dark bands that appear when the chromosome is stained in a certain way. The position is usually designated by two digits (representing a region and a band), which are sometimes followed by a decimal point and one or more additional digits (representing sub-bands within a light or dark area). The number indicating the gene position increases with distance from the centromere. For example: 14q21 represents position 21 on the long arm of chromosome 14. 14q21 is closer to the centromere than 14q22.

Sometimes, the abbreviations "cen" or "ter" are also used to describe a gene's cytogenetic location. "Cen" indicates that the gene is very close to the centromere. For example, 16pcen refers to the short arm of chromosome 16 near the centromere. "Ter" stands for terminus, which indicates that the gene is very close to the end of the p or q arm. For example, 14qter refers to the tip of the long arm of chromosome 14. ("Tel" is also sometimes used to describe a gene's location. "Tel" stands for telomeres, which are at the ends of each chromosome. The abbreviations "tel" and "ter" refer to the same location.)



The CFTR gene is located on the long arm of chromosome 7 at position 7q31.2.

Molecular Location

The Human Genome Project, an international research effort completed in 2003, determined the sequence of base pairs for each human chromosome. This sequence information allows researchers to provide a more specific address than the cytogenetic location for many genes. A gene's molecular address pinpoints the location of that gene in terms of base pairs. For example, the molecular location of the APOE gene on chromosome 19 begins with base pair 50,100,901 and ends with base pair 50,104,488. This range describes the gene's precise position on chromosome 19 and indicates the size of the gene (3,588 base pairs). Knowing a gene's molecular location also allows researchers to determine exactly how far the gene is from other genes on the same chromosome.

Different groups of researchers often present slightly different values for a gene's molecular location. Researchers interpret the sequence of the human genome using a variety of methods, which can result in small differences in a gene's molecular address. For example, the National Center for Biotechnology Information (NCBI) identifies the molecular location of the APOE gene as base pair 50,100,901 to base pair 50,104,488 on chromosome 19. The Ensembl database identifies the location of this gene as base pair 50,100,879 to base pair 50,104,489 on chromosome 19. Neither of these addresses is incorrect; they represent different interpretations of the same data. For consistency, Genetics Home Reference presents data from NCBI for the molecular location of genes.

What Are Proteins and What Do They Do?

Proteins are large, complex molecules that play many critical roles in the body. They do most of the work in cells and are required for the structure, function, and regulation of the body's tissues and organs.

Proteins are made up of hundreds or thousands of smaller units called amino acids, which are attached to one another in long chains. There are 20 different types of amino acids that can be combined to make a protein. The sequence of amino acids determines each protein's unique 3-dimensional structure and its specific function.

Examples of Protein Functions

Proteins can be described according to their large range of functions in the body, listed in alphabetical order:

Function	Description	Example
Antibody	Antibodies bind to specific foreign particles, such as viruses and bacteria, to help protect the body.	Immunoglobulin G (IgG)
Enzyme	Enzymes carry out almost all of the thousands of chemical reactions that take place in cells. They also assist with the formation of new molecules by reading the genetic information stored in DNA.	Phenylalanine hydroxylase
Messenger	Messenger proteins, such as some types of hormones, transmit signals to coordinate biological processes between different cells, tissues, and organs.	Growth hormone
Structural component	These proteins provide structure and support for cells. On a larger scale, they also allow the body to move.	Actin
Transport/storage	These proteins bind and carry atoms and small molecules within cells and throughout the body.	Ferritin

How Does a Gene Make a Protein?


Most genes contain the information needed to make functional molecules called proteins. (A few genes produce other molecules that help the cell assemble proteins.) The journey from gene to protein is complex and tightly controlled within each cell. It consists of two major steps: transcription and translation. Together, transcription and translation are known as gene expression.

During the process of transcription, the information stored in a gene's DNA is transferred to a similar molecule called RNA (ribonucleic acid) in the cell nucleus. Both RNA and DNA are made up of a chain of nucleotide bases, but they have slightly different chemical properties. The type of RNA that contains the information for making a protein is called messenger RNA (mRNA) because it carries the information, or message, from the DNA out of the nucleus into the cytoplasm.

Translation, the second step in getting from a gene to a protein, takes place in the cytoplasm. The mRNA interacts with a specialized complex called a ribosome, which "reads" the sequence of mRNA bases. Each sequence of three bases, called a codon, usually codes for

one particular amino acid. (Amino acids are the building blocks of proteins.) A type of RNA called transfer RNA (tRNA) assembles the protein, one amino acid at a time. Protein assembly continues until the ribosome encounters a “stop” codon (a sequence of three bases that does not code for an amino acid).

The flow of information from DNA to RNA to proteins is one of the fundamental principles of molecular biology. It is so important that it is sometimes called the “central dogma.”



Through the processes of transcription and translation, information from genes is used to make proteins.

Can Genes Be Turned On and Off in Cells?

Each cell expresses, or turns on, only a fraction of its genes. The rest of the genes are repressed, or turned off. The process of turning genes on and off is known as gene regulation. Gene regulation is an important part of normal development. Genes are turned on and off in different patterns during development to make a brain cell look and act different from a liver cell or a muscle cell, for example. Gene regulation also allows cells to react quickly to changes in their environments. Although we know that the regulation of genes is critical for life, this complex process is not yet fully understood.


Gene regulation can occur at any point during gene expression, but most commonly occurs at the level of transcription (when the information in a gene’s DNA is transferred to mRNA). Signals from the environment or from other cells activate proteins called transcription factors. These proteins bind to regulatory regions of a gene and increase or decrease the level of transcription. By controlling the level of transcription, this process can determine the amount of protein product that is made by a gene at any given time.

How Do Cells Divide?

There are two types of cell division: mitosis and meiosis. Most of the time when people refer to “cell division,” they mean mitosis, the process of making new body cells. Meiosis is the type of cell division that creates egg and sperm cells.

Mitosis is a fundamental process for life. During mitosis, a cell duplicates all of its contents, including its chromosomes, and splits to form two identical daughter cells. Because this process is so critical, the steps of mitosis are carefully controlled by a number of genes. When mitosis is not regulated correctly, health problems such as cancer can result.

The other type of cell division, meiosis, ensures that humans have the same number of chromosomes in each generation. It is a two-step process that reduces the chromosome number by half—from 46 to 23—to form sperm and egg cells. When the sperm and egg cells unite at conception, each contributes 23 chromosomes so the resulting embryo will have the usual 46. Meiosis also allows genetic variation through a process of DNA shuffling while the cells are dividing.



Mitosis and meiosis, the two types of cell division.

How Do Genes Control the Growth and Division of Cells?

A variety of genes are involved in the control of cell growth and division. The cell cycle is the cell’s way of replicating itself in an organized, step-by-step fashion. Tight regulation of this process ensures that a dividing cell’s DNA is copied properly, any errors in the DNA are repaired, and each daughter cell receives a full set of chromosomes. The cycle has checkpoints (also called restriction points), which allow certain genes to check for mistakes and halt the cycle for repairs if something goes wrong.

If a cell has an error in its DNA that cannot be repaired, it may undergo programmed cell death (apoptosis). Apoptosis is a common process throughout life that helps the body get rid of cells it doesn't need. Cells that undergo apoptosis break apart and are recycled by a type of white blood cell called a macrophage. Apoptosis protects the body by removing genetically damaged cells that could lead to cancer, and it plays an important role in the development of the embryo and the maintenance of adult tissues.

Cancer results from a disruption of the normal regulation of the cell cycle. When the cycle proceeds without control, cells can divide without order and accumulate genetic defects that can lead to a cancerous tumor.

Genetic Mutations and Health

This section presents basic information about gene mutations, chromosomal changes, and conditions that run in families.⁸

What Is a Gene Mutation and How Do Mutations Occur?

A gene mutation is a permanent change in the DNA sequence that makes up a gene. Mutations range in size from a single DNA building block (DNA base) to a large segment of a chromosome.

Gene mutations occur in two ways: they can be inherited from a parent or acquired during a person's lifetime. Mutations that are passed from parent to child are called hereditary mutations or germline mutations (because they are present in the egg and sperm cells, which are also called germ cells). This type of mutation is present throughout a person's life in virtually every cell in the body.

Mutations that occur only in an egg or sperm cell, or those that occur just after fertilization, are called new (de novo) mutations. De novo mutations may explain genetic disorders in which an affected child has a mutation in every cell, but has no family history of the disorder.

Acquired (or somatic) mutations occur in the DNA of individual cells at some time during a person's life. These changes can be caused by environmental factors such as ultraviolet radiation from the sun, or can occur if a mistake is made as DNA copies itself during cell division. Acquired mutations in somatic cells (cells other than sperm and egg cells) cannot be passed on to the next generation.

Mutations may also occur in a single cell within an early embryo. As all the cells divide during growth and development, the individual will have some cells with the mutation and some cells without the genetic change. This situation is called mosaicism.

Some genetic changes are very rare; others are common in the population. Genetic changes that occur in more than 1 percent of the population are called polymorphisms. They are common enough to be considered a normal variation in the DNA. Polymorphisms are

⁸ This section has been adapted from the National Library of Medicine's handbook, *Help Me Understand Genetics*, which presents basic information about genetics in clear language and provides links to online resources: <http://ghr.nlm.nih.gov/handbook>.

responsible for many of the normal differences between people such as eye color, hair color, and blood type. Although many polymorphisms have no negative effects on a person's health, some of these variations may influence the risk of developing certain disorders.

How Can Gene Mutations Affect Health and Development?

To function correctly, each cell depends on thousands of proteins to do their jobs in the right places at the right times. Sometimes, gene mutations prevent one or more of these proteins from working properly. By changing a gene's instructions for making a protein, a mutation can cause the protein to malfunction or to be missing entirely. When a mutation alters a protein that plays a critical role in the body, it can disrupt normal development or cause a medical condition. A condition caused by mutations in one or more genes is called a genetic disorder.

In some cases, gene mutations are so severe that they prevent an embryo from surviving until birth. These changes occur in genes that are essential for development, and often disrupt the development of an embryo in its earliest stages. Because these mutations have very serious effects, they are incompatible with life.

It is important to note that genes themselves do not cause disease—genetic disorders are caused by mutations that make a gene function improperly. For example, when people say that someone has “the cystic fibrosis gene,” they are usually referring to a mutated version of the CFTR gene, which causes the disease. All people, including those without cystic fibrosis, have a version of the CFTR gene.

Do All Gene Mutations Affect Health and Development?

No, only a small percentage of mutations cause genetic disorders—most have no impact on health or development. For example, some mutations alter a gene's DNA base sequence but do not change the function of the protein made by the gene.

Often, gene mutations that could cause a genetic disorder are repaired by certain enzymes before the gene is expressed (makes a protein). Each cell has a number of pathways through which enzymes recognize and repair mistakes in DNA. Because DNA can be damaged or mutated in many ways, DNA repair is an important process by which the body protects itself from disease.

A very small percentage of all mutations actually have a positive effect. These mutations lead to new versions of proteins that help an organism and its future generations better adapt to changes in their environment. For example, a beneficial mutation could result in a protein that protects the organism from a new strain of bacteria.

For More Information about DNA Repair and the Health Effects of Gene Mutations

- The University of Utah Genetic Science Learning Center provides information about genetic disorders that explains why some mutations cause disorders but others do not. (Refer to the questions in the far right column.)
See <http://learn.genetics.utah.edu/units/disorders/whataregd/>.

- Additional information about DNA repair is available from the NCBI Science Primer. In the chapter called “What Is A Cell?”, scroll down to the heading “DNA Repair Mechanisms.” See http://www.ncbi.nlm.nih.gov/About/primer/genetics_cell.html.

What Kinds of Gene Mutations Are Possible?

The DNA sequence of a gene can be altered in a number of ways. Gene mutations have varying effects on health, depending on where they occur and whether they alter the function of essential proteins. The types of mutations include:

- **Missense mutation:** This type of mutation is a change in one DNA base pair that results in the substitution of one amino acid for another in the protein made by a gene.
- **Nonsense mutation:** A nonsense mutation is also a change in one DNA base pair. Instead of substituting one amino acid for another, however, the altered DNA sequence prematurely signals the cell to stop building a protein. This type of mutation results in a shortened protein that may function improperly or not at all.
- **Insertion:** An insertion changes the number of DNA bases in a gene by adding a piece of DNA. As a result, the protein made by the gene may not function properly.
- **Deletion:** A deletion changes the number of DNA bases by removing a piece of DNA. Small deletions may remove one or a few base pairs within a gene, while larger deletions can remove an entire gene or several neighboring genes. The deleted DNA may alter the function of the resulting protein(s).
- **Duplication:** A duplication consists of a piece of DNA that is abnormally copied one or more times. This type of mutation may alter the function of the resulting protein.
- **Frameshift mutation:** This type of mutation occurs when the addition or loss of DNA bases changes a gene’s reading frame. A reading frame consists of groups of 3 bases that each code for one amino acid. A frameshift mutation shifts the grouping of these bases and changes the code for amino acids. The resulting protein is usually nonfunctional. Insertions, deletions, and duplications can all be frameshift mutations.
- **Repeat expansion:** Nucleotide repeats are short DNA sequences that are repeated a number of times in a row. For example, a trinucleotide repeat is made up of 3-base-pair sequences, and a tetranucleotide repeat is made up of 4-base-pair sequences. A repeat expansion is a mutation that increases the number of times that the short DNA sequence is repeated. This type of mutation can cause the resulting protein to function improperly.

Can Changes in Chromosomes Affect Health and Development?

Changes that affect entire chromosomes or segments of chromosomes can cause problems with growth, development, and function of the body’s systems. These changes can affect many genes along the chromosome and alter the proteins made by those genes. Conditions caused by a change in the number or structure of chromosomes are known as chromosomal disorders.

Human cells normally contain 23 pairs of chromosomes, for a total of 46 chromosomes in each cell. A change in the number of chromosomes leads to a chromosomal disorder. These

changes can occur during the formation of reproductive cells (eggs and sperm) or in early fetal development. A gain or loss of chromosomes from the normal 46 is called aneuploidy. The most common form of aneuploidy is trisomy, or the presence of an extra chromosome in each cell. “Tri-” is Greek for “three”; people with trisomy have three copies of a particular chromosome in each cell instead of the normal two copies. Down syndrome is an example of a condition caused by trisomy – people with Down syndrome typically have three copies of chromosome 21 in each cell, for a total of 47 chromosomes per cell.

Monosomy, or the loss of one chromosome from each cell, is another kind of aneuploidy. “Mono-” is Greek for “one”; people with monosomy have one copy of a particular chromosome in each cell instead of the normal two copies. Turner syndrome is a condition caused by monosomy. Women with Turner syndrome are often missing one copy of the X chromosome in every cell, for a total of 45 chromosomes per cell.

Chromosomal disorders can also be caused by changes in chromosome structure. These changes are caused by the breakage and reunion of chromosome segments when an egg or sperm cell is formed or in early fetal development. Pieces of DNA can be rearranged within one chromosome, or transferred between two or more chromosomes. The effects of structural changes depend on their size and location. Many different structural changes are possible; some cause medical problems, while others may have no effect on a person’s health.

Many cancer cells also have changes in their chromosome number or structure. These changes most often occur in somatic cells (cells other than eggs and sperm) during a person’s lifetime.

Can Changes in Mitochondrial DNA Affect Health and Development?

Mitochondria are structures within cells that convert the energy from food into a form that cells can use. Although most DNA is packaged in chromosomes within the nucleus, mitochondria also have a small amount of their own DNA (known as mitochondrial DNA or mtDNA). In some cases, inherited changes in mitochondrial DNA can cause problems with growth, development, and function of the body’s systems. These mutations disrupt the mitochondria’s ability to generate energy efficiently for the cell.

Conditions caused by mutations in mitochondrial DNA often involve multiple organ systems. The effects of these conditions are most pronounced in organs and tissues that require a lot of energy (such as the heart, brain, and muscles). Although the health consequences of inherited mitochondrial DNA mutations vary widely, frequently observed features include muscle weakness and wasting, problems with movement, diabetes, kidney failure, heart disease, loss of intellectual functions (dementia), hearing loss, and abnormalities involving the eyes and vision.

Mitochondrial DNA is also prone to noninherited (somatic) mutations. Somatic mutations occur in the DNA of certain cells during a person’s lifetime, and typically are not passed to future generations. Because mitochondrial DNA has a limited ability to repair itself when it is damaged, these mutations tend to build up over time. A buildup of somatic mutations in mitochondrial DNA has been associated with some forms of cancer and an increased risk of certain age-related disorders such as heart disease, Alzheimer disease, and Parkinson disease. Additionally, research suggests that the progressive accumulation of these mutations over a person’s lifetime may play a role in the normal process of aging.

What Are Complex or Multifactorial Disorders?

Researchers are learning that nearly all conditions and diseases have a genetic component. Some disorders, such as sickle cell anemia and cystic fibrosis, are caused by mutations in a single gene. The causes of many other disorders, however, are much more complex. Common medical problems such as heart disease, diabetes, and obesity do not have a single genetic cause—they are likely associated with the effects of multiple genes in combination with lifestyle and environmental factors. Conditions caused by many contributing factors are called complex or multifactorial disorders.

Although complex disorders often cluster in families, they do not have a clear-cut pattern of inheritance. This makes it difficult to determine a person's risk of inheriting or passing on these disorders. Complex disorders are also difficult to study and treat because the specific factors that cause most of these disorders have not yet been identified. By 2010, however, researchers predict they will have found the major contributing genes for many common complex disorders.

What Information about a Genetic Condition Can Statistics Provide?

Statistical data can provide general information about how common a condition is, how many people have the condition, or how likely it is that a person will develop the condition. Statistics are not personalized, however—they offer estimates based on groups of people. By taking into account a person's family history, medical history, and other factors, a genetics professional can help interpret what statistics mean for a particular patient.

Common Statistical Terms

Some statistical terms are commonly used when describing genetic conditions and other disorders. These terms include:

Statistical Term	Description	Examples
<i>Incidence</i>	The incidence of a gene mutation or a genetic disorder is the number of people who are born with the mutation or disorder in a specified group per year. Incidence is often written in the form "1 in [a number]" or as a total number of live births.	About 1 in 200,000 people in the United States are born with syndrome A each year. An estimated 15,000 infants with syndrome B were born last year worldwide.

<i>Prevalence</i>	The prevalence of a gene mutation or a genetic disorder is the total number of people in a specified group at a given time who have the mutation or disorder. This term includes both newly diagnosed and pre-existing cases in people of any age. Prevalence is often written in the form "1 in [a number]" or as a total number of people who have a condition.	Approximately 1 in 100,000 people in the United States have syndrome A at the present time. About 100,000 children worldwide currently have syndrome B.
<i>Mortality</i>	Mortality is the number of deaths from a particular disorder occurring in a specified group per year. Mortality is usually expressed as a total number of deaths.	An estimated 12,000 people worldwide died from syndrome C in 2002.
<i>Lifetime risk</i>	Lifetime risk is the average risk of developing a particular disorder at some point during a lifetime. Lifetime risk is often written as a percentage or as "1 in [a number]." It is important to remember that the risk per year or per decade is much lower than the lifetime risk. In addition, other factors may increase or decrease a person's risk as compared with the average.	Approximately 1 percent of people in the United States develop disorder D during their lifetimes. The lifetime risk of developing disorder D is 1 in 100.

Naming Genetic Conditions

Genetic conditions are not named in one standard way (unlike genes, which are given an official name and symbol by a formal committee). Doctors who treat families with a particular disorder are often the first to propose a name for the condition. Expert working groups may later revise the name to improve its usefulness. Naming is important because it allows accurate and effective communication about particular conditions, which will ultimately help researchers find new approaches to treatment.

Disorder names are often derived from one or a combination of sources:

- The basic genetic or biochemical defect that causes the condition (for example, alpha-1 antitrypsin deficiency)
- One or more major signs or symptoms of the disorder (for example, sickle cell anemia)
- The parts of the body affected by the condition (for example, retinoblastoma)
- The name of a physician or researcher, often the first person to describe the disorder (for example, Marfan syndrome, which was named after Dr. Antoine Bernard-Jean Marfan)

- A geographic area (for example, familial Mediterranean fever, which occurs mainly in populations bordering the Mediterranean Sea)
- The name of a patient or family with the condition (for example, amyotrophic lateral sclerosis, which is also called Lou Gehrig disease after a famous baseball player who had the condition).

Disorders named after a specific person or place are called eponyms. There is debate as to whether the possessive form (e.g., Alzheimer's disease) or the nonpossessive form (Alzheimer disease) of eponyms is preferred. As a rule, medical geneticists use the nonpossessive form, and this form may become the standard for doctors in all fields of medicine. Genetics Home Reference uses the nonpossessive form of eponyms.

Genetics Home Reference consults with experts in the field of medical genetics to provide the current, most accurate name for each disorder. Alternate names are included as synonyms.

Naming genes

The HUGO Gene Nomenclature Committee (HGNC) designates an official name and symbol (an abbreviation of the name) for each known human gene. Some official gene names include additional information in parentheses, such as related genetic conditions, subtypes of a condition, or inheritance pattern. The HGNC is a non-profit organization funded by the U.K. Medical Research Council and the U.S. National Institutes of Health. The Committee has named more than 13,000 of the estimated 20,000 to 25,000 genes in the human genome.

During the research process, genes often acquire several alternate names and symbols. Different researchers investigating the same gene may each give the gene a different name, which can cause confusion. The HGNC assigns a unique name and symbol to each human gene, which allows effective organization of genes in large databanks, aiding the advancement of research. For specific information about how genes are named, refer to the HGNC's Guidelines for Human Gene Nomenclature.

Genetics Home Reference describes genes using the HGNC's official gene names and gene symbols. Genetics Home Reference frequently presents the symbol and name separated with a colon (for example, FGFR4: Fibroblast growth factor receptor 4).


Inheriting Genetic Conditions

This section gives you information on inheritance patterns and understanding risk.

What Does It Mean If a Disorder Seems to Run in My Family?

A particular disorder might be described as "running in a family" if more than one person in the family has the condition. Some disorders that affect multiple family members are caused by gene mutations, which can be inherited (passed down from parent to child). Other conditions that appear to run in families are not inherited. Instead, environmental factors such as dietary habits or a combination of genetic and environmental factors are responsible for these disorders.

It is not always easy to determine whether a condition in a family is inherited. A genetics professional can use a person's family history (a record of health information about a person's immediate and extended family) to help determine whether a disorder has a genetic component.



U.S. National Library of Medicine

Some disorders are seen in more than one generation of a family.

Why Is It Important to Know My Family Medical History?

A family medical history is a record of health information about a person and his or her close relatives. A complete record includes information from three generations of relatives, including children, brothers and sisters, parents, aunts and uncles, nieces and nephews, grandparents, and cousins.

Families have many factors in common, including their genes, environment, and lifestyle. Together, these factors can give clues to medical conditions that may run in a family. By noticing patterns of disorders among relatives, healthcare professionals can determine whether an individual, other family members, or future generations may be at an increased risk of developing a particular condition.

A family medical history can identify people with a higher-than-usual chance of having common disorders, such as heart disease, high blood pressure, stroke, certain cancers, and diabetes. These complex disorders are influenced by a combination of genetic factors, environmental conditions, and lifestyle choices. A family history also can provide information about the risk of rarer conditions caused by mutations in a single gene, such as cystic fibrosis and sickle cell anemia.

While a family medical history provides information about the risk of specific health concerns, having relatives with a medical condition does not mean that an individual will definitely develop that condition. On the other hand, a person with no family history of a disorder may still be at risk of developing that disorder.

Knowing one's family medical history allows a person to take steps to reduce his or her risk. For people at an increased risk of certain cancers, healthcare professionals may recommend more frequent screening (such as mammography or colonoscopy) starting at an earlier age. Healthcare providers may also encourage regular checkups or testing for people with a medical condition that runs in their family. Additionally, lifestyle changes such as adopting a healthier diet, getting regular exercise, and quitting smoking help many people lower their chances of developing heart disease and other common illnesses.

The easiest way to get information about family medical history is to talk to relatives about their health. Have they had any medical problems, and when did they occur? A family gathering could be a good time to discuss these issues. Additionally, obtaining medical records and other documents (such as obituaries and death certificates) can help complete a family medical history. It is important to keep this information up-to-date and to share it with a healthcare professional regularly.

What Are the Different Ways in which a Genetic Condition Can Be Inherited?

Some genetic conditions are caused by mutations in a single gene. These conditions are usually inherited in one of several straightforward patterns, depending on the gene involved:

Inheritance Pattern	Description	Examples
Autosomal dominant	One mutated copy of the gene in each cell is sufficient for a person to be affected by an autosomal dominant disorder. Each affected person usually has one affected parent. Autosomal dominant disorders tend to occur in every generation of an affected family.	Huntington disease, neurofibromatosis type 1

Autosomal recessive	Two mutated copies of the gene are present in each cell when a person has an autosomal recessive disorder. An affected person usually has unaffected parents who each carry a single copy of the mutated gene (and are referred to as carriers). Autosomal recessive disorders are typically not seen in every generation of an affected family.	cystic fibrosis, sickle cell anemia
X-linked dominant	X-linked dominant disorders are caused by mutations in genes on the X chromosome. Females are more frequently affected than males, and the chance of passing on an X-linked dominant disorder differs between men and women. Families with an X-linked dominant disorder often have both affected males and affected females in each generation. A striking characteristic of X-linked inheritance is that fathers cannot pass X-linked traits to their sons (no male-to-male transmission).	fragile X syndrome
X-linked recessive	X-linked recessive disorders are also caused by mutations in genes on the X chromosome. Males are more frequently affected than females, and the chance of passing on the disorder differs between men and women. Families with an X-linked recessive disorder often have affected males, but rarely affected females, in each generation. A striking characteristic of X-linked inheritance is that fathers cannot pass X-linked traits to their sons (no male-to-male transmission).	hemophilia, Fabry disease
Codominant	In codominant inheritance, two different versions (alleles) of a gene can be expressed, and each version makes a slightly different protein. Both alleles influence the genetic trait or determine the characteristics of the genetic condition.	ABO blood group, alpha-1 antitrypsin deficiency
Mitochondrial	This type of inheritance, also known as maternal inheritance, applies to genes in mitochondrial DNA. Mitochondria, which are structures in each cell that convert molecules into energy, each contain a small amount of DNA. Because only egg cells contribute mitochondria to the developing embryo, only females can pass on mitochondrial conditions to their children. Mitochondrial disorders can appear in every generation of a family and can affect both males and females, but fathers do not pass mitochondrial traits to their children.	Leber hereditary optic neuropathy (LHON)

Many other disorders are caused by a combination of the effects of multiple genes or by interactions between genes and the environment. Such disorders are more difficult to analyze because their genetic causes are often unclear, and they do not follow the patterns of inheritance described above. Examples of conditions caused by multiple genes or gene/environment interactions include heart disease, diabetes, schizophrenia, and certain types of cancer. Disorders caused by changes in the number or structure of chromosomes do not follow the straightforward patterns of inheritance listed above. Other genetic factors can also influence how a disorder is inherited.

If a Genetic Disorder Runs in My Family, What Are the Chances That My Children Will Have the Condition?

When a genetic disorder is diagnosed in a family, family members often want to know the likelihood that they or their children will develop the condition. This can be difficult to predict in some cases because many factors influence a person's chances of developing a genetic condition. One important factor is how the condition is inherited. For example:

- **Autosomal dominant inheritance:** A person affected by an autosomal dominant disorder has a 50 percent chance of passing the mutated gene to each child. The chance that a child will not inherit the mutated gene is also 50 percent.
- **Autosomal recessive inheritance:** Two unaffected people who each carry one copy of the mutated gene for an autosomal recessive disorder (carriers) have a 25 percent chance with each pregnancy of having a child affected by the disorder. The chance with each pregnancy of having an unaffected child who is a carrier of the disorder is 50 percent, and the chance that a child will not have the disorder and will not be a carrier is 25 percent.
- **X-linked dominant inheritance:** The chance of passing on an X-linked dominant condition differs between men and women because men have one X chromosome and one Y chromosome, while women have two X chromosomes. A man passes on his Y chromosome to all of his sons and his X chromosome to all of his daughters. Therefore, the sons of a man with an X-linked dominant disorder will not be affected, but all of his daughters will inherit the condition. A woman passes on one or the other of her X chromosomes to each child. Therefore, a woman with an X-linked dominant disorder has a 50 percent chance of having an affected daughter or son with each pregnancy.
- **X-linked recessive inheritance:** Because of the difference in sex chromosomes, the probability of passing on an X-linked recessive disorder also differs between men and women. The sons of a man with an X-linked recessive disorder will not be affected, and his daughters will carry one copy of the mutated gene. With each pregnancy, a woman who carries an X-linked recessive disorder has a 50 percent chance of having sons who are affected and a 50 percent chance of having daughters who carry one copy of the mutated gene.
- **Codominant inheritance:** In codominant inheritance, each parent contributes a different version of a particular gene, and both versions influence the resulting genetic trait. The chance of developing a genetic condition with codominant inheritance, and the characteristic features of that condition, depend on which versions of the gene are passed from parents to their child.
- **Mitochondrial inheritance:** Mitochondria, which are the energy-producing centers inside cells, each contain a small amount of DNA. Disorders with mitochondrial inheritance result from mutations in mitochondrial DNA. Although mitochondrial

disorders can affect both males and females, only females can pass mutations in mitochondrial DNA to their children. A woman with a disorder caused by changes in mitochondrial DNA will pass the mutation to all of her daughters and sons, but the children of a man with such a disorder will not inherit the mutation.

It is important to note that the chance of passing on a genetic condition applies equally to each pregnancy. For example, if a couple has a child with an autosomal recessive disorder, the chance of having another child with the disorder is still 25 percent (or 1 in 4). Having one child with a disorder does not “protect” future children from inheriting the condition. Conversely, having a child without the condition does not mean that future children will definitely be affected.

Although the chances of inheriting a genetic condition appear straightforward, factors such as a person’s family history and the results of genetic testing can sometimes modify those chances. In addition, some people with a disease-causing mutation never develop any health problems or may experience only mild symptoms of the disorder. If a disease that runs in a family does not have a clear-cut inheritance pattern, predicting the likelihood that a person will develop the condition can be particularly difficult.

Estimating the chance of developing or passing on a genetic disorder can be complex. Genetics professionals can help people understand these chances and help them make informed decisions about their health.

Factors that Influence the Effects of Particular Genetic Changes

Reduced penetrance and variable expressivity are factors that influence the effects of particular genetic changes. These factors usually affect disorders that have an autosomal dominant pattern of inheritance, although they are occasionally seen in disorders with an autosomal recessive inheritance pattern.

Reduced Penetrance

Penetrance refers to the proportion of people with a particular genetic change (such as a mutation in a specific gene) who exhibit signs and symptoms of a genetic disorder. If some people with the mutation do not develop features of the disorder, the condition is said to have reduced (or incomplete) penetrance. Reduced penetrance often occurs with familial cancer syndromes. For example, many people with a mutation in the BRCA1 or BRCA2 gene will develop cancer during their lifetime, but some people will not. Doctors cannot predict which people with these mutations will develop cancer or when the tumors will develop.

Reduced penetrance probably results from a combination of genetic, environmental, and lifestyle factors, many of which are unknown. This phenomenon can make it challenging for genetics professionals to interpret a person’s family medical history and predict the risk of passing a genetic condition to future generations.

Variable Expressivity

Although some genetic disorders exhibit little variation, most have signs and symptoms that differ among affected individuals. Variable expressivity refers to the range of signs and

symptoms that can occur in different people with the same genetic condition. For example, the features of Marfan syndrome vary widely— some people have only mild symptoms (such as being tall and thin with long, slender fingers), while others also experience life-threatening complications involving the heart and blood vessels. Although the features are highly variable, most people with this disorder have a mutation in the same gene (FBN1).

As with reduced penetrance, variable expressivity is probably caused by a combination of genetic, environmental, and lifestyle factors, most of which have not been identified. If a genetic condition has highly variable signs and symptoms, it may be challenging to diagnose.

What Do Geneticists Mean by Anticipation?

The signs and symptoms of some genetic conditions tend to become more severe and appear at an earlier age as the disorder is passed from one generation to the next. This phenomenon is called anticipation. Anticipation is most often seen with certain genetic disorders of the nervous system, such as Huntington disease, myotonic dystrophy, and fragile X syndrome.

Anticipation typically occurs with disorders that are caused by an unusual type of mutation called a trinucleotide repeat expansion. A trinucleotide repeat is a sequence of three DNA building blocks (nucleotides) that is repeated a number of times in a row. DNA segments with an abnormal number of these repeats are unstable and prone to errors during cell division. The number of repeats can change as the gene is passed from parent to child. If the number of repeats increases, it is known as a trinucleotide repeat expansion. In some cases, the trinucleotide repeat may expand until the gene stops functioning normally. This expansion causes the features of some disorders to become more severe with each successive generation.

Most genetic disorders have signs and symptoms that differ among affected individuals, including affected people in the same family. Not all of these differences can be explained by anticipation. A combination of genetic, environmental, and lifestyle factors is probably responsible for the variability, although many of these factors have not been identified. Researchers study multiple generations of affected family members and consider the genetic cause of a disorder before determining that it shows anticipation.

What Is Genomic Imprinting?

Genomic imprinting is a factor that influences how some genetic conditions are inherited.

People inherit two copies of their genes—one from their mother and one from their father. Usually both copies of each gene are active, or “turned on,” in cells. In some cases, however, only one of the two copies is normally turned on. Which copy is active depends on the parent of origin: some genes are normally active only when they are inherited from a person’s father; others are active only when inherited from a person’s mother. This phenomenon is known as genomic imprinting.

In genes that undergo genomic imprinting, the parent of origin is often marked, or “stamped,” on the gene during the formation of egg and sperm cells. This stamping process, called methylation, is a chemical reaction that attaches small molecules called methyl groups to certain segments of DNA. These molecules identify which copy of a gene was inherited

from the mother and which was inherited from the father. The addition and removal of methyl groups can be used to control the activity of genes.

Only a small percentage of all human genes undergo genomic imprinting. Researchers are not yet certain why some genes are imprinted and others are not. They do know that imprinted genes tend to cluster together in the same regions of chromosomes. Two major clusters of imprinted genes have been identified in humans, one on the short (p) arm of chromosome 11 (at position 11p15) and another on the long (q) arm of chromosome 15 (in the region 15q11 to 15q13).

What Is Uniparental Disomy?

Uniparental disomy is a factor that influences how some genetic conditions are inherited.

Uniparental disomy (UPD) occurs when a person receives two copies of a chromosome, or part of a chromosome, from one parent and no copies from the other parent. UPD can occur as a random event during the formation of egg or sperm cells or may happen in early fetal development.

In many cases, UPD likely has no effect on health or development. Because most genes are not imprinted, it doesn't matter if a person inherits both copies from one parent instead of one copy from each parent. In some cases, however, it does make a difference whether a gene is inherited from a person's mother or father. A person with UPD may lack any active copies of essential genes that undergo genomic imprinting. This loss of gene function can lead to delayed development, mental retardation, or other medical problems.

Several genetic disorders can result from UPD or a disruption of normal genomic imprinting. The most well-known conditions include Prader-Willi syndrome, which is characterized by uncontrolled eating and obesity, and Angelman syndrome, which causes mental retardation and impaired speech. Both of these disorders can be caused by UPD or other errors in imprinting involving genes on the long arm of chromosome 15. Other conditions, such as Beckwith-Wiedemann syndrome (a disorder characterized by accelerated growth and an increased risk of cancerous tumors), are associated with abnormalities of imprinted genes on the short arm of chromosome 11.

Are Chromosomal Disorders Inherited?

Although it is possible to inherit some types of chromosomal abnormalities, most chromosomal disorders (such as Down syndrome and Turner syndrome) are not passed from one generation to the next.

Some chromosomal conditions are caused by changes in the number of chromosomes. These changes are not inherited, but occur as random events during the formation of reproductive cells (eggs and sperm). An error in cell division called nondisjunction results in reproductive cells with an abnormal number of chromosomes. For example, a reproductive cell may accidentally gain or lose one copy of a chromosome. If one of these atypical reproductive cells contributes to the genetic makeup of a child, the child will have an extra or missing chromosome in each of the body's cells.

Changes in chromosome structure can also cause chromosomal disorders. Some changes in chromosome structure can be inherited, while others occur as random accidents during the formation of reproductive cells or in early fetal development. Because the inheritance of these changes can be complex, people concerned about this type of chromosomal abnormality may want to talk with a genetics professional.

Some cancer cells also have changes in the number or structure of their chromosomes. Because these changes occur in somatic cells (cells other than eggs and sperm), they cannot be passed from one generation to the next.

Why Are Some Genetic Conditions More Common in Particular Ethnic Groups?

Some genetic disorders are more likely to occur among people who trace their ancestry to a particular geographic area. People in an ethnic group often share certain versions of their genes, which have been passed down from common ancestors. If one of these shared genes contains a disease-causing mutation, a particular genetic disorder may be more frequently seen in the group.

Examples of genetic conditions that are more common in particular ethnic groups are sickle cell anemia, which is more common in people of African, African-American, or Mediterranean heritage; and Tay-Sachs disease, which is more likely to occur among people of Ashkenazi (eastern and central European) Jewish or French Canadian ancestry. It is important to note, however, that these disorders can occur in any ethnic group.

Genetic Consultation

This section presents information on finding and visiting a genetic counselor or other genetics professional.

What Is a Genetic Consultation?

A genetic consultation is a health service that provides information and support to people who have, or may be at risk for, genetic disorders. During a consultation, a genetics professional meets with an individual or family to discuss genetic risks or to diagnose, confirm, or rule out a genetic condition.

Genetics professionals include medical geneticists (doctors who specialize in genetics) and genetic counselors (certified healthcare workers with experience in medical genetics and counseling). Other healthcare professionals such as nurses, psychologists, and social workers trained in genetics can also provide genetic consultations.

Consultations usually take place in a doctor's office, hospital, genetics center, or other type of medical center. These meetings are most often in-person visits with individuals or families, but they are occasionally conducted in a group or over the telephone.

Why Might Someone Have a Genetic Consultation?

Individuals or families who are concerned about an inherited condition may benefit from a genetic consultation. The reasons that a person might be referred to a genetic counselor, medical geneticist, or other genetics professional include:

- A personal or family history of a genetic condition, birth defect, chromosomal disorder, or hereditary cancer.
- Two or more pregnancy losses (miscarriages), a stillbirth, or a baby who died.
- A child with a known inherited disorder, a birth defect, mental retardation, or developmental delay.
- A woman who is pregnant or plans to become pregnant at or after age 35. (Some chromosomal disorders occur more frequently in children born to older women.)
- Abnormal test results that suggest a genetic or chromosomal condition.
- An increased risk of developing or passing on a particular genetic disorder on the basis of a person's ethnic background.
- People related by blood (for example, cousins) who plan to have children together. (A child whose parents are related may be at an increased risk of inheriting certain genetic disorders.)

A genetic consultation is also an important part of the decision-making process for genetic testing. A visit with a genetics professional may be helpful even if testing is not available for a specific condition, however.

What Happens during a Genetic Consultation?

A genetic consultation provides information, offers support, and addresses a patient's specific questions and concerns. To help determine whether a condition has a genetic component, a genetics professional asks about a person's medical history and takes a detailed family history (a record of health information about a person's immediate and extended family). The genetics professional may also perform a physical examination and recommend appropriate tests.

If a person is diagnosed with a genetic condition, the genetics professional provides information about the diagnosis, how the condition is inherited, the chance of passing the condition to future generations, and the options for testing and treatment.

During a consultation, a genetics professional will:

- Interpret and communicate complex medical information.
- Help each person make informed, independent decisions about their health care and reproductive options.
- Respect each person's individual beliefs, traditions, and feelings.

A genetics professional will NOT:

- Tell a person which decision to make.
- Advise a couple not to have children.

- Recommend that a woman continue or end a pregnancy.
- Tell someone whether to undergo testing for a genetic disorder.

How Can I Find a Genetics Professional in My Area?

To find a genetics professional in your community, you may wish to ask your doctor for a referral. If you have health insurance, you can also contact your insurance company to find a medical geneticist or genetic counselor in your area who participates in your plan.

Several resources for locating a genetics professional in your community are available online:

- GeneTests from the University of Washington provides a list of genetics clinics around the United States and international genetics clinics. You can also access the list by clicking on “Clinic Directory” at the top of the GeneTests home page. Clinics can be chosen by state or country, by service, and/or by specialty. State maps can help you locate a clinic in your area. See <http://www.genetests.org/>.
- The National Society of Genetic Counselors offers a searchable directory of genetic counselors in the United States. You can search by location, name, area of practice/specialization, and/or ZIP Code. See <http://www.nsgc.org/resource/link.cfm>.
- The National Cancer Institute provides a Cancer Genetics Services Directory, which lists professionals who provide services related to cancer genetics. You can search by type of cancer or syndrome, location, and/or provider name at the following Web site: http://cancer.gov/search/genetics_services/.

Genetic Testing

This section presents information on the benefits, costs, risks, and limitations of genetic testing.

What Is Genetic Testing?

Genetic testing is a type of medical test that identifies changes in chromosomes, genes, or proteins. Most of the time, testing is used to find changes that are associated with inherited disorders. The results of a genetic test can confirm or rule out a suspected genetic condition or help determine a person’s chance of developing or passing on a genetic disorder. Several hundred genetic tests are currently in use, and more are being developed.

Genetic testing is voluntary. Because testing has both benefits and limitations, the decision about whether to be tested is a personal and complex one. A genetic counselor can help by providing information about the pros and cons of the test and discussing the social and emotional aspects of testing.

What Are the Types of Genetic Tests?

Genetic testing can provide information about a person's genes and chromosomes. Available types of testing include:

- **Newborn screening** is used just after birth to identify genetic disorders that can be treated early in life. Millions of babies are tested each year in the United States. All states currently test infants for phenylketonuria (a genetic disorder that causes mental retardation if left untreated) and congenital hypothyroidism (a disorder of the thyroid gland). Most states also test for other genetic disorders.
- **Diagnostic testing** is used to identify or rule out a specific genetic or chromosomal condition. In many cases, genetic testing is used to confirm a diagnosis when a particular condition is suspected based on physical signs and symptoms. Diagnostic testing can be performed before birth or at any time during a person's life, but is not available for all genes or all genetic conditions. The results of a diagnostic test can influence a person's choices about health care and the management of the disorder.
- **Carrier testing** is used to identify people who carry one copy of a gene mutation that, when present in two copies, causes a genetic disorder. This type of testing is offered to individuals who have a family history of a genetic disorder and to people in certain ethnic groups with an increased risk of specific genetic conditions. If both parents are tested, the test can provide information about a couple's risk of having a child with a genetic condition.
- **Prenatal testing** is used to detect changes in a fetus's genes or chromosomes before birth. This type of testing is offered during pregnancy if there is an increased risk that the baby will have a genetic or chromosomal disorder. In some cases, prenatal testing can lessen a couple's uncertainty or help them make decisions about a pregnancy. It cannot identify all possible inherited disorders and birth defects, however.
- **Preimplantation testing**, also called preimplantation genetic diagnosis (PGD), is a specialized technique that can reduce the risk of having a child with a particular genetic or chromosomal disorder. It is used to detect genetic changes in embryos that were created using assisted reproductive techniques such as in-vitro fertilization. In-vitro fertilization involves removing egg cells from a woman's ovaries and fertilizing them with sperm cells outside the body. To perform preimplantation testing, a small number of cells are taken from these embryos and tested for certain genetic changes. Only embryos without these changes are implanted in the uterus to initiate a pregnancy.
- **Predictive and presymptomatic types of testing** are used to detect gene mutations associated with disorders that appear after birth, often later in life. These tests can be helpful to people who have a family member with a genetic disorder, but who have no features of the disorder themselves at the time of testing. Predictive testing can identify mutations that increase a person's risk of developing disorders with a genetic basis, such as certain types of cancer. Presymptomatic testing can determine whether a person will develop a genetic disorder, such as hemochromatosis (an iron overload disorder), before any signs or symptoms appear. The results of predictive and presymptomatic testing can provide information about a person's risk of developing a specific disorder and help with making decisions about medical care.
- **Forensic testing** uses DNA sequences to identify an individual for legal purposes. Unlike the tests described above, forensic testing is not used to detect gene mutations associated with disease. This type of testing can identify crime or catastrophe victims, rule out or implicate a crime suspect, or establish biological relationships between people (for example, paternity).

How Is Genetic Testing Done?

Once a person decides to proceed with genetic testing, a medical geneticist, primary care doctor, specialist, or nurse practitioner can order the test. Genetic testing is often done as part of a genetic consultation.

Genetic tests are performed on a sample of blood, hair, skin, amniotic fluid (the fluid that surrounds a fetus during pregnancy), or other tissue. For example, a procedure called a buccal smear uses a small brush or cotton swab to collect a sample of cells from the inside surface of the cheek. The sample is sent to a laboratory where technicians look for specific changes in chromosomes, DNA, or proteins, depending on the suspected disorder. The laboratory reports the test results in writing to a person's doctor or genetic counselor.

Newborn screening tests are done on a small blood sample, which is taken by pricking the baby's heel. Unlike other types of genetic testing, a parent will usually only receive the result if it is positive. If the test result is positive, additional testing is needed to determine whether the baby has a genetic disorder.

Before a person has a genetic test, it is important that he or she understands the testing procedure, the benefits and limitations of the test, and the possible consequences of the test results. The process of educating a person about the test and obtaining permission is called informed consent.

What Is Direct-to-Consumer Genetic Testing?

Traditionally, genetic tests have been available only through healthcare providers such as physicians, nurse practitioners, and genetic counselors. Healthcare providers order the appropriate test from a laboratory, collect and send the samples, and interpret the test results. Direct-to-consumer genetic testing refers to genetic tests that are marketed directly to consumers via television, print advertisements, or the Internet. This form of testing, which is also known as at-home genetic testing, provides access to a person's genetic information without necessarily involving a doctor or insurance company in the process.

If a consumer chooses to purchase a genetic test directly, the test kit is mailed to the consumer instead of being ordered through a doctor's office. The test typically involves collecting a DNA sample at home, often by swabbing the inside of the cheek, and mailing the sample back to the laboratory. In some cases, the person must visit a health clinic to have blood drawn. Consumers are notified of their results by mail or over the telephone, or the results are posted online. In some cases, a genetic counselor or other healthcare provider is available to explain the results and answer questions. The price for this type of at-home genetic testing ranges from several hundred dollars to more than a thousand dollars.

The growing market for direct-to-consumer genetic testing may promote awareness of genetic diseases, allow consumers to take a more proactive role in their health care, and offer a means for people to learn about their ancestral origins. At-home genetic tests, however, have significant risks and limitations. Consumers are vulnerable to being misled by the results of unproven or invalid tests. Without guidance from a healthcare provider, they may make important decisions about treatment or prevention based on inaccurate, incomplete, or misunderstood information about their health. Consumers may also experience an invasion

of genetic privacy if testing companies use their genetic information in an unauthorized way.

Genetic testing provides only one piece of information about a person's health—other genetic and environmental factors, lifestyle choices, and family medical history also affect a person's risk of developing many disorders. These factors are discussed during a consultation with a doctor or genetic counselor, but in many cases are not addressed by at-home genetic tests. More research is needed to fully understand the benefits and limitations of direct-to-consumer genetic testing.

What Do the Results of Genetic Tests Mean?

The results of genetic tests are not always straightforward, which often makes them challenging to interpret and explain. Therefore, it is important for patients and their families to ask questions about the potential meaning of genetic test results both before and after the test is performed. When interpreting test results, healthcare professionals consider a person's medical history, family history, and the type of genetic test that was done.

A positive test result means that the laboratory found a change in a particular gene, chromosome, or protein of interest. Depending on the purpose of the test, this result may confirm a diagnosis, indicate that a person is a carrier of a particular genetic mutation, identify an increased risk of developing a disease (such as cancer) in the future, or suggest a need for further testing. Because family members have some genetic material in common, a positive test result may also have implications for certain blood relatives of the person undergoing testing. It is important to note that a positive result of a predictive or presymptomatic genetic test usually cannot establish the exact risk of developing a disorder. Also, health professionals typically cannot use a positive test result to predict the course or severity of a condition.

A negative test result means that the laboratory did not find a change in the gene, chromosome, or protein under consideration. This result can indicate that a person is not affected by a particular disorder, is not a carrier of a specific genetic mutation, or does not have an increased risk of developing a certain disease. It is possible, however, that the test missed a disease-causing genetic alteration because many tests cannot detect all genetic changes that can cause a particular disorder. Further testing may be required to confirm a negative result.

In some cases, a negative result might not give any useful information. This type of result is called uninformative, indeterminate, inconclusive, or ambiguous. Uninformative test results sometimes occur because everyone has common, natural variations in their DNA, called polymorphisms, that do not affect health. If a genetic test finds a change in DNA that has not been associated with a disorder in other people, it can be difficult to tell whether it is a natural polymorphism or a disease-causing mutation. An uninformative result cannot confirm or rule out a specific diagnosis, and it cannot indicate whether a person has an increased risk of developing a disorder. In some cases, testing other affected and unaffected family members can help clarify this type of result.

What Is the Cost of Genetic Testing, and How Long Does It Take to Get the Results?

The cost of genetic testing can range from under \$100 to more than \$2,000, depending on the nature and complexity of the test. The cost increases if more than one test is necessary or if multiple family members must be tested to obtain a meaningful result. For newborn screening, costs vary by state. Some states cover part of the total cost, but most charge a fee of \$15 to \$60 per infant.

From the date that a sample is taken, it may take a few weeks to several months to receive the test results. Results for prenatal testing are usually available more quickly because time is an important consideration in making decisions about a pregnancy. The doctor or genetic counselor who orders a particular test can provide specific information about the cost and time frame associated with that test.

Will Health Insurance Cover the Costs of Genetic Testing?

In many cases, health insurance plans will cover the costs of genetic testing when it is recommended by a person's doctor. Health insurance providers have different policies about which tests are covered, however. A person interested in submitting the costs of testing may wish to contact his or her insurance company beforehand to ask about coverage.

Some people may choose not to use their insurance to pay for testing because the results of a genetic test can affect a person's health insurance coverage. Instead, they may opt to pay out-of-pocket for the test. People considering genetic testing may want to find out more about their state's privacy protection laws before they ask their insurance company to cover the costs.

What Are the Benefits of Genetic Testing?

Genetic testing has potential benefits whether the results are positive or negative for a gene mutation. Test results can provide a sense of relief from uncertainty and help people make informed decisions about managing their health care. For example, a negative result can eliminate the need for unnecessary checkups and screening tests in some cases. A positive result can direct a person toward available prevention, monitoring, and treatment options. Some test results can also help people make decisions about having children. Newborn screening can identify genetic disorders early in life so treatment can be started as early as possible.

What Are the Risks and Limitations of Genetic Testing?

The physical risks associated with most genetic tests are very small, particularly for those tests that require only a blood sample or buccal smear (a procedure that samples cells from the inside surface of the cheek). The procedures used for prenatal testing carry a small but real risk of losing the pregnancy (miscarriage) because they require a sample of amniotic fluid or tissue from around the fetus.

Many of the risks associated with genetic testing involve the emotional, social, or financial consequences of the test results. People may feel angry, depressed, anxious, or guilty about

their results. In some cases, genetic testing creates tension within a family because the results can reveal information about other family members in addition to the person who is tested. The possibility of genetic discrimination in employment or insurance is also a concern.

Genetic testing can provide only limited information about an inherited condition. The test often can't determine if a person will show symptoms of a disorder, how severe the symptoms will be, or whether the disorder will progress over time. Another major limitation is the lack of treatment strategies for many genetic disorders once they are diagnosed.

A genetics professional can explain in detail the benefits, risks, and limitations of a particular test. It is important that any person who is considering genetic testing understand and weigh these factors before making a decision.

What Is Genetic Discrimination?

Genetic discrimination occurs when people are treated differently by their employer or insurance company because they have a gene mutation that causes or increases the risk of an inherited disorder. People who undergo genetic testing may be at risk for genetic discrimination.

The results of a genetic test are normally included in a person's medical records. When a person applies for life, disability, or health insurance, the insurance company may ask to look at these records before making a decision about coverage. An employer may also have the right to look at an employee's medical records. As a result, genetic test results could affect a person's insurance coverage or employment. People making decisions about genetic testing should be aware that when test results are placed in their medical records, the results might not be kept private.

Fear of discrimination is a common concern among people considering genetic testing. Several laws at the federal and state levels help protect people against genetic discrimination; however, genetic testing is a fast-growing field and these laws don't cover every situation.

How Does Genetic Testing in a Research Setting Differ from Clinical Genetic Testing?

The main differences between clinical genetic testing and research testing are the purpose of the test and who receives the results. The goals of research testing include finding unknown genes, learning how genes work, and advancing our understanding of genetic conditions. The results of testing done as part of a research study are usually not available to patients or their healthcare providers. Clinical testing, on the other hand, is done to find out about an inherited disorder in an individual patient or family. People receive the results of a clinical test and can use them to help them make decisions about medical care or reproductive issues.

It is important for people considering genetic testing to know whether the test is available on a clinical or research basis. Clinical and research testing both involve a process of informed consent in which patients learn about the testing procedure, the risks and benefits of the test, and the potential consequences of testing.

Gene Therapy

This section presents information on experimental techniques, safety, ethics, and availability of gene therapy.

What Is Gene Therapy?

Gene therapy is an experimental technique that uses genes to treat or prevent disease. In the future, this technique may allow doctors to treat a disorder by inserting a gene into a patient's cells instead of using drugs or surgery. Researchers are testing several approaches to gene therapy, including:

- Replacing a mutated gene that causes disease with a healthy copy of the gene.
- Inactivating, or "knocking out," a mutated gene that is functioning improperly.
- Introducing a new gene into the body to help fight a disease.

Although gene therapy is a promising treatment option for a number of diseases (including inherited disorders, some types of cancer, and certain viral infections), the technique remains risky and is still under study to make sure that it will be safe and effective. Gene therapy is currently only being tested for the treatment of diseases that have no other cures.


How Does Gene Therapy Work?

Gene therapy is designed to introduce genetic material into cells to compensate for abnormal genes or to make a beneficial protein. If a mutated gene causes a necessary protein to be faulty or missing, gene therapy may be able to introduce a normal copy of the gene to restore the function of the protein.

A gene that is inserted directly into a cell usually does not function. Instead, a carrier called a vector is genetically engineered to deliver the gene. Certain viruses are often used as vectors because they can deliver the new gene by infecting the cell. The viruses are modified so they can't cause disease when used in people. Some types of virus, such as retroviruses, integrate their genetic material (including the new gene) into a chromosome in the human cell. Other viruses, such as adenoviruses, introduce their DNA into the nucleus of the cell, but the DNA is not integrated into a chromosome.

The vector can be injected or given intravenously (by IV) directly into a specific tissue in the body, where it is taken up by individual cells. Alternately, a sample of the patient's cells can be removed and exposed to the vector in a laboratory setting. The cells containing the vector are then returned to the patient. If the treatment is successful, the new gene delivered by the vector will make a functioning protein.

Researchers must overcome many technical challenges before gene therapy will be a practical approach to treating disease. For example, scientists must find better ways to deliver genes and target them to particular cells. They must also ensure that new genes are precisely controlled by the body.



A new gene is injected into an adenovirus vector, which is used to introduce the modified DNA into a human cell. If the treatment is successful, the new gene will make a functional protein.

Is Gene Therapy Safe?

Gene therapy is under study to determine whether it could be used to treat disease. Current research is evaluating the safety of gene therapy; future studies will test whether it is an effective treatment option. Several studies have already shown that this approach can have very serious health risks, such as toxicity, inflammation, and cancer. Because the techniques are relatively new, some of the risks may be unpredictable; however, medical researchers, institutions, and regulatory agencies are working to ensure that gene therapy research is as safe as possible.

Comprehensive federal laws, regulations, and guidelines help protect people who participate in research studies (called clinical trials). The U.S. Food and Drug Administration (FDA) regulates all gene therapy products in the United States and oversees research in this area. Researchers who wish to test an approach in a clinical trial must first obtain permission from the FDA. The FDA has the authority to reject or suspend clinical trials that are suspected of being unsafe for participants.

The National Institutes of Health (NIH) also plays an important role in ensuring the safety of gene therapy research. NIH provides guidelines for investigators and institutions (such as universities and hospitals) to follow when conducting clinical trials with gene therapy. These guidelines state that clinical trials at institutions receiving NIH funding for this type of research must be registered with the NIH Office of Biotechnology Activities. The protocol, or plan, for each clinical trial is then reviewed by the NIH Recombinant DNA Advisory Committee (RAC) to determine whether it raises medical, ethical, or safety issues that warrant further discussion at one of the RAC's public meetings.

An Institutional Review Board (IRB) and an Institutional Biosafety Committee (IBC) must approve each gene therapy clinical trial before it can be carried out. An IRB is a committee of scientific and medical advisors and consumers that reviews all research within an institution. An IBC is a group that reviews and approves an institution's potentially hazardous research studies. Multiple levels of evaluation and oversight ensure that safety concerns are a top priority in the planning and carrying out of gene therapy research.

What Are the Ethical Issues surrounding Gene Therapy?

Because gene therapy involves making changes to the body's set of basic instructions, it raises many unique ethical concerns. The ethical questions surrounding gene therapy include:

- How can "good" and "bad" uses of gene therapy be distinguished?
- Who decides which traits are normal and which constitute a disability or disorder?
- Will the high costs of gene therapy make it available only to the wealthy?
- Could the widespread use of gene therapy make society less accepting of people who are different?
- Should people be allowed to use gene therapy to enhance basic human traits such as height, intelligence, or athletic ability?

Current gene therapy research has focused on treating individuals by targeting the therapy to body cells such as bone marrow or blood cells. This type of gene therapy cannot be passed on to a person's children. Gene therapy could be targeted to egg and sperm cells (germ cells), however, which would allow the inserted gene to be passed on to future generations. This approach is known as germline gene therapy.

The idea of germline gene therapy is controversial. While it could spare future generations in a family from having a particular genetic disorder, it might affect the development of a fetus in unexpected ways or have long-term side effects that are not yet known. Because people who would be affected by germline gene therapy are not yet born, they can't choose whether to have the treatment. Because of these ethical concerns, the U.S. Government does not allow federal funds to be used for research on germline gene therapy in people.

Is Gene Therapy Available to Treat My Disorder?

Gene therapy is currently available only in a research setting. The U.S. Food and Drug Administration (FDA) has not yet approved any gene therapy products for sale in the United States.

Hundreds of research studies (clinical trials) are under way to test gene therapy as a treatment for genetic conditions, cancer, and HIV/AIDS. If you are interested in participating in a clinical trial, talk with your doctor or a genetics professional about how to participate.

You can also search for clinical trials online. ClinicalTrials.gov, a service of the National Institutes of Health, provides easy access to information on clinical trials. You can search for

specific trials or browse by condition or trial sponsor. You may wish to refer to a list of gene therapy trials that are accepting (or will accept) patients.

The Human Genome Project and Genomic Research

This section presents information on the goals, accomplishments, and next steps in understanding the human genome.

What Is a Genome?

A genome is an organism's complete set of DNA, including all of its genes. Each genome contains all of the information needed to build and maintain that organism. In humans, a copy of the entire genome—more than 3 billion DNA base pairs—is contained in all cells that have a nucleus.

What Was the Human Genome Project and Why Has It Been Important?

The Human Genome Project was an international research effort to determine the sequence of the human genome and identify the genes that it contains. The Project was coordinated by the National Institutes of Health and the U.S. Department of Energy. Additional contributors included universities across the United States and international partners in the United Kingdom, France, Germany, Japan, and China. The Human Genome Project formally began in 1990 and was completed in 2003, 2 years ahead of its original schedule.

The work of the Human Genome Project has allowed researchers to begin to understand the blueprint for building a person. As researchers learn more about the functions of genes and proteins, this knowledge will have a major impact in the fields of medicine, biotechnology, and the life sciences.

What Were the Goals of the Human Genome Project?

The main goals of the Human Genome Project were to provide a complete and accurate sequence of the 3 billion DNA base pairs that make up the human genome and to find all of the estimated 20,000 to 25,000 human genes. The Project also aimed to sequence the genomes of several other organisms that are important to medical research, such as the mouse and the fruit fly.

In addition to sequencing DNA, the Human Genome Project sought to develop new tools to obtain and analyze the data and to make this information widely available. Also, because advances in genetics have consequences for individuals and society, the Human Genome Project committed to exploring the consequences of genomic research through its Ethical, Legal, and Social Implications (ELSI) program.

What Did the Human Genome Project Accomplish?

In April 2003, researchers announced that the Human Genome Project had completed a high-quality sequence of essentially the entire human genome. This sequence closed the gaps from a working draft of the genome, which was published in 2001. It also identified the

locations of many human genes and provided information about their structure and organization. The Project made the sequence of the human genome and tools to analyze the data freely available via the Internet.

In addition to the human genome, the Human Genome Project sequenced the genomes of several other organisms, including brewers' yeast, the roundworm, and the fruit fly. In 2002, researchers announced that they had also completed a working draft of the mouse genome. By studying the similarities and differences between human genes and those of other organisms, researchers can discover the functions of particular genes and identify which genes are critical for life.

The Project's Ethical, Legal, and Social Implications (ELSI) program became the world's largest bioethics program and a model for other ELSI programs worldwide.

What Were Some of the Ethical, Legal, and Social Implications Addressed by the Human Genome Project?

The Ethical, Legal, and Social Implications (ELSI) program was founded in 1990 as an integral part of the Human Genome Project. The mission of the ELSI program was to identify and address issues raised by genomic research that would affect individuals, families, and society. A percentage of the Human Genome Project budget at the National Institutes of Health and the U.S. Department of Energy was devoted to ELSI research.

The ELSI program focused on the possible consequences of genomic research in four main areas:

- Privacy and fairness in the use of genetic information, including the potential for genetic discrimination in employment and insurance.
- The integration of new genetic technologies, such as genetic testing, into the practice of clinical medicine.
- Ethical issues surrounding the design and conduct of genetic research with people, including the process of informed consent.
- The education of healthcare professionals, policy makers, students, and the public about genetics and the complex issues that result from genomic research.

What Are the Next Steps in Genomic Research?

Discovering the sequence of the human genome was only the first step in understanding how the instructions coded in DNA lead to a functioning human being. The next stage of genomic research will begin to derive meaningful knowledge from the DNA sequence. Research studies that build on the work of the Human Genome Project are under way worldwide.

The objectives of continued genomic research include the following:

- Determine the function of genes and the elements that regulate genes throughout the genome.

- Find variations in the DNA sequence among people and determine their significance. These variations may one day provide information about a person's disease risk and response to certain medications.
- Discover the 3-dimensional structures of proteins and identify their functions.
- Explore how DNA and proteins interact with one another and with the environment to create complex living systems.
- Develop and apply genome-based strategies for the early detection, diagnosis, and treatment of disease.
- Sequence the genomes of other organisms, such as the rat, cow, and chimpanzee, in order to compare similar genes between species.
- Develop new technologies to study genes and DNA on a large scale and store genomic data efficiently.
- Continue to explore the ethical, legal, and social issues raised by genomic research.

What Is Pharmacogenomics?

Pharmacogenomics is the study of how genes affect a person's response to drugs. This relatively new field combines pharmacology (the science of drugs) and genomics (the study of genes and their functions) to develop effective, safe medications and doses that will be tailored to a person's genetic makeup.

Many drugs that are currently available are "one size fits all," but they don't work the same way for everyone. It can be difficult to predict who will benefit from a medication, who will not respond at all, and who will experience negative side effects (called adverse drug reactions). Adverse drug reactions are a significant cause of hospitalizations and deaths in the United States. With the knowledge gained from the Human Genome Project, researchers are learning how inherited differences in genes affect the body's response to medications. These genetic differences will be used to predict whether a medication will be effective for a particular person and to help prevent adverse drug reactions.

The field of pharmacogenomics is still in its infancy. Its use is currently quite limited, but new approaches are under study in clinical trials. In the future, pharmacogenomics will allow the development of tailored drugs to treat a wide range of health problems, including cardiovascular disease, Alzheimer disease, cancer, HIV/AIDS, and asthma.

APPENDIX B. PHYSICIAN RESOURCES

Overview

In this chapter, we focus on databases and Internet-based guidelines and information resources created or written for a professional audience.

NIH Guidelines

Commonly referred to as “clinical” or “professional” guidelines, the National Institutes of Health publish physician guidelines for the most common diseases. Publications are available at the following by relevant Institute⁹:

- National Institutes of Health (NIH); guidelines consolidated across agencies available at <http://health.nih.gov/>
- National Institute of General Medical Sciences (NIGMS); fact sheets available at <http://www.nigms.nih.gov/Publications/FactSheets.htm>
- National Library of Medicine (NLM); extensive encyclopedia (A.D.A.M., Inc.) with guidelines: <http://www.nlm.nih.gov/medlineplus/healthtopics.html>
- National Cancer Institute (NCI); guidelines available at <http://www.cancer.gov/cancertopics/pdq>
- National Eye Institute (NEI); guidelines available at <http://www.nei.nih.gov/health/>
- National Heart, Lung, and Blood Institute (NHLBI); guidelines available at <http://www.nhlbi.nih.gov/guidelines/index.htm>
- National Human Genome Research Institute (NHGRI); research available at <http://www.genome.gov/page.cfm?pageID=10000375>
- National Institute on Aging (NIA); guidelines available at <http://www.nia.nih.gov/HealthInformation/Publications/>
- National Institute on Alcohol Abuse and Alcoholism (NIAAA); guidelines available at <http://www.niaaa.nih.gov/Publications/>

⁹ These publications are typically written by one or more of the various NIH Institutes.

- National Institute of Allergy and Infectious Diseases (NIAID); guidelines available at <http://www.niaid.nih.gov/publications/>
- National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS); fact sheets and guidelines available at <http://www.niams.nih.gov/hi/index.htm>
- National Institute of Child Health and Human Development (NICHD); guidelines available at <http://www.nichd.nih.gov/publications/pubskey.cfm>
- National Institute on Deafness and Other Communication Disorders (NIDCD); fact sheets and guidelines at <http://www.nidcd.nih.gov/health/>
- National Institute of Dental and Craniofacial Research (NIDCR); guidelines available at <http://www.nidcr.nih.gov/HealthInformation/>
- National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK); guidelines available at <http://www.niddk.nih.gov/health/health.htm>
- National Institute on Drug Abuse (NIDA); guidelines available at <http://www.nida.nih.gov/DrugAbuse.html>
- National Institute of Environmental Health Sciences (NIEHS); environmental health information available at <http://www.niehs.nih.gov/external/facts.htm>
- National Institute of Mental Health (NIMH); guidelines available at <http://www.nimh.nih.gov/healthinformation/index.cfm>
- National Institute of Neurological Disorders and Stroke (NINDS); neurological disorder information pages available at http://www.ninds.nih.gov/health_and_medical/disorder_index.htm
- National Institute of Biomedical Imaging and Bioengineering; general information at <http://www.nibib.nih.gov/HealthEdu>
- National Center for Complementary and Alternative Medicine (NCCAM); health information available at <http://nccam.nih.gov/health/>
- National Center for Research Resources (NCRR); various information directories available at <http://www.ncrr.nih.gov/publications.asp>
- Office of Rare Diseases; various fact sheets available at http://rarediseases.info.nih.gov/html/resources/rep_pubs.html
- Centers for Disease Control and Prevention; various fact sheets on infectious diseases available at <http://www.cdc.gov/publications.htm>

NIH Databases

In addition to the various Institutes of Health that publish professional guidelines, the NIH has designed a number of databases for professionals.¹⁰ Physician-oriented resources provide a wide variety of information related to the biomedical and health sciences, both past and present. The format of these resources varies. Searchable databases, bibliographic

¹⁰ Remember, for the general public, the National Library of Medicine recommends the databases referenced in MEDLINEplus (<http://medlineplus.gov/> or <http://www.nlm.nih.gov/medlineplus/databases.html>).

citations, full-text articles (when available), archival collections, and images are all available. The following are referenced by the National Library of Medicine¹¹:

- **Bioethics:** Access to published literature on the ethical, legal, and public policy issues surrounding healthcare and biomedical research. This information is provided in conjunction with the Kennedy Institute of Ethics located at Georgetown University, Washington, D.C.: http://www.nlm.nih.gov/databases/databases_bioethics.html
- **HIV/AIDS Resources:** Describes various links and databases dedicated to HIV/AIDS research: <http://www.nlm.nih.gov/pubs/factsheets/aidsinfs.html>
- **NLM Online Exhibitions:** Describes “Exhibitions in the History of Medicine”: <http://www.nlm.nih.gov/exhibition/exhibition.html>. Additional resources for historical scholarship in medicine: <http://www.nlm.nih.gov/hmd/index.html>
- **Biotechnology Information:** Access to public databases. The National Center for Biotechnology Information conducts research in computational biology, develops software tools for analyzing genome data, and disseminates biomedical information for the better understanding of molecular processes affecting human health and disease: <http://www.ncbi.nlm.nih.gov/>
- **Population Information:** The National Library of Medicine provides access to worldwide coverage of population, family planning, and related health issues, including family planning technology and programs, fertility, and population law and policy: http://www.nlm.nih.gov/databases/databases_population.html
- **Cancer Information:** Access to cancer-oriented databases: http://www.nlm.nih.gov/databases/databases_cancer.html
- **Profiles in Science:** Offering the archival collections of prominent twentieth-century biomedical scientists to the public through modern digital technology: <http://www.profiles.nlm.nih.gov/>
- **Chemical Information:** Provides links to various chemical databases and references: <http://sis.nlm.nih.gov/Chem/ChemMain.html>
- **Clinical Alerts:** Reports the release of findings from the NIH-funded clinical trials where such release could significantly affect morbidity and mortality: http://www.nlm.nih.gov/databases/alerts/clinical_alerts.html
- **Space Life Sciences:** Provides links and information to space-based research (including NASA): http://www.nlm.nih.gov/databases/databases_space.html
- **MEDLINE:** Bibliographic database covering the fields of medicine, nursing, dentistry, veterinary medicine, the healthcare system, and the pre-clinical sciences: http://www.nlm.nih.gov/databases/databases_medline.html
- **Toxicology and Environmental Health Information (TOXNET):** Databases covering toxicology and environmental health: <http://sis.nlm.nih.gov/Tox/ToxMain.html>
- **Visible Human Interface:** Anatomically detailed, three-dimensional representations of normal male and female human bodies: http://www.nlm.nih.gov/research/visible/visible_human.html

¹¹ See <http://www.nlm.nih.gov/databases/index.html>.

The NLM Gateway¹²

The NLM (National Library of Medicine) Gateway is a Web-based system that lets users search simultaneously in multiple retrieval systems at the U.S. National Library of Medicine (NLM). It allows users of NLM services to initiate searches from one Web interface, providing one-stop searching for many of NLM's information resources or databases.¹³ To use the NLM Gateway, simply go to the search site at <http://gateway.nlm.nih.gov/gw/Cmd>. Type **alkaptonuria** (or synonyms) into the search box and click **Search**. The results will be presented in a tabular form, indicating the number of references in each database category.

Results Summary

Category	Items Found
Journal Articles	725
Books / Periodicals / Audio Visual	5
Consumer Health	1
Meeting Abstracts	0
Other Collections	0
Total	731

HSTAT¹⁴

HSTAT is a free, Web-based resource that provides access to full-text documents used in healthcare decision-making.¹⁵ These documents include clinical practice guidelines, quick-reference guides for clinicians, consumer health brochures, evidence reports and technology assessments from the Agency for Healthcare Research and Quality (AHRQ), as well as AHRQ's Put Prevention Into Practice.¹⁶ Simply search by **alkaptonuria** (or synonyms) at the following Web site: <http://text.nlm.nih.gov>.

Coffee Break: Tutorials for Biologists¹⁷

Coffee Break is a general healthcare site that takes a scientific view of the news and covers recent breakthroughs in biology that may one day assist physicians in developing treatments. Here you will find a collection of short reports on recent biological discoveries. Each report incorporates interactive tutorials that demonstrate how bioinformatics tools are

¹² Adapted from NLM: <http://gateway.nlm.nih.gov/gw/Cmd?Overview.x>.

¹³ The NLM Gateway is currently being developed by the Lister Hill National Center for Biomedical Communications (LHNCBC) at the National Library of Medicine (NLM) of the National Institutes of Health (NIH).

¹⁴ Adapted from HSTAT: <http://www.nlm.nih.gov/pubs/factsheets/hstat.html>.

¹⁵ The HSTAT URL is <http://hstat.nlm.nih.gov/>.

¹⁶ Other important documents in HSTAT include: the National Institutes of Health (NIH) Consensus Conference Reports and Technology Assessment Reports; the HIV/AIDS Treatment Information Service (ATIS) resource documents; the Substance Abuse and Mental Health Services Administration's Center for Substance Abuse Treatment (SAMHSA/CSAT) Treatment Improvement Protocols (TIP) and Center for Substance Abuse Prevention (SAMHSA/CSAP) Prevention Enhancement Protocols System (PEPS); the Public Health Service (PHS) Preventive Services Task Force's *Guide to Clinical Preventive Services*; the independent, nonfederal Task Force on Community Services' *Guide to Community Preventive Services*; and the Health Technology Advisory Committee (HTAC) of the Minnesota Health Care Commission (MHCC) health technology evaluations.

¹⁷ Adapted from <http://www.ncbi.nlm.nih.gov/Coffeefbreak/Archive/FAQ.html>.

used as a part of the research process. Currently, all Coffee Breaks are written by NCBI staff.¹⁸ Each report is about 400 words and is usually based on a discovery reported in one or more articles from recently published, peer-reviewed literature.¹⁹ This site has new articles every few weeks, so it can be considered an online magazine of sorts. It is intended for general background information. You can access the Coffee Break Web site at the following hyperlink: <http://www.ncbi.nlm.nih.gov/Coffeebreak/>.

Other Commercial Databases

In addition to resources maintained by official agencies, other databases exist that are commercial ventures addressing medical professionals. Here are some examples that may interest you:

- **MD Consult:** Access to electronic clinical resources, see <http://www.mdconsult.com/>.
- **Medical Matrix:** Lists over 6000 medical Web sites and links to over 1.5 million documents with clinical content, see <http://www.medmatrix.org/>.
- **Medical World Search:** Searches full text from thousands of selected medical sites on the Internet; see <http://www.mwsearch.com/>.

¹⁸ The figure that accompanies each article is frequently supplied by an expert external to NCBI, in which case the source of the figure is cited. The result is an interactive tutorial that tells a biological story.

¹⁹ After a brief introduction that sets the work described into a broader context, the report focuses on how a molecular understanding can provide explanations of observed biology and lead to therapies for diseases. Each vignette is accompanied by a figure and hypertext links that lead to a series of pages that interactively show how NCBI tools and resources are used in the research process.

APPENDIX C. PATIENT RESOURCES

Overview

Official agencies, as well as federally funded institutions supported by national grants, frequently publish a variety of guidelines written with the patient in mind. These are typically called **Fact Sheets** or **Guidelines**. They can take the form of a brochure, information kit, pamphlet, or flyer. Often they are only a few pages in length. Since new guidelines on alkaptonuria can appear at any moment and be published by a number of sources, the best approach to finding guidelines is to systematically scan the Internet-based services that post them.

Patient Guideline Sources

This section directs you to sources which either publish fact sheets or can help you find additional guidelines on topics related to alkaptonuria. Due to space limitations, these sources are listed in a concise manner. Do not hesitate to consult the following sources by either using the Internet hyperlink provided, or, in cases where the contact information is provided, contacting the publisher or author directly.

The National Institutes of Health

The NIH gateway to patients is located at <http://health.nih.gov/>. From this site, you can search across various sources and institutes, a number of which are summarized below.

Topic Pages: MEDLINEplus

The National Library of Medicine has created a vast and patient-oriented healthcare information portal called MEDLINEplus. Within this Internet-based system are **health topic pages** which list links to available materials relevant to alkaptonuria. To access this system, log on to <http://www.nlm.nih.gov/medlineplus/healthtopics.html>. From there you can either search using the alphabetical index or browse by broad topic areas. Recently, MEDLINEplus listed the following when searched for **alkaptonuria**:

Metabolic Disorders

<http://www.nlm.nih.gov/medlineplus/metabolicdisorders.html>

You may also choose to use the search utility provided by MEDLINEplus at the following Web address: <http://www.nlm.nih.gov/medlineplus/>. Simply type a keyword into the search box and click **Search**. This utility is similar to the NIH search utility, with the exception that it only includes materials that are linked within the MEDLINEplus system (mostly patient-oriented information). It also has the disadvantage of generating unstructured results. We recommend, therefore, that you use this method only if you have a very targeted search.

The NIH Search Utility

The NIH search utility allows you to search for documents on over 100 selected Web sites that comprise the NIH-WEB-SPACE. Each of these servers is “crawled” and indexed on an ongoing basis. Your search will produce a list of various documents, all of which will relate in some way to alkaptonuria. The drawbacks of this approach are that the information is not organized by theme and that the references are often a mix of information for professionals and patients. Nevertheless, a large number of the listed Web sites provide useful background information. We can only recommend this route, therefore, for relatively rare or specific disorders, or when using highly targeted searches. To use the NIH search utility, visit the following Web page: <http://health.nih.gov/index.asp>. Under **Search Health Topics**, type **alkaptonuria** (or synonyms) into the search box, and click **Search**.

NORD (The National Organization of Rare Disorders, Inc.)

NORD provides an invaluable service to the public by publishing short yet comprehensive guidelines on over 1,000 diseases. NORD primarily focuses on rare diseases that might not be covered by the previously listed sources. NORD’s Web address is <http://www.rarediseases.org/>. A complete guide on alkaptonuria can be purchased from NORD for a nominal fee.

Additional Web Sources

A number of Web sites are available to the public that often link to government sites. These can also point you in the direction of essential information. The following is a representative sample:

- Family Village: <http://www.familyvillage.wisc.edu/specific.htm>
- Google: http://directory.google.com/Top/Health/Conditions_and_Diseases/
- Med Help International: <http://www.medhelp.org/HealthTopics/A.html>
- Open Directory Project: http://dmoz.org/Health/Conditions_and_Diseases/
- Yahoo.com: http://dir.yahoo.com/Health/Diseases_and_Conditions/
- WebMD®Health: http://www.webmd.com/diseases_and_conditions/default.htm

Finding Associations

There are several Internet directories that provide lists of medical associations with information on or resources relating to alkaptonuria. By consulting all of associations listed in this chapter, you will have nearly exhausted all sources for patient associations concerned with alkaptonuria.

The National Health Information Center (NHIC)

The National Health Information Center (NHIC) offers a free referral service to help people find organizations that provide information about alkaptonuria. For more information, see the NHIC's Web site at <http://www.health.gov/NHIC/> or contact an information specialist by calling 1-800-336-4797.

Directory of Health Organizations

The Directory of Health Organizations, provided by the National Library of Medicine Specialized Information Services, is a comprehensive source of information on associations. The Directory of Health Organizations database can be accessed via the Internet at <http://sis.nlm.nih.gov/dirline.html>. It is composed of two parts: DIRLINE and Health Hotlines.

The DIRLINE database comprises some 10,000 records of organizations, research centers, and government institutes and associations that primarily focus on health and biomedicine. Simply type in **alkaptonuria** (or a synonym), and you will receive information on all relevant organizations listed in the database.

Health Hotlines directs you to toll-free numbers to over 300 organizations. You can access this database directly at <http://healthhotlines.nlm.nih.gov/>. On this page, you are given the option to search by keyword or by browsing the subject list. When you have received your search results, click on the name of the organization for its description and contact information.

The National Organization for Rare Disorders, Inc.

The National Organization for Rare Disorders, Inc. has prepared a Web site that provides, at no charge, lists of associations organized by health topic. You can access this database at the following Web site: <http://www.rarediseases.org/search/orgsearch.html>. Type **alkaptonuria** (or a synonym) into the search box, and click **Submit Query**.

Resources for Patients and Families

The following are organizations that provide support and advocacy for patient with genetic conditions and their families²⁰:

²⁰ Adapted from the National Library of Medicine: <http://ghr.nlm.nih.gov/ghr/resource/patients>.

- Genetic Alliance: <http://geneticalliance.org>
- Genetic and Rare Diseases Information Center: http://rarediseases.info.nih.gov/html/resources/info_cntr.html
- Madisons Foundation: <http://www.madisonsfoundation.org/>
- March of Dimes: <http://www.marchofdimes.com>
- National Organization for Rare Disorders (NORD): <http://www.rarediseases.org/>

For More Information on Genetics

The following publications offer detailed information for patients about the science of genetics:

- What Is a Genome?: http://www.ncbi.nlm.nih.gov/About/primer/genetics_genome.html
- A Science Called Genetics: <http://publications.nigms.nih.gov/genetics/science.html>
- Genetic Mapping: <http://www.genome.gov/10000715>

ONLINE GLOSSARIES

The Internet provides access to a number of free-to-use medical dictionaries. The National Library of Medicine has compiled the following list of online dictionaries:

- ADAM Medical Encyclopedia (A.D.A.M., Inc.), comprehensive medical reference:
<http://www.nlm.nih.gov/medlineplus/encyclopedia.html>
- MedicineNet.com Medical Dictionary (MedicineNet, Inc.):
<http://www.medterms.com/Script/Main/hp.asp>
- Merriam-Webster Medical Dictionary (Inteli-Health, Inc.):
<http://www.intelihealth.com/IH/>
- Multilingual Glossary of Technical and Popular Medical Terms in Eight European Languages (European Commission) - Danish, Dutch, English, French, German, Italian, Portuguese, and Spanish: <http://allserv.rug.ac.be/~rvdstich/eugloss/welcome.html>
- On-line Medical Dictionary (CancerWEB): <http://cancerweb.ncl.ac.uk/omd/>
- Rare Diseases Terms (Office of Rare Diseases):
<http://ord.aspensys.com/asp/diseases/diseases.asp>
- Technology Glossary (National Library of Medicine) - Health Care Technology:
<http://www.nlm.nih.gov/archive/20040831/nichsr/ta101/ta10108.html>

Beyond these, MEDLINEplus contains a very patient-friendly encyclopedia covering every aspect of medicine (licensed from A.D.A.M., Inc.). The ADAM Medical Encyclopedia can be accessed at <http://www.nlm.nih.gov/medlineplus/encyclopedia.html>. ADAM is also available on commercial Web sites such as drkoop.com (<http://www.drkoop.com/>) and Web MD (http://my.webmd.com/adam/asset/adam_disease_articles/a_to_z/a). The NIH suggests the following Web sites in the ADAM Medical Encyclopedia when searching for information on alkaptonuria:

- **Basic Guidelines for Alkaptonuria**

Alcaptonuria

Web site: <http://www.nlm.nih.gov/medlineplus/ency/article/001200.htm>

- **Diagnostics and Tests for Alkaptonuria**

Urinalysis

Web site: <http://www.nlm.nih.gov/medlineplus/ency/article/003579.htm>

- **Background Topics for Alkaptonuria**

Autosomal recessive

Web site: <http://www.nlm.nih.gov/medlineplus/ency/article/002052.htm>

Enzyme

Web site: <http://www.nlm.nih.gov/medlineplus/ency/article/002353.htm>

Metabolism

Web site: <http://www.nlm.nih.gov/medlineplus/ency/article/002257.htm>

Sclera

Web site: <http://www.nlm.nih.gov/medlineplus/ency/article/002295.htm>

Online Dictionary Directories

The following are additional online directories compiled by the National Library of Medicine, including a number of specialized medical dictionaries:

- Medical Dictionaries: Medical & Biological (World Health Organization):
<http://www.who.int/hlt/virtuallibrary/English/diction.htm#Medical>
- Patient Education: Glossaries (DMOZ Open Directory Project):
http://dmoz.org/Health/Education/Patient_Education/Glossaries/
- Web of Online Dictionaries (Bucknell University):
<http://www.yourdictionary.com/diction5.html#medicine>

ALKAPTONURIA DICTIONARY

The definitions below are derived from official public sources, including the National Institutes of Health [NIH] and the European Union [EU].

3-dimensional: 3-D. A graphic display of depth, width, and height. Three-dimensional radiation therapy uses computers to create a 3-dimensional picture of the tumor. This allows doctors to give the highest possible dose of radiation to the tumor, while sparing the normal tissue as much as possible. [NIH]

Abdomen: That portion of the body that lies between the thorax and the pelvis. [NIH]

Acceptor: A substance which, while normally not oxidized by oxygen or reduced by hydrogen, can be oxidized or reduced in presence of a substance which is itself undergoing oxidation or reduction. [NIH]

Acromonium: A mitosporic fungal genus with many reported ascomycetous teleomorphs. Cephalosporin antibiotics are derived from this genus. [NIH]

Adaptability: Ability to develop some form of tolerance to conditions extremely different from those under which a living organism evolved. [NIH]

Adenine: A purine base and a fundamental unit of adenine nucleotides. [NIH]

Adenosine: A nucleoside that is composed of adenine and d-ribose. Adenosine or adenosine derivatives play many important biological roles in addition to being components of DNA and RNA. Adenosine itself is a neurotransmitter. [NIH]

Adenosine Triphosphate: Adenosine 5'-(tetrahydrogen triphosphate). An adenine nucleotide containing three phosphate groups esterified to the sugar moiety. In addition to its crucial roles in metabolism adenosine triphosphate is a neurotransmitter. [NIH]

Adenovirus: A group of viruses that cause respiratory tract and eye infections. Adenoviruses used in gene therapy are altered to carry a specific tumor-fighting gene. [NIH]

Adverse Effect: An unwanted side effect of treatment. [NIH]

Aerobic: In biochemistry, reactions that need oxygen to happen or happen when oxygen is present. [NIH]

Algorithms: A procedure consisting of a sequence of algebraic formulas and/or logical steps to calculate or determine a given task. [NIH]

Alleles: Mutually exclusive forms of the same gene, occupying the same locus on homologous chromosomes, and governing the same biochemical and developmental process. [NIH]

Alpha-1: A protein with the property of inactivating proteolytic enzymes such as leucocyte collagenase and elastase. [NIH]

Alternative medicine: Practices not generally recognized by the medical community as standard or conventional medical approaches and used instead of standard treatments. Alternative medicine includes the taking of dietary supplements, megadose vitamins, and herbal preparations; the drinking of special teas; and practices such as massage therapy, magnet therapy, spiritual healing, and meditation. [NIH]

Amino Acids: Organic compounds that generally contain an amino (-NH₂) and a carboxyl (-COOH) group. Twenty alpha-amino acids are the subunits which are polymerized to form proteins. [NIH]

Amino Acids: Organic compounds that generally contain an amino (-NH₂) and a carboxyl (-COOH) group. Twenty alpha-amino acids are the subunits which are polymerized to form proteins. [NIH]

Amnion: The extraembryonic membrane which contains the embryo and amniotic fluid. [NIH]

Amniotic Fluid: Amniotic cavity fluid which is produced by the amnion and fetal lungs and kidneys. [NIH]

Anatomical: Pertaining to anatomy, or to the structure of the organism. [EU]

Anemia: A reduction in the number of circulating erythrocytes or in the quantity of hemoglobin. [NIH]

Aneuploidy: The chromosomal constitution of cells which deviate from the normal by the addition or subtraction of chromosomes or chromosome pairs. In a normally diploid cell the loss of a chromosome pair is termed nullisomy (symbol: 2N-2), the loss of a single chromosome is monosomy (symbol: 2N-1), the addition of a chromosome pair is tetrasomy (symbol: 2N+2), the addition of a single chromosome is trisomy (symbol: 2N+1). [NIH]

Anions: Negatively charged atoms, radicals or groups of atoms which travel to the anode or positive pole during electrolysis. [NIH]

Antibacterial: A substance that destroys bacteria or suppresses their growth or reproduction. [EU]

Antibiotic: A drug used to treat infections caused by bacteria and other microorganisms. [NIH]

Antibody: A type of protein made by certain white blood cells in response to a foreign substance (antigen). Each antibody can bind to only a specific antigen. The purpose of this binding is to help destroy the antigen. Antibodies can work in several ways, depending on the nature of the antigen. Some antibodies destroy antigens directly. Others make it easier for white blood cells to destroy the antigen. [NIH]

Antigen: Any substance which is capable, under appropriate conditions, of inducing a specific immune response and of reacting with the products of that response, that is, with specific antibody or specifically sensitized T-lymphocytes, or both. Antigens may be soluble substances, such as toxins and foreign proteins, or particulate, such as bacteria and tissue cells; however, only the portion of the protein or polysaccharide molecule known as the antigenic determinant (q.v.) combines with antibody or a specific receptor on a lymphocyte. Abbreviated Ag. [EU]

Antineoplastic: Inhibiting or preventing the development of neoplasms, checking the maturation and proliferation of malignant cells. [EU]

Antioxidant: A substance that prevents damage caused by free radicals. Free radicals are highly reactive chemicals that often contain oxygen. They are produced when molecules are split to give products that have unpaired electrons. This process is called oxidation. [NIH]

Anuria: Inability to form or excrete urine. [NIH]

Anus: The opening of the rectum to the outside of the body. [NIH]

Aorta: The main trunk of the systemic arteries. [NIH]

Aortic Valve: The valve between the left ventricle and the ascending aorta which prevents backflow into the left ventricle. [NIH]

Aortic Valve Stenosis: Constriction in the opening of the aortic valve or of the supra- or subvalvular regions. [NIH]

Apoptosis: One of the two mechanisms by which cell death occurs (the other being the

pathological process of necrosis). Apoptosis is the mechanism responsible for the physiological deletion of cells and appears to be intrinsically programmed. It is characterized by distinctive morphologic changes in the nucleus and cytoplasm, chromatin cleavage at regularly spaced sites, and the endonucleolytic cleavage of genomic DNA (DNA fragmentation) at internucleosomal sites. This mode of cell death serves as a balance to mitosis in regulating the size of animal tissues and in mediating pathologic processes associated with tumor growth. [NIH]

Aqueous: Having to do with water. [NIH]

Arginine: An essential amino acid that is physiologically active in the L-form. [NIH]

Arteries: The vessels carrying blood away from the heart. [NIH]

Arterioles: The smallest divisions of the arteries located between the muscular arteries and the capillaries. [NIH]

Artery: Vessel-carrying blood from the heart to various parts of the body. [NIH]

Arthropathy: Any joint disease. [EU]

Arthroplasty: Surgical reconstruction of a joint to relieve pain or restore motion. [NIH]

Ascorbic Acid: A six carbon compound related to glucose. It is found naturally in citrus fruits and many vegetables. Ascorbic acid is an essential nutrient in human diets, and necessary to maintain connective tissue and bone. Its biologically active form, vitamin C, functions as a reducing agent and coenzyme in several metabolic pathways. Vitamin C is considered an antioxidant. [NIH]

Assay: Determination of the amount of a particular constituent of a mixture, or of the biological or pharmacological potency of a drug. [EU]

Atypical: Irregular; not conformable to the type; in microbiology, applied specifically to strains of unusual type. [EU]

Autopsy: Postmortem examination of the body. [NIH]

Back Pain: Acute or chronic pain located in the posterior regions of the trunk, including the thoracic, lumbar, sacral, or adjacent regions. [NIH]

Bacteria: Unicellular prokaryotic microorganisms which generally possess rigid cell walls, multiply by cell division, and exhibit three principal forms: round or coccid, rodlike or bacillary, and spiral or spirochetal. [NIH]

Bacteriuria: The presence of bacteria in the urine with or without consequent urinary tract infection. Since bacteriuria is a clinical entity, the term does not preclude the use of urine/microbiology for technical discussions on the isolation and segregation of bacteria in the urine. [NIH]

Base: In chemistry, the nonacid part of a salt; a substance that combines with acids to form salts; a substance that dissociates to give hydroxide ions in aqueous solutions; a substance whose molecule or ion can combine with a proton (hydrogen ion); a substance capable of donating a pair of electrons (to an acid) for the formation of a coordinate covalent bond. [EU]

Base Sequence: The sequence of purines and pyrimidines in nucleic acids and polynucleotides. It is also called nucleotide or nucleoside sequence. [NIH]

Bewilderment: Impairment or loss of will power. [NIH]

Bile: An emulsifying agent produced in the liver and secreted into the duodenum. Its composition includes bile acids and salts, cholesterol, and electrolytes. It aids digestion of fats in the duodenum. [NIH]

Biochemical: Relating to biochemistry; characterized by, produced by, or involving chemical reactions in living organisms. [EU]

Biosynthesis: The building up of a chemical compound in the physiologic processes of a living organism. [EU]

Biotechnology: Body of knowledge related to the use of organisms, cells or cell-derived constituents for the purpose of developing products which are technically, scientifically and clinically useful. Alteration of biologic function at the molecular level (i.e., genetic engineering) is a central focus; laboratory methods used include transfection and cloning technologies, sequence and structure analysis algorithms, computer databases, and gene and protein structure function analysis and prediction. [NIH]

Bladder: The organ that stores urine. [NIH]

Blastocyst: The mammalian embryo in the post-morula stage in which a fluid-filled cavity, enclosed primarily by trophoblast, contains an inner cell mass which becomes the embryonic disc. [NIH]

Bleomycin: A complex of related glycopeptide antibiotics from *Streptomyces verticillus* consisting of bleomycin A2 and B2. It inhibits DNA metabolism and is used as an antineoplastic, especially for solid tumors. [NIH]

Blood Glucose: Glucose in blood. [NIH]

Blood pressure: The pressure of blood against the walls of a blood vessel or heart chamber. Unless there is reference to another location, such as the pulmonary artery or one of the heart chambers, it refers to the pressure in the systemic arteries, as measured, for example, in the forearm. [NIH]

Blood vessel: A tube in the body through which blood circulates. Blood vessels include a network of arteries, arterioles, capillaries, venules, and veins. [NIH]

Bone Marrow: The soft tissue filling the cavities of bones. Bone marrow exists in two types, yellow and red. Yellow marrow is found in the large cavities of large bones and consists mostly of fat cells and a few primitive blood cells. Red marrow is a hematopoietic tissue and is the site of production of erythrocytes and granular leukocytes. Bone marrow is made up of a framework of connective tissue containing branching fibers with the frame being filled with marrow cells. [NIH]

Broad-spectrum: Effective against a wide range of microorganisms; said of an antibiotic. [EU]

Bronchi: The larger air passages of the lungs arising from the terminal bifurcation of the trachea. [NIH]

Bronchoscopy: Endoscopic examination, therapy or surgery of the bronchi. [NIH]

Buccal: Pertaining to or directed toward the cheek. In dental anatomy, used to refer to the buccal surface of a tooth. [EU]

Calcium: A basic element found in nearly all organized tissues. It is a member of the alkaline earth family of metals with the atomic symbol Ca, atomic number 20, and atomic weight 40. Calcium is the most abundant mineral in the body and combines with phosphorus to form calcium phosphate in the bones and teeth. It is essential for the normal functioning of nerves and muscles and plays a role in blood coagulation (as factor IV) and in many enzymatic processes. [NIH]

Calcium Oxalate: The calcium salt of oxalic acid, occurring in the urine as crystals and in certain calculi. [NIH]

Cardiovascular: Having to do with the heart and blood vessels. [NIH]

Cardiovascular disease: Any abnormal condition characterized by dysfunction of the heart and blood vessels. CVD includes atherosclerosis (especially coronary heart disease, which can lead to heart attacks), cerebrovascular disease (e.g., stroke), and hypertension (high

blood pressure). [NIH]

Case report: A detailed report of the diagnosis, treatment, and follow-up of an individual patient. Case reports also contain some demographic information about the patient (for example, age, gender, ethnic origin). [NIH]

Catabolism: Any destructive metabolic process by which organisms convert substances into excreted compounds. [EU]

Cataract: An opacity, partial or complete, of one or both eyes, on or in the lens or capsule, especially an opacity impairing vision or causing blindness. The many kinds of cataract are classified by their morphology (size, shape, location) or etiology (cause and time of occurrence). [EU]

Cations: Positively charged atoms, radicals or groups of atoms which travel to the cathode or negative pole during electrolysis. [NIH]

Cause of Death: Factors which produce cessation of all vital bodily functions. They can be analyzed from an epidemiologic viewpoint. [NIH]

Cell: The individual unit that makes up all of the tissues of the body. All living things are made up of one or more cells. [NIH]

Cell Cycle: The complex series of phenomena, occurring between the end of one cell division and the end of the next, by which cellular material is divided between daughter cells. [NIH]

Cell Death: The termination of the cell's ability to carry out vital functions such as metabolism, growth, reproduction, responsiveness, and adaptability. [NIH]

Cell Division: The fission of a cell. [NIH]

Cell Respiration: The metabolic process of all living cells (animal and plant) in which oxygen is used to provide a source of energy for the cell. [NIH]

Centromere: The clear constricted portion of the chromosome at which the chromatids are joined and by which the chromosome is attached to the spindle during cell division. [NIH]

Cephalosporins: A group of broad-spectrum antibiotics first isolated from the Mediterranean fungus *Acremonium* (*Cephalosporium acremonium*). They contain the beta-lactam moiety thia-azabicyclo-octenecarboxylic acid also called 7-aminocephalosporanic acid. [NIH]

Cerebrovascular: Pertaining to the blood vessels of the cerebrum, or brain. [EU]

Cerumen: The yellow or brown waxy secretions produced by vestigial apocrine sweat glands in the external ear canal. [NIH]

Chin: The anatomical frontal portion of the mandible, also known as the mentum, that contains the line of fusion of the two separate halves of the mandible (symphysis menti). This line of fusion divides inferiorly to enclose a triangular area called the mental protuberance. On each side, inferior to the second premolar tooth, is the mental foramen for the passage of blood vessels and a nerve. [NIH]

Cholesterol: The principal sterol of all higher animals, distributed in body tissues, especially the brain and spinal cord, and in animal fats and oils. [NIH]

Chromatin: The material of chromosomes. It is a complex of DNA, histones, and nonhistone proteins (chromosomal proteins, non-histone) found within the nucleus of a cell. [NIH]

Chromosomal: Pertaining to chromosomes. [EU]

Chromosome: Part of a cell that contains genetic information. Except for sperm and eggs, all human cells contain 46 chromosomes. [NIH]

Chromosome Fragility: Susceptibility of chromosomes to breakage and translocation or other aberrations. Chromosome fragile sites are regions that show up in karyotypes as a gap (uncondensed stretch) on the chromatid arm. They are associated with chromosome break sites and other aberrations. A fragile site on the X chromosome is associated with fragile X syndrome. Fragile sites are designated by the letters "FRA" followed by the designation for the specific chromosome and a letter which refers to the different fragile sites on a chromosome (e.g. FRAXA). [NIH]

Chronic: A disease or condition that persists or progresses over a long period of time. [NIH]

Cirrhosis: A type of chronic, progressive liver disease. [NIH]

CIS: Cancer Information Service. The CIS is the National Cancer Institute's link to the public, interpreting and explaining research findings in a clear and understandable manner, and providing personalized responses to specific questions about cancer. Access the CIS by calling 1-800-4-CANCER, or by using the Web site at <http://cis.nci.nih.gov>. [NIH]

Clavulanic Acid: Clavulanic acid (C₈H₉O₅N) and its salts and esters. The acid is a suicide inhibitor of bacterial beta-lactamase enzymes from *Streptomyces clavuligerus*. Administered alone, it has only weak antibacterial activity against most organisms, but given in combination with beta-lactam antibiotics prevents antibiotic inactivation by microbial lactamase. [NIH]

Clinical Medicine: The study and practice of medicine by direct examination of the patient. [NIH]

Clinical trial: A research study that tests how well new medical treatments or other interventions work in people. Each study is designed to test new methods of screening, prevention, diagnosis, or treatment of a disease. [NIH]

Cloning: The production of a number of genetically identical individuals; in genetic engineering, a process for the efficient replication of a great number of identical DNA molecules. [NIH]

Codon: A set of three nucleotides in a protein coding sequence that specifies individual amino acids or a termination signal (codon, terminator). Most codons are universal, but some organisms do not produce the transfer RNAs (RNA, transfer) complementary to all codons. These codons are referred to as unassigned codons (codons, nonsense). [NIH]

Coenzyme: An organic nonprotein molecule, frequently a phosphorylated derivative of a water-soluble vitamin, that binds with the protein molecule (apoenzyme) to form the active enzyme (holoenzyme). [EU]

Colon: The long, coiled, tubelike organ that removes water from digested food. The remaining material, solid waste called stool, moves through the colon to the rectum and leaves the body through the anus. [NIH]

Colonoscopy: Endoscopic examination, therapy or surgery of the luminal surface of the colon. [NIH]

Complement: A term originally used to refer to the heat-labile factor in serum that causes immune cytolysis, the lysis of antibody-coated cells, and now referring to the entire functionally related system comprising at least 20 distinct serum proteins that is the effector not only of immune cytolysis but also of other biologic functions. Complement activation occurs by two different sequences, the classic and alternative pathways. The proteins of the classic pathway are termed 'components of complement' and are designated by the symbols C1 through C9. C1 is a calcium-dependent complex of three distinct proteins C1q, C1r and C1s. The proteins of the alternative pathway (collectively referred to as the properdin system) and complement regulatory proteins are known by semisystematic or trivial names. Fragments resulting from proteolytic cleavage of complement proteins are designated with

lower-case letter suffixes, e.g., C3a. Inactivated fragments may be designated with the suffix 'i', e.g. C3bi. Activated components or complexes with biological activity are designated by a bar over the symbol e.g. C1 or C4b,2a. The classic pathway is activated by the binding of C1 to classic pathway activators, primarily antigen-antibody complexes containing IgM, IgG1, IgG3; C1q binds to a single IgM molecule or two adjacent IgG molecules. The alternative pathway can be activated by IgA immune complexes and also by nonimmunologic materials including bacterial endotoxins, microbial polysaccharides, and cell walls. Activation of the classic pathway triggers an enzymatic cascade involving C1, C4, C2 and C3; activation of the alternative pathway triggers a cascade involving C3 and factors B, D and P. Both result in the cleavage of C5 and the formation of the membrane attack complex. Complement activation also results in the formation of many biologically active complement fragments that act as anaphylatoxins, opsonins, or chemotactic factors. [EU]

Complementary and alternative medicine: CAM. Forms of treatment that are used in addition to (complementary) or instead of (alternative) standard treatments. These practices are not considered standard medical approaches. CAM includes dietary supplements, megadose vitamins, herbal preparations, special teas, massage therapy, magnet therapy, spiritual healing, and meditation. [NIH]

Complementary medicine: Practices not generally recognized by the medical community as standard or conventional medical approaches and used to enhance or complement the standard treatments. Complementary medicine includes the taking of dietary supplements, megadose vitamins, and herbal preparations; the drinking of special teas; and practices such as massage therapy, magnet therapy, spiritual healing, and meditation. [NIH]

Computational Biology: A field of biology concerned with the development of techniques for the collection and manipulation of biological data, and the use of such data to make biological discoveries or predictions. This field encompasses all computational methods and theories applicable to molecular biology and areas of computer-based techniques for solving biological problems including manipulation of models and datasets. [NIH]

Concentric: Having a common center of curvature or symmetry. [NIH]

Conception: The onset of pregnancy, marked by implantation of the blastocyst; the formation of a viable zygote. [EU]

Confusion: A mental state characterized by bewilderment, emotional disturbance, lack of clear thinking, and perceptual disorientation. [NIH]

Connective Tissue: Tissue that supports and binds other tissues. It consists of connective tissue cells embedded in a large amount of extracellular matrix. [NIH]

Connective Tissue: Tissue that supports and binds other tissues. It consists of connective tissue cells embedded in a large amount of extracellular matrix. [NIH]

Connective Tissue Cells: A group of cells that includes fibroblasts, cartilage cells, adipocytes, smooth muscle cells, and bone cells. [NIH]

Consciousness: Sense of awareness of self and of the environment. [NIH]

Constriction: The act of constricting. [NIH]

Consultation: A deliberation between two or more physicians concerning the diagnosis and the proper method of treatment in a case. [NIH]

Contraindications: Any factor or sign that it is unwise to pursue a certain kind of action or treatment, e. g. giving a general anesthetic to a person with pneumonia. [NIH]

Coronary: Encircling in the manner of a crown; a term applied to vessels; nerves, ligaments, etc. The term usually denotes the arteries that supply the heart muscle and, by extension, a pathologic involvement of them. [EU]

Coronary Artery Bypass: Surgical therapy of ischemic coronary artery disease achieved by grafting a section of saphenous vein, internal mammary artery, or other substitute between the aorta and the obstructed coronary artery distal to the obstructive lesion. [NIH]

Coronary heart disease: A type of heart disease caused by narrowing of the coronary arteries that feed the heart, which needs a constant supply of oxygen and nutrients carried by the blood in the coronary arteries. When the coronary arteries become narrowed or clogged by fat and cholesterol deposits and cannot supply enough blood to the heart, CHD results. [NIH]

Cytochrome: Any electron transfer hemoprotein having a mode of action in which the transfer of a single electron is effected by a reversible valence change of the central iron atom of the heme prosthetic group between the +2 and +3 oxidation states; classified as cytochromes a in which the heme contains a formyl side chain, cytochromes b, which contain protoheme or a closely similar heme that is not covalently bound to the protein, cytochromes c in which protoheme or other heme is covalently bound to the protein, and cytochromes d in which the iron-tetrapyrrole has fewer conjugated double bonds than the hemes have. Well-known cytochromes have been numbered consecutively within groups and are designated by subscripts (beginning with no subscript), e.g. cytochromes c, c1, C2, . New cytochromes are named according to the wavelength in nanometres of the absorption maximum of the a-band of the iron (II) form in pyridine, e.g., c-555. [EU]

Cytoplasm: The protoplasm of a cell exclusive of that of the nucleus; it consists of a continuous aqueous solution (cytosol) and the organelles and inclusions suspended in it (phaneroplasm), and is the site of most of the chemical activities of the cell. [EU]

Cytosine: A pyrimidine base that is a fundamental unit of nucleic acids. [NIH]

De novo: In cancer, the first occurrence of cancer in the body. [NIH]

Death Certificates: Official records of individual deaths including the cause of death certified by a physician, and any other required identifying information. [NIH]

Degenerative: Undergoing degeneration : tending to degenerate; having the character of or involving degeneration; causing or tending to cause degeneration. [EU]

Deletion: A genetic rearrangement through loss of segments of DNA (chromosomes), bringing sequences, which are normally separated, into close proximity. [NIH]

Dementia: An acquired organic mental disorder with loss of intellectual abilities of sufficient severity to interfere with social or occupational functioning. The dysfunction is multifaceted and involves memory, behavior, personality, judgment, attention, spatial relations, language, abstract thought, and other executive functions. The intellectual decline is usually progressive, and initially spares the level of consciousness. [NIH]

Deoxyribonucleic: A polymer of subunits called deoxyribonucleotides which is the primary genetic material of a cell, the material equivalent to genetic information. [NIH]

Deoxyribonucleic acid: A polymer of subunits called deoxyribonucleotides which is the primary genetic material of a cell, the material equivalent to genetic information. [NIH]

Deoxyribonucleotides: A purine or pyrimidine base bonded to a deoxyribose containing a bond to a phosphate group. [NIH]

Diabetes Mellitus: A heterogeneous group of disorders that share glucose intolerance in common. [NIH]

Diagnostic procedure: A method used to identify a disease. [NIH]

Digestion: The process of breakdown of food for metabolism and use by the body. [NIH]

Diploid: Having two sets of chromosomes. [NIH]

Direct: 1. Straight; in a straight line. 2. Performed immediately and without the intervention of subsidiary means. [EU]

Discrete: Made up of separate parts or characterized by lesions which do not become blended; not running together; separate. [NIH]

Discrimination: The act of qualitative and/or quantitative differentiation between two or more stimuli. [NIH]

Disorientation: The loss of proper bearings, or a state of mental confusion as to time, place, or identity. [EU]

Distal: Remote; farther from any point of reference; opposed to proximal. In dentistry, used to designate a position on the dental arch farther from the median line of the jaw. [EU]

Dopamine: An endogenous catecholamine and prominent neurotransmitter in several systems of the brain. In the synthesis of catecholamines from tyrosine, it is the immediate precursor to norepinephrine and epinephrine. Dopamine is a major transmitter in the extrapyramidal system of the brain, and important in regulating movement. A family of dopaminergic receptor subtypes mediate its action. Dopamine is used pharmacologically for its direct (beta adrenergic agonist) and indirect (adrenergic releasing) sympathomimetic effects including its actions as an inotropic agent and as a renal vasodilator. [NIH]

Duct: A tube through which body fluids pass. [NIH]

Effector: It is often an enzyme that converts an inactive precursor molecule into an active second messenger. [NIH]

Effector cell: A cell that performs a specific function in response to a stimulus; usually used to describe cells in the immune system. [NIH]

Electrolytes: Substances that break up into ions (electrically charged particles) when they are dissolved in body fluids or water. Some examples are sodium, potassium, chloride, and calcium. Electrolytes are primarily responsible for the movement of nutrients into cells, and the movement of wastes out of cells. [NIH]

Electrons: Stable elementary particles having the smallest known negative charge, present in all elements; also called negatrons. Positively charged electrons are called positrons. The numbers, energies and arrangement of electrons around atomic nuclei determine the chemical identities of elements. Beams of electrons are called cathode rays or beta rays, the latter being a high-energy biproduct of nuclear decay. [NIH]

Embryo: The prenatal stage of mammalian development characterized by rapid morphological changes and the differentiation of basic structures. [NIH]

Environmental Health: The science of controlling or modifying those conditions, influences, or forces surrounding man which relate to promoting, establishing, and maintaining health. [NIH]

Enzyme: A protein that speeds up chemical reactions in the body. [NIH]

Epidermis: Nonvascular layer of the skin. It is made up, from within outward, of five layers: 1) basal layer (stratum basale epidermidis); 2) spinous layer (stratum spinosum epidermidis); 3) granular layer (stratum granulosum epidermidis); 4) clear layer (stratum lucidum epidermidis); and 5) horny layer (stratum corneum epidermidis). [NIH]

Epinephrine: The active sympathomimetic hormone from the adrenal medulla in most species. It stimulates both the alpha- and beta- adrenergic systems, causes systemic vasoconstriction and gastrointestinal relaxation, stimulates the heart, and dilates bronchi and cerebral vessels. It is used in asthma and cardiac failure and to delay absorption of local anesthetics. [NIH]

Erythrocytes: Red blood cells. Mature erythrocytes are non-nucleated, biconcave disks

containing hemoglobin whose function is to transport oxygen. [NIH]

Ethnic Groups: A group of people with a common cultural heritage that sets them apart from others in a variety of social relationships. [NIH]

Eukaryotic Cells: Cells of the higher organisms, containing a true nucleus bounded by a nuclear membrane. [NIH]

Excrete: To get rid of waste from the body. [NIH]

Extracellular: Outside a cell or cells. [EU]

Extracellular Matrix: A meshwork-like substance found within the extracellular space and in association with the basement membrane of the cell surface. It promotes cellular proliferation and provides a supporting structure to which cells or cell lysates in culture dishes adhere. [NIH]

Eye Color: Color of the iris. [NIH]

Eye Infections: Infection, moderate to severe, caused by bacteria, fungi, or viruses, which occurs either on the external surface of the eye or intraocularly with probable inflammation, visual impairment, or blindness. [NIH]

Family Planning: Programs or services designed to assist the family in controlling reproduction by either improving or diminishing fertility. [NIH]

Fat: Total lipids including phospholipids. [NIH]

Fathers: Male parents, human or animal. [NIH]

Fetus: The developing offspring from 7 to 8 weeks after conception until birth. [NIH]

Fibrosis: Any pathological condition where fibrous connective tissue invades any organ, usually as a consequence of inflammation or other injury. [NIH]

Flatus: Gas passed through the rectum. [NIH]

Forearm: The part between the elbow and the wrist. [NIH]

Frameshift: A type of mutation which causes out-of-phase transcription of the base sequence; such mutations arise from the addition or deletion of nucleotide(s) in numbers other than 3 or multiples of 3. [NIH]

Frameshift Mutation: A type of mutation in which a number of nucleotides not divisible by three is deleted from or inserted into a coding sequence, thereby causing an alteration in the reading frame of the entire sequence downstream of the mutation. These mutations may be induced by certain types of mutagens or may occur spontaneously. [NIH]

Fungus: A general term used to denote a group of eukaryotic protists, including mushrooms, yeasts, rusts, moulds, smuts, etc., which are characterized by the absence of chlorophyll and by the presence of a rigid cell wall composed of chitin, mannans, and sometimes cellulose. They are usually of simple morphological form or show some reversible cellular specialization, such as the formation of pseudoparenchymatous tissue in the fruiting body of a mushroom. The dimorphic fungi grow, according to environmental conditions, as moulds or yeasts. [EU]

Ganglia: Clusters of multipolar neurons surrounded by a capsule of loosely organized connective tissue located outside the central nervous system. [NIH]

Gas: Air that comes from normal breakdown of food. The gases are passed out of the body through the rectum (flatus) or the mouth (burp). [NIH]

Gastrin: A hormone released after eating. Gastrin causes the stomach to produce more acid. [NIH]

Gene: The functional and physical unit of heredity passed from parent to offspring. Genes

are pieces of DNA, and most genes contain the information for making a specific protein. [NIH]

Gene Expression: The phenotypic manifestation of a gene or genes by the processes of gene action. [NIH]

Gene Products, rev: Trans-acting nuclear proteins whose functional expression are required for HIV viral replication. Specifically, the rev gene products are required for processing and translation of the HIV gag and env mRNAs, and thus rev regulates the expression of the viral structural proteins. rev can also regulate viral regulatory proteins. A cis-acting antirepression sequence (CAR) in env, also known as the rev-responsive element (RRE), is responsive to the rev gene product. rev is short for regulator of virion. [NIH]

Gene Therapy: The introduction of new genes into cells for the purpose of treating disease by restoring or adding gene expression. Techniques include insertion of retroviral vectors, transfection, homologous recombination, and injection of new genes into the nuclei of single cell embryos. The entire gene therapy process may consist of multiple steps. The new genes may be introduced into proliferating cells in vivo (e.g., bone marrow) or in vitro (e.g., fibroblast cultures) and the modified cells transferred to the site where the gene expression is required. Gene therapy may be particularly useful for treating enzyme deficiency diseases, hemoglobinopathies, and leukemias and may also prove useful in restoring drug sensitivity, particularly for leukemia. [NIH]

Genes, env: DNA sequences that form the coding region for the viral envelope (env) proteins in retroviruses. The env genes contain a cis-acting RNA target sequence for the rev protein (= gene products, rev), termed the rev-responsive element (RRE). [NIH]

Genetic testing: Analyzing DNA to look for a genetic alteration that may indicate an increased risk for developing a specific disease or disorder. [NIH]

Genetics: The biological science that deals with the phenomena and mechanisms of heredity. [NIH]

Genomics: The systematic study of the complete DNA sequences (genome) of organisms. [NIH]

Genotype: The genetic constitution of the individual; the characterization of the genes. [NIH]

Germ Cells: The reproductive cells in multicellular organisms. [NIH]

Germline mutation: A gene change in the body's reproductive cells (egg or sperm) that becomes incorporated into the DNA of every cell in the body of offspring; germline mutations are passed on from parents to offspring. Also called hereditary mutation. [NIH]

Gland: An organ that produces and releases one or more substances for use in the body. Some glands produce fluids that affect tissues or organs. Others produce hormones or participate in blood production. [NIH]

Glomerular: Pertaining to or of the nature of a glomerulus, especially a renal glomerulus. [EU]

Glucose: D-Glucose. A primary source of energy for living organisms. It is naturally occurring and is found in fruits and other parts of plants in its free state. It is used therapeutically in fluid and nutrient replacement. [NIH]

Governing Board: The group in which legal authority is vested for the control of health-related institutions and organizations. [NIH]

Grafting: The operation of transfer of tissue from one site to another. [NIH]

Granule: A small pill made from sucrose. [EU]

Granulocytes: Leukocytes with abundant granules in the cytoplasm. They are divided into three groups: neutrophils, eosinophils, and basophils. [NIH]

Guanine: One of the four DNA bases. [NIH]

Hair Color: Color of hair or fur. [NIH]

Heart attack: A seizure of weak or abnormal functioning of the heart. [NIH]

Hemochromatosis: A disease that occurs when the body absorbs too much iron. The body stores the excess iron in the liver, pancreas, and other organs. May cause cirrhosis of the liver. Also called iron overload disease. [NIH]

Hemodialysis: The use of a machine to clean wastes from the blood after the kidneys have failed. The blood travels through tubes to a dialyzer, which removes wastes and extra fluid. The cleaned blood then flows through another set of tubes back into the body. [NIH]

Hemoglobin: One of the fractions of glycosylated hemoglobin A1c. Glycosylated hemoglobin is formed when linkages of glucose and related monosaccharides bind to hemoglobin A and its concentration represents the average blood glucose level over the previous several weeks. HbA1c levels are used as a measure of long-term control of plasma glucose (normal, 4 to 6 percent). In controlled diabetes mellitus, the concentration of glycosylated hemoglobin A is within the normal range, but in uncontrolled cases the level may be 3 to 4 times the normal concentration. Generally, complications are substantially lower among patients with Hb levels of 7 percent or less than in patients with HbA1c levels of 9 percent or more. [NIH]

Hemoglobinopathies: A group of inherited disorders characterized by structural alterations within the hemoglobin molecule. [NIH]

Hemophilia: Refers to a group of hereditary disorders in which affected individuals fail to make enough of certain proteins needed to form blood clots. [NIH]

Hemorrhage: Bleeding or escape of blood from a vessel. [NIH]

Hepatic: Refers to the liver. [NIH]

Hepatitis: Inflammation of the liver and liver disease involving degenerative or necrotic alterations of hepatocytes. [NIH]

Hepatocytes: The main structural component of the liver. They are specialized epithelial cells that are organized into interconnected plates called lobules. [NIH]

Hereditary: Of, relating to, or denoting factors that can be transmitted genetically from one generation to another. [NIH]

Hereditary mutation: A gene change in the body's reproductive cells (egg or sperm) that becomes incorporated into the DNA of every cell in the body of offspring; hereditary mutations are passed on from parents to offspring. Also called germline mutation. [NIH]

Heredity: 1. The genetic transmission of a particular quality or trait from parent to offspring. 2. The genetic constitution of an individual. [EU]

Heterogeneity: The property of one or more samples or populations which implies that they are not identical in respect of some or all of their parameters, e. g. heterogeneity of variance. [NIH]

Heterozygote: An individual having different alleles at one or more loci in homologous chromosome segments. [NIH]

Histones: Small chromosomal proteins (approx 12-20 kD) possessing an open, unfolded structure and attached to the DNA in cell nuclei by ionic linkages. Classification into the various types (designated histone I, histone II, etc.) is based on the relative amounts of arginine and lysine in each. [NIH]

Homologous: Corresponding in structure, position, origin, etc., as (a) the feathers of a bird and the scales of a fish, (b) antigen and its specific antibody, (c) allelic chromosomes. [EU]

Hormone: A substance in the body that regulates certain organs. Hormones such as gastrin help in breaking down food. Some hormones come from cells in the stomach and small intestine. [NIH]

Hydrogen: The first chemical element in the periodic table. It has the atomic symbol H, atomic number 1, and atomic weight 1. It exists, under normal conditions, as a colorless, odorless, tasteless, diatomic gas. Hydrogen ions are protons. Besides the common H1 isotope, hydrogen exists as the stable isotope deuterium and the unstable, radioactive isotope tritium. [NIH]

Hydroxylation: Hydroxylate, to introduce hydroxyl into (a compound or radical) usually by replacement of hydrogen. [EU]

Hypercalcemia: Abnormally high level of calcium in the blood. [NIH]

Hyperpigmentation: Excessive pigmentation of the skin, usually as a result of increased melanization of the epidermis rather than as a result of an increased number of melanocytes. Etiology is varied and the condition may arise from exposure to light, chemicals or other substances, or from a primary metabolic imbalance. [NIH]

Hypertension: Persistently high arterial blood pressure. Currently accepted threshold levels are 140 mm Hg systolic and 90 mm Hg diastolic pressure. [NIH]

Hypokinesia: Slow or diminished movement of body musculature. It may be associated with basal ganglia diseases; mental disorders; prolonged inactivity due to illness; experimental protocols used to evaluate the physiologic effects of immobility; and other conditions. [NIH]

Immune response: The activity of the immune system against foreign substances (antigens). [NIH]

Immune system: The organs, cells, and molecules responsible for the recognition and disposal of foreign ("non-self") material which enters the body. [NIH]

Impairment: In the context of health experience, an impairment is any loss or abnormality of psychological, physiological, or anatomical structure or function. [NIH]

Implantation: The insertion or grafting into the body of biological, living, inert, or radioactive material. [EU]

In vitro: In the laboratory (outside the body). The opposite of in vivo (in the body). [NIH]

In vivo: In the body. The opposite of in vitro (outside the body or in the laboratory). [NIH]

Individuality: Those psychological characteristics which differentiate individuals from one another. [NIH]

Infancy: The period of complete dependency prior to the acquisition of competence in walking, talking, and self-feeding. [NIH]

Infection: 1. Invasion and multiplication of microorganisms in body tissues, which may be clinically unapparent or result in local cellular injury due to competitive metabolism, toxins, intracellular replication, or antigen-antibody response. The infection may remain localized, subclinical, and temporary if the body's defensive mechanisms are effective. A local infection may persist and spread by extension to become an acute, subacute, or chronic clinical infection or disease state. A local infection may also become systemic when the microorganisms gain access to the lymphatic or vascular system. 2. An infectious disease. [EU]

Inflammation: A pathological process characterized by injury or destruction of tissues caused by a variety of cytologic and chemical reactions. It is usually manifested by typical signs of pain, heat, redness, swelling, and loss of function. [NIH]

Informed Consent: Voluntary authorization, given to the physician by the patient, with full

comprehension of the risks involved, for diagnostic or investigative procedures and medical and surgical treatment. [NIH]

Initiation: Mutation induced by a chemical reactive substance causing cell changes; being a step in a carcinogenic process. [NIH]

Inner ear: The labyrinth, comprising the vestibule, cochlea, and semicircular canals. [NIH]

Insight: The capacity to understand one's own motives, to be aware of one's own psychodynamics, to appreciate the meaning of symbolic behavior. [NIH]

Interstitial: Pertaining to or situated between parts or in the interspaces of a tissue. [EU]

Intravenous: IV. Into a vein. [NIH]

Ion Channels: Gated, ion-selective glycoproteins that traverse membranes. The stimulus for channel gating can be a membrane potential, drug, transmitter, cytoplasmic messenger, or a mechanical deformation. Ion channels which are integral parts of ionotropic neurotransmitter receptors are not included. [NIH]

Ions: An atom or group of atoms that have a positive or negative electric charge due to a gain (negative charge) or loss (positive charge) of one or more electrons. Atoms with a positive charge are known as cations; those with a negative charge are anions. [NIH]

Iris: The most anterior portion of the uveal layer, separating the anterior chamber from the posterior. It consists of two layers - the stroma and the pigmented epithelium. Color of the iris depends on the amount of melanin in the stroma on reflection from the pigmented epithelium. [NIH]

Irradiation: The use of high-energy radiation from x-rays, neutrons, and other sources to kill cancer cells and shrink tumors. Radiation may come from a machine outside the body (external-beam radiation therapy) or from materials called radioisotopes. Radioisotopes produce radiation and can be placed in or near the tumor or in the area near cancer cells. This type of radiation treatment is called internal radiation therapy, implant radiation, interstitial radiation, or brachytherapy. Systemic radiation therapy uses a radioactive substance, such as a radiolabeled monoclonal antibody, that circulates throughout the body. Irradiation is also called radiation therapy, radiotherapy, and x-ray therapy. [NIH]

Karyotype: The characteristic chromosome complement of an individual, race, or species as defined by their number, size, shape, etc. [NIH]

Kidney Failure: The inability of a kidney to excrete metabolites at normal plasma levels under conditions of normal loading, or the inability to retain electrolytes under conditions of normal intake. In the acute form (kidney failure, acute), it is marked by uremia and usually by oliguria or anuria, with hyperkalemia and pulmonary edema. The chronic form (kidney failure, chronic) is irreversible and requires hemodialysis. [NIH]

Kidney Failure, Acute: A clinical syndrome characterized by a sudden decrease in glomerular filtration rate, often to values of less than 1 to 2 ml per minute. It is usually associated with oliguria (urine volumes of less than 400 ml per day) and is always associated with biochemical consequences of the reduction in glomerular filtration rate such as a rise in blood urea nitrogen (BUN) and serum creatinine concentrations. [NIH]

Kidney Failure, Chronic: An irreversible and usually progressive reduction in renal function in which both kidneys have been damaged by a variety of diseases to the extent that they are unable to adequately remove the metabolic products from the blood and regulate the body's electrolyte composition and acid-base balance. Chronic kidney failure requires hemodialysis or surgery, usually kidney transplantation. [NIH]

Kidney stone: A stone that develops from crystals that form in urine and build up on the inner surfaces of the kidney, in the renal pelvis, or in the ureters. [NIH]

Lens: The transparent, double convex (outward curve on both sides) structure suspended between the aqueous and vitreous; helps to focus light on the retina. [NIH]

Lesion: An area of abnormal tissue change. [NIH]

Leucocyte: All the white cells of the blood and their precursors (myeloid cell series, lymphoid cell series) but commonly used to indicate granulocytes exclusive of lymphocytes. [NIH]

Leukemia: Cancer of blood-forming tissue. [NIH]

Ligaments: Shiny, flexible bands of fibrous tissue connecting together articular extremities of bones. They are pliant, tough, and inextensible. [NIH]

Linkages: The tendency of two or more genes in the same chromosome to remain together from one generation to the next more frequently than expected according to the law of independent assortment. [NIH]

Liver: A large, glandular organ located in the upper abdomen. The liver cleanses the blood and aids in digestion by secreting bile. [NIH]

Liver cancer: A disease in which malignant (cancer) cells are found in the tissues of the liver. [NIH]

Liver Transplantation: The transference of a part of or an entire liver from one human or animal to another. [NIH]

Lumbar: Pertaining to the loins, the part of the back between the thorax and the pelvis. [EU]

Lymphocytes: White blood cells formed in the body's lymphoid tissue. The nucleus is round or ovoid with coarse, irregularly clumped chromatin while the cytoplasm is typically pale blue with azurophilic (if any) granules. Most lymphocytes can be classified as either T or B (with subpopulations of each); those with characteristics of neither major class are called null cells. [NIH]

Lysine: An essential amino acid. It is often added to animal feed. [NIH]

Macrophage: A type of white blood cell that surrounds and kills microorganisms, removes dead cells, and stimulates the action of other immune system cells. [NIH]

Malignant: Cancerous; a growth with a tendency to invade and destroy nearby tissue and spread to other parts of the body. [NIH]

Mammary: Pertaining to the mamma, or breast. [EU]

Mammography: Radiographic examination of the breast. [NIH]

Medical Records: Recording of pertinent information concerning patient's illness or illnesses. [NIH]

MEDLINE: An online database of MEDLARS, the computerized bibliographic Medical Literature Analysis and Retrieval System of the National Library of Medicine. [NIH]

Meiosis: A special method of cell division, occurring in maturation of the germ cells, by means of which each daughter nucleus receives half the number of chromosomes characteristic of the somatic cells of the species. [NIH]

Melanin: The substance that gives the skin its color. [NIH]

Melanocytes: Epidermal dendritic pigment cells which control long-term morphological color changes by alteration in their number or in the amount of pigment they produce and store in the pigment containing organelles called melanosomes. Melanophores are larger cells which do not exist in mammals. [NIH]

Melanoma: A form of skin cancer that arises in melanocytes, the cells that produce pigment. Melanoma usually begins in a mole. [NIH]

Membrane: A very thin layer of tissue that covers a surface. [NIH]

Memory: Complex mental function having four distinct phases: (1) memorizing or learning, (2) retention, (3) recall, and (4) recognition. Clinically, it is usually subdivided into immediate, recent, and remote memory. [NIH]

Mental: Pertaining to the mind; psychic. 2. (L. mentum chin) pertaining to the chin. [EU]

Mental Retardation: Refers to sub-average general intellectual functioning which originated during the developmental period and is associated with impairment in adaptive behavior. [NIH]

Microbe: An organism which cannot be observed with the naked eye; e. g. unicellular animals, lower algae, lower fungi, bacteria. [NIH]

Microbiology: The study of microorganisms such as fungi, bacteria, algae, archaea, and viruses. [NIH]

Microorganism: An organism that can be seen only through a microscope. Microorganisms include bacteria, protozoa, algae, and fungi. Although viruses are not considered living organisms, they are sometimes classified as microorganisms. [NIH]

Miscarriage: Spontaneous expulsion of the products of pregnancy before the middle of the second trimester. [NIH]

Mitochondria: Parts of a cell where aerobic production (also known as cell respiration) takes place. [NIH]

Mitosis: A method of indirect cell division by means of which the two daughter nuclei normally receive identical complements of the number of chromosomes of the somatic cells of the species. [NIH]

Molecular: Of, pertaining to, or composed of molecules : a very small mass of matter. [EU]

Molecule: A chemical made up of two or more atoms. The atoms in a molecule can be the same (an oxygen molecule has two oxygen atoms) or different (a water molecule has two hydrogen atoms and one oxygen atom). Biological molecules, such as proteins and DNA, can be made up of many thousands of atoms. [NIH]

Monitor: An apparatus which automatically records such physiological signs as respiration, pulse, and blood pressure in an anesthetized patient or one undergoing surgical or other procedures. [NIH]

Monosomy: The condition in which one chromosome of a pair is missing. In a normally diploid cell it is represented symbolically as $2N-1$. [NIH]

Morphological: Relating to the configuration or the structure of live organs. [NIH]

Morphology: The science of the form and structure of organisms (plants, animals, and other forms of life). [NIH]

Mosaicism: The occurrence in an individual of two or more cell populations of different chromosomal constitutions, derived from a single zygote, as opposed to chimerism in which the different cell populations are derived from more than one zygote. [NIH]

Mutagens: Chemical agents that increase the rate of genetic mutation by interfering with the function of nucleic acids. A clastogen is a specific mutagen that causes breaks in chromosomes. [NIH]

Myotonic Dystrophy: A condition presenting muscle weakness and wasting which may be progressive. [NIH]

NCI: National Cancer Institute. NCI, part of the National Institutes of Health of the United States Department of Health and Human Services, is the federal government's principal agency for cancer research. NCI conducts, coordinates, and funds cancer research, training,

health information dissemination, and other programs with respect to the cause, diagnosis, prevention, and treatment of cancer. Access the NCI Web site at <http://cancer.gov>. [NIH]

Necrosis: A pathological process caused by the progressive degradative action of enzymes that is generally associated with severe cellular trauma. It is characterized by mitochondrial swelling, nuclear flocculation, uncontrolled cell lysis, and ultimately cell death. [NIH]

Neonatal: Pertaining to the first four weeks after birth. [EU]

Nephropathy: Disease of the kidneys. [EU]

Nervous System: The entire nerve apparatus composed of the brain, spinal cord, nerves and ganglia. [NIH]

Neuroblastoma: Cancer that arises in immature nerve cells and affects mostly infants and children. [NIH]

Neurons: The basic cellular units of nervous tissue. Each neuron consists of a body, an axon, and dendrites. Their purpose is to receive, conduct, and transmit impulses in the nervous system. [NIH]

Neuropathy: A problem in any part of the nervous system except the brain and spinal cord. Neuropathies can be caused by infection, toxic substances, or disease. [NIH]

Neurotransmitter: Any of a group of substances that are released on excitation from the axon terminal of a presynaptic neuron of the central or peripheral nervous system and travel across the synaptic cleft to either excite or inhibit the target cell. Among the many substances that have the properties of a neurotransmitter are acetylcholine, norepinephrine, epinephrine, dopamine, glycine, γ -aminobutyrate, glutamic acid, substance P, enkephalins, endorphins, and serotonin. [EU]

Nuclear: A test of the structure, blood flow, and function of the kidneys. The doctor injects a mildly radioactive solution into an arm vein and uses x-rays to monitor its progress through the kidneys. [NIH]

Nuclear Envelope: The membrane system of the cell nucleus that surrounds the nucleoplasm. It consists of two concentric membranes separated by the perinuclear space. The structures of the envelope where it opens to the cytoplasm are called the nuclear pores (nuclear pore). [NIH]

Nuclear Pore: An opening through the nuclear envelope formed by the nuclear pore complex which transports nuclear proteins or RNA into or out of the cell nucleus and which, under some conditions, acts as an ion channel. [NIH]

Nuclei: A body of specialized protoplasm found in nearly all cells and containing the chromosomes. [NIH]

Nucleic acid: Either of two types of macromolecule (DNA or RNA) formed by polymerization of nucleotides. Nucleic acids are found in all living cells and contain the information (genetic code) for the transfer of genetic information from one generation to the next. [NIH]

Nucleus: A body of specialized protoplasm found in nearly all cells and containing the chromosomes. [NIH]

Nurse Practitioners: Nurses who are specially trained to assume an expanded role in providing medical care under the supervision of a physician. [NIH]

Ochronosis: Deposition of polymerized homogentisic acid as a brown-black pigment in the connective tissue. It occurs in alkaptonuria, but has also been observed in connection with exposure to certain chemicals (e.g., phenol, trinitrophenol, benzene derivatives). [NIH]

Oliguria: Clinical manifestation of the urinary system consisting of a decrease in the amount

of urine secreted. [NIH]

Opacity: Degree of density (area most dense taken for reading). [NIH]

Organelles: Specific particles of membrane-bound organized living substances present in eukaryotic cells, such as the mitochondria; the golgi apparatus; endoplasmic reticulum; lysosomes; plastids; and vacuoles. [NIH]

Ovaries: The pair of female reproductive glands in which the ova, or eggs, are formed. The ovaries are located in the pelvis, one on each side of the uterus. [NIH]

Oxalate: A chemical that combines with calcium in urine to form the most common type of kidney stone (calcium oxalate stone). [NIH]

Oxidation: The act of oxidizing or state of being oxidized. Chemically it consists in the increase of positive charges on an atom or the loss of negative charges. Most biological oxidations are accomplished by the removal of a pair of hydrogen atoms (dehydrogenation) from a molecule. Such oxidations must be accompanied by reduction of an acceptor molecule. Univalent o. indicates loss of one electron; divalent o., the loss of two electrons. [EU]

Oxidative Phosphorylation: Electron transfer through the cytochrome system liberating free energy which is transformed into high-energy phosphate bonds. [NIH]

Pancreas: A mixed exocrine and endocrine gland situated transversely across the posterior abdominal wall in the epigastric and hypochondriac regions. The endocrine portion is comprised of the Islets of Langerhans, while the exocrine portion is a compound acinar gland that secretes digestive enzymes. [NIH]

Parkinsonism: A group of neurological disorders characterized by hypokinesia, tremor, and muscular rigidity. [EU]

Paternity: Establishing the father relationship of a man and a child. [NIH]

Pathologic: 1. Indicative of or caused by a morbid condition. 2. Pertaining to pathology (= branch of medicine that treats the essential nature of the disease, especially the structural and functional changes in tissues and organs of the body caused by the disease). [EU]

Pathologic Processes: The abnormal mechanisms and forms involved in the dysfunctions of tissues and organs. [NIH]

Pathophysiology: Altered functions in an individual or an organ due to disease. [NIH]

PDQ: Physician Data Query. PDQ is an online database developed and maintained by the National Cancer Institute. Designed to make the most current, credible, and accurate cancer information available to health professionals and the public, PDQ contains peer-reviewed summaries on cancer treatment, screening, prevention, genetics, and supportive care; a registry of cancer clinical trials from around the world; and directories of physicians, professionals who provide genetics services, and organizations that provide cancer care. Most of this information is available on the CancerNet Web site, and more specific information about PDQ can be found at <http://cancer.net.nci.nih.gov/pdq.html>. [NIH]

Pelvis: The lower part of the abdomen, located between the hip bones. [NIH]

Penicillin: An antibiotic drug used to treat infection. [NIH]

Peptide: Any compound consisting of two or more amino acids, the building blocks of proteins. Peptides are combined to make proteins. [NIH]

Peroxidase: A hemeprotein from leukocytes. Deficiency of this enzyme leads to a hereditary disorder coupled with disseminated monilia. It catalyzes the conversion of a donor and peroxide to an oxidized donor and water. EC 1.11.1.7. [NIH]

Peroxide: Chemical compound which contains an atom group with two oxygen atoms tied

to each other. [NIH]

Pharmacologic: Pertaining to pharmacology or to the properties and reactions of drugs. [EU]

Phenotype: The outward appearance of the individual. It is the product of interactions between genes and between the genotype and the environment. This includes the killer phenotype, characteristic of yeasts. [NIH]

Phenylalanine: An aromatic amino acid that is essential in the animal diet. It is a precursor of melanin, dopamine, noradrenalin, and thyroxine. [NIH]

Phosphorus: A non-metallic element that is found in the blood, muscles, nerves, bones, and teeth, and is a component of adenosine triphosphate (ATP; the primary energy source for the body's cells.) [NIH]

Phosphorylation: The introduction of a phosphoryl group into a compound through the formation of an ester bond between the compound and a phosphorus moiety. [NIH]

Physical Examination: Systematic and thorough inspection of the patient for physical signs of disease or abnormality. [NIH]

Physiologic: Having to do with the functions of the body. When used in the phrase "physiologic age," it refers to an age assigned by general health, as opposed to calendar age. [NIH]

Pigment: A substance that gives color to tissue. Pigments are responsible for the color of skin, eyes, and hair. [NIH]

Pigmentation: Coloration or discoloration of a part by a pigment. [NIH]

Plasma: The clear, yellowish, fluid part of the blood that carries the blood cells. The proteins that form blood clots are in plasma. [NIH]

Plastids: Self-replicating cytoplasmic organelles of plant and algal cells that contain pigments and may synthesize and accumulate various substances. Plastids are used in phylogenetic studies. [NIH]

Pneumonia: Inflammation of the lungs. [NIH]

Point Mutation: A mutation caused by the substitution of one nucleotide for another. This results in the DNA molecule having a change in a single base pair. [NIH]

Polymorphic: Occurring in several or many forms; appearing in different forms at different stages of development. [EU]

Polymorphism: The occurrence together of two or more distinct forms in the same population. [NIH]

Posterior: Situated in back of, or in the back part of, or affecting the back or dorsal surface of the body. In lower animals, it refers to the caudal end of the body. [EU]

Practice Guidelines: Directions or principles presenting current or future rules of policy for the health care practitioner to assist him in patient care decisions regarding diagnosis, therapy, or related clinical circumstances. The guidelines may be developed by government agencies at any level, institutions, professional societies, governing boards, or by the convening of expert panels. The guidelines form a basis for the evaluation of all aspects of health care and delivery. [NIH]

Precursor: Something that precedes. In biological processes, a substance from which another, usually more active or mature substance is formed. In clinical medicine, a sign or symptom that heralds another. [EU]

Prenatal: Existing or occurring before birth, with reference to the fetus. [EU]

Prevalence: The total number of cases of a given disease in a specified population at a

designated time. It is differentiated from incidence, which refers to the number of new cases in the population at a given time. [NIH]

Progressive: Advancing; going forward; going from bad to worse; increasing in scope or severity. [EU]

Prolapse: The protrusion of an organ or part of an organ into a natural or artificial orifice. [NIH]

Prone: Having the front portion of the body downwards. [NIH]

Protein C: A vitamin-K dependent zymogen present in the blood, which, upon activation by thrombin and thrombomodulin exerts anticoagulant properties by inactivating factors Va and VIIIa at the rate-limiting steps of thrombin formation. [NIH]

Protein S: The vitamin K-dependent cofactor of activated protein C. Together with protein C, it inhibits the action of factors VIIIa and Va. A deficiency in protein S can lead to recurrent venous and arterial thrombosis. [NIH]

Proteins: Polymers of amino acids linked by peptide bonds. The specific sequence of amino acids determines the shape and function of the protein. [NIH]

Proteolytic: 1. Pertaining to, characterized by, or promoting proteolysis. 2. An enzyme that promotes proteolysis (= the splitting of proteins by hydrolysis of the peptide bonds with formation of smaller polypeptides). [EU]

Protocol: The detailed plan for a clinical trial that states the trial's rationale, purpose, drug or vaccine dosages, length of study, routes of administration, who may participate, and other aspects of trial design. [NIH]

Psychic: Pertaining to the psyche or to the mind; mental. [EU]

Public Policy: A course or method of action selected, usually by a government, from among alternatives to guide and determine present and future decisions. [NIH]

Publishing: "The business or profession of the commercial production and issuance of literature" (Webster's 3d). It includes the publisher, publication processes, editing and editors. Production may be by conventional printing methods or by electronic publishing. [NIH]

Pulmonary: Relating to the lungs. [NIH]

Pulmonary Artery: The short wide vessel arising from the conus arteriosus of the right ventricle and conveying unaerated blood to the lungs. [NIH]

Pulmonary Edema: An accumulation of an excessive amount of watery fluid in the lungs, may be caused by acute exposure to dangerous concentrations of irritant gasses. [NIH]

Purines: A series of heterocyclic compounds that are variously substituted in nature and are known also as purine bases. They include adenine and guanine, constituents of nucleic acids, as well as many alkaloids such as caffeine and theophylline. Uric acid is the metabolic end product of purine metabolism. [NIH]

Pyrimidines: A family of 6-membered heterocyclic compounds occurring in nature in a wide variety of forms. They include several nucleic acid constituents (cytosine, thymine, and uracil) and form the basic structure of the barbiturates. [NIH]

Race: A population within a species which exhibits general similarities within itself, but is both discontinuous and distinct from other populations of that species, though not sufficiently so as to achieve the status of a taxon. [NIH]

Radiation: Emission or propagation of electromagnetic energy (waves/rays), or the waves/rays themselves; a stream of electromagnetic particles (electrons, neutrons, protons, alpha particles) or a mixture of these. The most common source is the sun. [NIH]

Radiation therapy: The use of high-energy radiation from x-rays, gamma rays, neutrons, and other sources to kill cancer cells and shrink tumors. Radiation may come from a machine outside the body (external-beam radiation therapy), or it may come from radioactive material placed in the body in the area near cancer cells (internal radiation therapy, implant radiation, or brachytherapy). Systemic radiation therapy uses a radioactive substance, such as a radiolabeled monoclonal antibody, that circulates throughout the body. Also called radiotherapy. [NIH]

Radioactive: Giving off radiation. [NIH]

Receptor: A molecule inside or on the surface of a cell that binds to a specific substance and causes a specific physiologic effect in the cell. [NIH]

Recombinant: A cell or an individual with a new combination of genes not found together in either parent; usually applied to linked genes. [EU]

Recombination: The formation of new combinations of genes as a result of segregation in crosses between genetically different parents; also the rearrangement of linked genes due to crossing-over. [NIH]

Rectum: The last 8 to 10 inches of the large intestine. [NIH]

Refer: To send or direct for treatment, aid, information, de decision. [NIH]

Regurgitation: A backward flowing, as the casting up of undigested food, or the backward flowing of blood into the heart, or between the chambers of the heart when a valve is incompetent. [EU]

Renal failure: Progressive renal insufficiency and uremia, due to irreversible and progressive renal glomerular tubular or interstitial disease. [NIH]

Renal tubular: A defect in the kidneys that hinders their normal excretion of acids. Failure to excrete acids can lead to weak bones, kidney stones, and poor growth in children. [NIH]

Repopulation: The replacement of functional cells, usually by proliferation, following or during irradiation. [NIH]

Reproductive cells: Egg and sperm cells. Each mature reproductive cell carries a single set of 23 chromosomes. [NIH]

Retinoblastoma: An eye cancer that most often occurs in children younger than 5 years. It occurs in hereditary and nonhereditary (sporadic) forms. [NIH]

Retroviral vector: RNA from a virus that is used to insert genetic material into cells. [NIH]

Ribonucleic acid: RNA. One of the two nucleic acids found in all cells. The other is deoxyribonucleic acid (DNA). Ribonucleic acid transfers genetic information from DNA to proteins produced by the cell. [NIH]

Ribose: A pentose active in biological systems usually in its D-form. [NIH]

Ribosome: A granule of protein and RNA, synthesized in the nucleolus and found in the cytoplasm of cells. Ribosomes are the main sites of protein synthesis. Messenger RNA attaches to them and there receives molecules of transfer RNA bearing amino acids. [NIH]

Rigidity: Stiffness or inflexibility, chiefly that which is abnormal or morbid; rigor. [EU]

Ristocetin: An antibiotic mixture of two components, A and B, obtained from *Nocardia lurida* (or the same substance produced by any other means). It is no longer used clinically because of its toxicity. It causes platelet agglutination and blood coagulation and is used to assay those functions in vitro. [NIH]

Saphenous: Applied to certain structures in the leg, e. g. nerve vein. [NIH]

Saphenous Vein: The vein which drains the foot and leg. [NIH]

Scatter: The extent to which relative success and failure are divergently manifested in qualitatively different tests. [NIH]

Schizophrenia: A mental disorder characterized by a special type of disintegration of the personality. [NIH]

Sclerosis: A pathological process consisting of hardening or fibrosis of an anatomical structure, often a vessel or a nerve. [NIH]

Screening: Checking for disease when there are no symptoms. [NIH]

Second Messenger Systems: Systems in which an intracellular signal is generated in response to an intercellular primary messenger such as a hormone or neurotransmitter. They are intermediate signals in cellular processes such as metabolism, secretion, contraction, phototransduction, and cell growth. Examples of second messenger systems are the adenylyl cyclase-cyclic AMP system, the phosphatidylinositol diphosphate-inositol triphosphate system, and the cyclic GMP system. [NIH]

Sediment: A precipitate, especially one that is formed spontaneously. [EU]

Sequencing: The determination of the order of nucleotides in a DNA or RNA chain. [NIH]

Side effect: A consequence other than the one(s) for which an agent or measure is used, as the adverse effects produced by a drug, especially on a tissue or organ system other than the one sought to be benefited by its administration. [EU]

Signs and Symptoms: Clinical manifestations that can be either objective when observed by a physician, or subjective when perceived by the patient. [NIH]

Skin Pigmentation: Coloration of the skin. [NIH]

Small intestine: The part of the digestive tract that is located between the stomach and the large intestine. [NIH]

Social Work: The use of community resources, individual case work, or group work to promote the adaptive capacities of individuals in relation to their social and economic environments. It includes social service agencies. [NIH]

Soft tissue: Refers to muscle, fat, fibrous tissue, blood vessels, or other supporting tissue of the body. [NIH]

Solid tumor: Cancer of body tissues other than blood, bone marrow, or the lymphatic system. [NIH]

Soma: The body as distinct from the mind; all the body tissue except the germ cells; all the axial body. [NIH]

Somatic: 1. Pertaining to or characteristic of the soma or body. 2. Pertaining to the body wall in contrast to the viscera. [EU]

Somatic cells: All the body cells except the reproductive (germ) cells. [NIH]

Somatic mutations: Alterations in DNA that occur after conception. Somatic mutations can occur in any of the cells of the body except the germ cells (sperm and egg) and therefore are not passed on to children. These alterations can (but do not always) cause cancer or other diseases. [NIH]

Specialist: In medicine, one who concentrates on 1 special branch of medical science. [NIH]

Species: A taxonomic category subordinate to a genus (or subgenus) and superior to a subspecies or variety, composed of individuals possessing common characters distinguishing them from other categories of individuals of the same taxonomic level. In taxonomic nomenclature, species are designated by the genus name followed by a Latin or Latinized adjective or noun. [EU]

Spectroscopic: The recognition of elements through their emission spectra. [NIH]

Sperm: The fecundating fluid of the male. [NIH]

Spinal cord: The main trunk or bundle of nerves running down the spine through holes in the spinal bone (the vertebrae) from the brain to the level of the lower back. [NIH]

Sporadic: Neither endemic nor epidemic; occurring occasionally in a random or isolated manner. [EU]

Stenosis: Narrowing or stricture of a duct or canal. [EU]

Stillbirth: The birth of a dead fetus or baby. [NIH]

Stimulant: 1. Producing stimulation; especially producing stimulation by causing tension on muscle fibre through the nervous tissue. 2. An agent or remedy that produces stimulation. [EU]

Stomach: An organ of digestion situated in the left upper quadrant of the abdomen between the termination of the esophagus and the beginning of the duodenum. [NIH]

Stool: The waste matter discharged in a bowel movement; feces. [NIH]

Strand: DNA normally exists in the bacterial nucleus in a helix, in which two strands are coiled together. [NIH]

Stricture: The abnormal narrowing of a body opening. Also called stenosis. [NIH]

Stroke: Sudden loss of function of part of the brain because of loss of blood flow. Stroke may be caused by a clot (thrombosis) or rupture (hemorrhage) of a blood vessel to the brain. [NIH]

Subspecies: A category intermediate in rank between species and variety, based on a smaller number of correlated characters than are used to differentiate species and generally conditioned by geographical and/or ecological occurrence. [NIH]

Supportive care: Treatment given to prevent, control, or relieve complications and side effects and to improve the comfort and quality of life of people who have cancer. [NIH]

Sweat: The fluid excreted by the sweat glands. It consists of water containing sodium chloride, phosphate, urea, ammonia, and other waste products. [NIH]

Sweat Glands: Sweat-producing structures that are embedded in the dermis. Each gland consists of a single tube, a coiled body, and a superficial duct. [NIH]

Synapses: Specialized junctions at which a neuron communicates with a target cell. At classical synapses, a neuron's presynaptic terminal releases a chemical transmitter stored in synaptic vesicles which diffuses across a narrow synaptic cleft and activates receptors on the postsynaptic membrane of the target cell. The target may be a dendrite, cell body, or axon of another neuron, or a specialized region of a muscle or secretory cell. Neurons may also communicate through direct electrical connections which are sometimes called electrical synapses; these are not included here but rather in gap junctions. [NIH]

Systemic: Affecting the entire body. [NIH]

Tendon: A discrete band of connective tissue mainly composed of parallel bundles of collagenous fibers by which muscles are attached, or two muscles bellies joined. [NIH]

Terminator: A DNA sequence sited at the end of a transcriptional unit that signals the end of transcription. [NIH]

Thoracic: Having to do with the chest. [NIH]

Thrombosis: The formation or presence of a blood clot inside a blood vessel. [NIH]

Thyroid: A gland located near the windpipe (trachea) that produces thyroid hormone, which helps regulate growth and metabolism. [NIH]

Thyroid Gland: A highly vascular endocrine gland consisting of two lobes, one on either side of the trachea, joined by a narrow isthmus; it produces the thyroid hormones which are concerned in regulating the metabolic rate of the body. [NIH]

Thyroid Hormones: Hormones secreted by the thyroid gland. [NIH]

Thyroxine: An amino acid of the thyroid gland which exerts a stimulating effect on thyroid metabolism. [NIH]

Tissue: A group or layer of cells that are alike in type and work together to perform a specific function. [NIH]

Tone: 1. The normal degree of vigour and tension; in muscle, the resistance to passive elongation or stretch; tonus. 2. A particular quality of sound or of voice. 3. To make permanent, or to change, the colour of silver stain by chemical treatment, usually with a heavy metal. [EU]

Toxic: Having to do with poison or something harmful to the body. Toxic substances usually cause unwanted side effects. [NIH]

Toxicity: The quality of being poisonous, especially the degree of virulence of a toxic microbe or of a poison. [EU]

Toxicology: The science concerned with the detection, chemical composition, and pharmacologic action of toxic substances or poisons and the treatment and prevention of toxic manifestations. [NIH]

Trachea: The cartilaginous and membranous tube descending from the larynx and branching into the right and left main bronchi. [NIH]

Transcription Factors: Endogenous substances, usually proteins, which are effective in the initiation, stimulation, or termination of the genetic transcription process. [NIH]

Transfection: The uptake of naked or purified DNA into cells, usually eukaryotic. It is analogous to bacterial transformation. [NIH]

Translation: The process whereby the genetic information present in the linear sequence of ribonucleotides in mRNA is converted into a corresponding sequence of amino acids in a protein. It occurs on the ribosome and is unidirectional. [NIH]

Transplantation: Transference of a tissue or organ, alive or dead, within an individual, between individuals of the same species, or between individuals of different species. [NIH]

Tremor: Cyclical movement of a body part that can represent either a physiologic process or a manifestation of disease. Intention or action tremor, a common manifestation of cerebellar diseases, is aggravated by movement. In contrast, resting tremor is maximal when there is no attempt at voluntary movement, and occurs as a relatively frequent manifestation of Parkinson disease. [NIH]

Trinucleotide Repeat Expansion: DNA region comprised of a variable number of repetitive, contiguous trinucleotide sequences. The presence of these regions is associated with diseases such as Fragile X Syndrome and myotonic dystrophy. Many chromosome fragile sites (chromosome fragility) contain expanded trinucleotide repeats. [NIH]

Trinucleotide Repeats: Microsatellite repeats consisting of three nucleotides dispersed in the euchromatic arms of chromosomes. [NIH]

Trisomy: The possession of a third chromosome of any one type in an otherwise diploid cell. [NIH]

Tyrosine: A non-essential amino acid. In animals it is synthesized from phenylalanine. It is also the precursor of epinephrine, thyroid hormones, and melanin. [NIH]

Ultraviolet radiation: Invisible rays that are part of the energy that comes from the sun. UV

radiation can damage the skin and cause melanoma and other types of skin cancer. UV radiation that reaches the earth's surface is made up of two types of rays, called UVA and UVB rays. UVB rays are more likely than UVA rays to cause sunburn, but UVA rays pass deeper into the skin. Scientists have long thought that UVB radiation can cause melanoma and other types of skin cancer. They now think that UVA radiation also may add to skin damage that can lead to skin cancer and cause premature aging. For this reason, skin specialists recommend that people use sunscreens that reflect, absorb, or scatter both kinds of UV radiation. [NIH]

Uremia: The illness associated with the buildup of urea in the blood because the kidneys are not working effectively. Symptoms include nausea, vomiting, loss of appetite, weakness, and mental confusion. [NIH]

Urethra: The tube through which urine leaves the body. It empties urine from the bladder. [NIH]

Uric: A kidney stone that may result from a diet high in animal protein. When the body breaks down this protein, uric acid levels rise and can form stones. [NIH]

Urinalysis: Examination of urine by chemical, physical, or microscopic means. Routine urinalysis usually includes performing chemical screening tests, determining specific gravity, observing any unusual color or odor, screening for bacteriuria, and examining the sediment microscopically. [NIH]

Urinary: Having to do with urine or the organs of the body that produce and get rid of urine. [NIH]

Urinary tract: The organs of the body that produce and discharge urine. These include the kidneys, ureters, bladder, and urethra. [NIH]

Urine: Fluid containing water and waste products. Urine is made by the kidneys, stored in the bladder, and leaves the body through the urethra. [NIH]

Urography: Radiography of any part of the urinary tract. [NIH]

Urolithiasis: Stones in the urinary system. [NIH]

Uterus: The small, hollow, pear-shaped organ in a woman's pelvis. This is the organ in which a fetus develops. Also called the womb. [NIH]

Vaccine: A substance or group of substances meant to cause the immune system to respond to a tumor or to microorganisms, such as bacteria or viruses. [NIH]

Vacuoles: Any spaces or cavities within a cell. They may function in digestion, storage, secretion, or excretion. [NIH]

Valine: A branched-chain essential amino acid that has stimulant activity. It promotes muscle growth and tissue repair. It is a precursor in the penicillin biosynthetic pathway. [NIH]

Vancomycin: Antibacterial obtained from *Streptomyces orientalis*. It is a glycopeptide related to ristocetin that inhibits bacterial cell wall assembly and is toxic to kidneys and the inner ear. [NIH]

Vascular: Pertaining to blood vessels or indicative of a copious blood supply. [EU]

Vector: Plasmid or other self-replicating DNA molecule that transfers DNA between cells in nature or in recombinant DNA technology. [NIH]

Vein: Vessel-carrying blood from various parts of the body to the heart. [NIH]

Ventricle: One of the two pumping chambers of the heart. The right ventricle receives oxygen-poor blood from the right atrium and pumps it to the lungs through the pulmonary artery. The left ventricle receives oxygen-rich blood from the left atrium and pumps it to the

body through the aorta. [NIH]

Venules: The minute vessels that collect blood from the capillary plexuses and join together to form veins. [NIH]

Veterinary Medicine: The medical science concerned with the prevention, diagnosis, and treatment of diseases in animals. [NIH]

Viral: Pertaining to, caused by, or of the nature of virus. [EU]

Virulence: The degree of pathogenicity within a group or species of microorganisms or viruses as indicated by case fatality rates and/or the ability of the organism to invade the tissues of the host. [NIH]

Virus: Submicroscopic organism that causes infectious disease. In cancer therapy, some viruses may be made into vaccines that help the body build an immune response to, and kill, tumor cells. [NIH]

Viscera: Any of the large interior organs in any one of the three great cavities of the body, especially in the abdomen. [NIH]

Vitro: Descriptive of an event or enzyme reaction under experimental investigation occurring outside a living organism. Parts of an organism or microorganism are used together with artificial substrates and/or conditions. [NIH]

Vivo: Outside of or removed from the body of a living organism. [NIH]

White blood cell: A type of cell in the immune system that helps the body fight infection and disease. White blood cells include lymphocytes, granulocytes, macrophages, and others. [NIH]

Windpipe: A rigid tube, 10 cm long, extending from the cricoid cartilage to the upper border of the fifth thoracic vertebra. [NIH]

Womb: A hollow, thick-walled, muscular organ in which the impregnated ovum is developed into a child. [NIH]

X-ray: High-energy radiation used in low doses to diagnose diseases and in high doses to treat cancer. [NIH]

Yeasts: A general term for single-celled rounded fungi that reproduce by budding. Brewers' and bakers' yeasts are *Saccharomyces cerevisiae*; therapeutic dried yeast is dried yeast. [NIH]

Zygote: The fertilized ovum. [NIH]

INDEX

3

3-dimensional, 41, 72, 84

A

Abdomen, 84, 98, 101, 106, 109
 Acceptor, 84, 101
 Acremonium, 84, 88
 Adaptability, 84, 88
 Adenine, 35, 84, 103
 Adenosine, 36, 84, 102
 Adenosine Triphosphate, 36, 84, 102
 Adenovirus, 68, 84
 Adverse Effect, 84, 105
 Aerobic, 84, 99
 Algorithms, 84, 87
 Alleles, 37, 54, 84, 95
 Alternative medicine, 84
 Amino Acids, 4, 6, 37, 41, 47, 84, 85, 89, 101, 103, 104, 107
 Amnion, 85
 Amniotic Fluid, 63, 65, 85
 Anatomical, 85, 88, 96, 105
 Anemia, 49, 50, 53, 54, 59, 85
 Aneuploidy, 48, 85
 Anions, 85, 97
 Antibacterial, 85, 89, 108
 Antibiotic, 85, 87, 89, 101, 104
 Antibody, 42, 85, 89, 95, 96, 97, 104
 Antigen, 85, 90, 95, 96
 Antineoplastic, 85, 87
 Antioxidant, 85, 86
 Anuria, 85, 97
 Anus, 85, 89
 Aorta, 28, 85, 91, 109
 Aortic Valve, 25, 85
 Aortic Valve Stenosis, 25, 85
 Apoptosis, 36, 45, 85
 Aqueous, 86, 91, 98
 Arginine, 86, 95
 Arteries, 85, 86, 87, 90, 91
 Arterioles, 86, 87
 Artery, 20, 86, 91
 Arthropathy, 5, 10, 12, 15, 22, 26, 86
 Arthroplasty, 20, 86
 Ascorbic Acid, 22, 27, 86
 Assay, 23, 86, 104
 Atypical, 14, 58, 86
 Autopsy, 19, 86

B

Back Pain, 15, 86

Bacteria, 34, 42, 46, 85, 86, 93, 99, 108
 Bacteriuria, 86, 108
 Base, 6, 7, 35, 36, 39, 41, 45, 46, 47, 70, 84, 86, 91, 93, 97, 102
 Base Sequence, 46, 86, 93
 Bewilderment, 86, 90
 Bile, 86, 98
 Biochemical, 50, 84, 86, 97
 Biosynthesis, 9, 87
 Biotechnology, 9, 41, 68, 70, 75, 87
 Bladder, 87, 108
 Blastocyst, 87, 90
 Bleomycin, 9, 87
 Blood Glucose, 87, 95
 Blood pressure, 53, 87, 88, 96, 99
 Blood vessel, 57, 87, 88, 105, 106, 108
 Bone Marrow, 69, 87, 94, 105
 Broad-spectrum, 87, 88
 Bronchi, 87, 92, 107
 Bronchoscopy, 13, 87
 Buccal, 63, 65, 87

C

Calcium, 87, 89, 92, 96, 101
 Calcium Oxalate, 87, 101
 Cardiovascular, 20, 72, 87
 Cardiovascular disease, 72, 87
 Case report, 11, 12, 14, 15, 17, 25, 88
 Catabolism, 8, 88
 Cataract, 19, 88
 Cations, 88, 97
 Cause of Death, 88, 91
 Cell Cycle, 44, 45, 88
 Cell Death, 45, 85, 88, 100
 Cell Division, 37, 44, 45, 57, 58, 86, 88, 98, 99
 Cell Respiration, 88, 99
 Centromere, 37, 40, 88
 Cephalosporins, 9, 88
 Cerebrovascular, 87, 88
 Cerumen, 21, 88
 Chin, 88, 99
 Cholesterol, 36, 86, 88, 91
 Chromatin, 86, 88, 98
 Chromosomal, 45, 47, 48, 58, 59, 60, 62, 85, 88, 95, 99
 Chromosome, 3, 6, 7, 11, 18, 23, 29, 37, 38, 39, 40, 41, 44, 45, 47, 48, 54, 55, 58, 59, 64, 67, 85, 88, 89, 95, 97, 98, 99, 107
 Chromosome Fragility, 89, 107

- Chronic, 86, 89, 96, 97
 Cirrhosis, 26, 29, 89, 95
 Clavulanic Acid, 9, 89
 Clinical Medicine, 71, 89, 102
 Clinical trial, 8, 68, 69, 72, 75, 89, 101, 103
 Cloning, 21, 87, 89
 Codon, 42, 89
 Coenzyme, 86, 89
 Colon, 51, 89
 Colonoscopy, 53, 89
 Complement, 89, 90, 97
 Complementary and alternative
 medicine, 31, 32, 90
 Complementary medicine, 31, 90
 Computational Biology, 75, 90
 Concentric, 90, 100
 Conception, 44, 90, 93, 105
 Confusion, 51, 90, 92, 108
 Connective Tissue, 3, 4, 6, 86, 87, 90, 93,
 100, 106
 Connective Tissue Cells, 90
 Consciousness, 90, 91
 Constriction, 37, 40, 85, 90
 Consultation, 59, 60, 63, 64, 90
 Contraindications, ii, 90
 Coronary, 20, 28, 87, 90, 91
 Coronary Artery Bypass, 28, 91
 Coronary heart disease, 87, 91
 Cytochrome, 91, 101
 Cytoplasm, 34, 35, 36, 42, 86, 91, 94, 98,
 100, 104
 Cytosine, 35, 91, 103
- D**
- De novo, 45, 91
 Death Certificates, 53, 91
 Degenerative, 91, 95
 Deletion, 47, 86, 91
 Dementia, 48, 91
 Deoxyribonucleic, 35, 91, 104
 Deoxyribonucleic acid, 35, 91, 104
 Deoxyribonucleotides, 91
 Diabetes Mellitus, 91, 95
 Diagnostic procedure, 91
 Digestion, 86, 91, 98, 106, 108
 Diploid, 85, 91, 99, 107
 Direct, iii, 63, 64, 65, 89, 92, 104, 106
 Discrete, 92, 106
 Discrimination, 66, 71, 92
 Disorientation, 90, 92
 Distal, 91, 92
 Dopamine, 92, 100, 102
 Duct, 92, 106
- E**
- Effector, 89, 92
 Effector cell, 92
 Electrolytes, 86, 92, 97
 Electrons, 85, 86, 92, 97, 101, 103
 Embryo, 44, 45, 46, 54, 85, 87, 92
 Environmental Health, 74, 75, 92
 Enzyme, 4, 6, 8, 21, 36, 42, 82, 89, 92, 94,
 101, 103, 109
 Epidermis, 92, 96
 Epinephrine, 92, 100, 107
 Erythrocytes, 85, 87, 92
 Ethnic Groups, 59, 62, 93
 Eukaryotic Cells, 93, 101
 Excrete, 85, 93, 97, 104
 Extracellular, 90, 93
 Extracellular Matrix, 90, 93
 Eye Color, 46, 93
 Eye Infections, 84, 93
- F**
- Family Planning, 75, 93
 Fat, 87, 91, 93, 105
 Fathers, 54, 93
 Fetus, 62, 63, 65, 69, 93, 102, 106, 108
 Fibrosis, 46, 49, 53, 54, 93, 105
 Flatus, 93
 Forearm, 87, 93
 Frameshift, 47, 93
 Frameshift Mutation, 47, 93
 Fungus, 88, 93
- G**
- Ganglia, 93, 96, 100
 Gas, 29, 93, 96
 Gastrin, 93, 96
 Gene Expression, 42, 43, 94
 Gene Products, rev, 94
 Gene Therapy, 8, 67, 68, 69, 70, 84, 94
 Genes, env, 53, 94
 Genetic testing, 56, 60, 61, 62, 63, 64, 65,
 66, 71, 94
 Genomics, 11, 18, 29, 72, 94
 Genotype, 7, 14, 94, 102
 Germ Cells, 45, 69, 94, 98, 105
 Germline mutation, 45, 94, 95
 Gland, 94, 101, 106, 107
 Glomerular, 94, 97, 104
 Glucose, 86, 87, 91, 94, 95
 Governing Board, 94, 102
 Grafting, 91, 94, 96
 Granule, 94, 104
 Granulocytes, 94, 98, 109
 Guanine, 35, 95, 103

H

Hair Color, 46, 95
 Heart attack, 87, 95
 Hemochromatosis, 62, 95
 Hemodialysis, 95, 97
 Hemoglobin, 36, 85, 93, 95
 Hemoglobinopathies, 94, 95
 Hemophilia, 54, 95
 Hemorrhage, 95, 106
 Hepatic, 8, 11, 95
 Hepatitis, 26, 95
 Hepatocytes, 95
 Hereditary, 34, 35, 45, 54, 60, 94, 95, 101, 104
 Hereditary mutation, 45, 94, 95
 Heredity, 12, 37, 93, 94, 95
 Heterogeneity, 19, 95
 Heterozygote, 12, 95
 Histones, 37, 88, 95
 Homologous, 11, 84, 94, 95
 Hormone, 42, 92, 93, 96, 105, 106
 Hydrogen, 84, 86, 96, 99, 101
 Hydroxylation, 9, 96
 Hypercalcemia, 25, 96
 Hyperpigmentation, 24, 96
 Hypertension, 87, 96
 Hypokinesia, 96, 101

I

Immune response, 85, 96, 109
 Immune system, 92, 96, 98, 108, 109
 Impairment, 86, 93, 96, 99
 Implantation, 90, 96
 In vitro, 94, 96, 104
 In vivo, 8, 94, 96
 Individuality, 29, 96
 Infancy, 22, 72, 96
 Infection, 86, 93, 96, 100, 101, 109
 Inflammation, 68, 93, 95, 96, 102
 Informed Consent, 63, 66, 71, 96
 Initiation, 97, 107
 Inner ear, 97, 108
 Insight, 9, 97
 Interstitial, 97, 104
 Intravenous, 20, 97
 Ion Channels, 97
 Ions, 9, 86, 92, 96, 97
 Iris, 93, 97
 Irradiation, 97, 104

K

Karyotype, 39, 97
 Kidney Failure, 48, 97
 Kidney Failure, Acute, 97

Kidney Failure, Chronic, 97
 Kidney stone, 97, 101, 104, 108

L

Lens, 88, 98
 Lesion, 91, 98
 Leucocyte, 84, 98
 Leukemia, 94, 98
 Ligaments, 90, 98
 Linkages, 95, 98
 Liver, 6, 8, 26, 43, 86, 89, 95, 98
 Liver cancer, 8, 98
 Liver Transplantation, 26, 98
 Lumbar, 11, 14, 86, 98
 Lymphocytes, 85, 98, 109
 Lysine, 95, 98

M

Macrophage, 45, 98
 Malignant, 85, 98
 Mammary, 91, 98
 Mammography, 53, 98
 Medical Records, 53, 66, 98
 MEDLINE, 75, 98
 Meiosis, 44, 98
 Melanin, 97, 98, 102, 107
 Melanocytes, 96, 98
 Melanoma, 98, 108
 Membrane, 35, 85, 90, 93, 97, 99, 100, 101, 106
 Memory, 91, 99
 Mental, iv, 8, 58, 60, 62, 74, 76, 88, 90, 91, 92, 96, 99, 103, 105, 108
 Mental Retardation, 58, 60, 62, 99
 Microbe, 99, 107
 Microbiology, 86, 99
 Microorganism, 99, 109
 Miscarriage, 65, 99
 Mitochondria, 35, 36, 48, 54, 55, 99, 101
 Mitosis, 44, 86, 99
 Molecular, 5, 6, 7, 8, 14, 22, 24, 28, 29, 39, 41, 43, 75, 77, 87, 90, 99
 Molecule, 6, 35, 36, 37, 42, 85, 86, 89, 90, 92, 95, 99, 101, 102, 104, 108
 Monitor, 99, 100
 Monosomy, 48, 85, 99
 Morphological, 92, 93, 98, 99
 Morphology, 88, 99
 Mosaicism, 45, 99
 Mutagens, 93, 99
 Myotonic Dystrophy, 57, 99, 107

N

NCI, 1, 73, 89, 99, 101
 Necrosis, 86, 100

- Neonatal, 8, 25, 100
 Nephropathy, 15, 100
 Nervous System, 57, 93, 100
 Neuroblastoma, 13, 27, 100
 Neurons, 93, 100, 106
 Neuropathy, 54, 100
 Neurotransmitter, 84, 92, 97, 100, 105
 Nuclear, 35, 92, 93, 94, 100
 Nuclear Envelope, 35, 100
 Nuclear Pore, 100
 Nuclei, 92, 94, 95, 99, 100
 Nucleic acid, 86, 91, 99, 100, 103, 104
 Nucleus, 35, 36, 37, 42, 48, 67, 70, 86, 88,
 91, 93, 98, 100, 106
 Nurse Practitioners, 63, 100
- O**
- Ochronosis, 3, 5, 10, 11, 12, 14, 15, 19, 21,
 22, 25, 26, 100
 Oliguria, 97, 100
 Opacity, 88, 101
 Organelles, 34, 35, 91, 98, 101, 102
 Ovaries, 62, 101
 Oxalate, 23, 101
 Oxidation, 9, 84, 85, 91, 101
 Oxidative Phosphorylation, 36, 101
- P**
- Pancreas, 95, 101
 Parkinsonism, 22, 101
 Paternity, 62, 101
 Pathologic, 86, 90, 101
 Pathologic Processes, 86, 101
 Pathophysiology, 8, 101
 PDQ, 73, 101
 Pelvis, 84, 97, 98, 101, 108
 Penicillin, 101, 108
 Peptide, 101, 103
 Peroxidase, 26, 101
 Peroxide, 101
 Pharmacologic, 102, 107
 Phenotype, 7, 8, 14, 102
 Phenylalanine, 4, 6, 42, 102, 107
 Phosphorus, 87, 102
 Phosphorylation, 36, 102
 Physical Examination, 60, 102
 Physiologic, 87, 96, 102, 104, 107
 Pigment, 3, 21, 98, 100, 102
 Pigmentation, 10, 21, 96, 102
 Plasma, 35, 95, 97, 102
 Plastids, 101, 102
 Pneumonia, 90, 102
 Point Mutation, 10, 102
 Polymorphic, 13, 102
 Polymorphism, 24, 64, 102
 Posterior, 86, 97, 101, 102
 Practice Guidelines, 76, 102
 Precursor, 92, 102, 107, 108
 Prenatal, 62, 65, 92, 102
 Prevalence, 18, 50, 102
 Progressive, 48, 89, 91, 97, 99, 100, 103, 104
 Prolapse, 14, 103
 Prone, 48, 57, 103
 Proteolytic, 84, 89, 103
 Protocol, 68, 103
 Psychic, 99, 103
 Public Policy, 75, 103
 Publishing, 79, 103
 Pulmonary, 87, 97, 103, 108
 Pulmonary Artery, 87, 103, 108
 Pulmonary Edema, 97, 103
 Purines, 86, 103
 Pyrimidines, 86, 103
- R**
- Race, 97, 103
 Radiation, 84, 97, 103, 104, 108, 109
 Radiation therapy, 84, 97, 104
 Radioactive, 96, 97, 100, 104
 Receptor, 51, 85, 92, 104
 Recombinant, 68, 104, 108
 Recombination, 94, 104
 Rectum, 85, 89, 93, 104
 Refer, 1, 40, 44, 46, 51, 70, 87, 89, 104
 Regurgitation, 20, 104
 Renal failure, 12, 104
 Renal tubular, 8, 104
 Repopulation, 8, 104
 Reproductive cells, 48, 58, 59, 94, 95, 104
 Retinoblastoma, 50, 104
 Retroviral vector, 94, 104
 Ribonucleic acid, 42, 104
 Ribose, 84, 104
 Ribosome, 42, 104, 107
 Rigidity, 101, 104
 Ristocetin, 104, 108
- S**
- Saphenous, 91, 104
 Saphenous Vein, 91, 104
 Scatter, 105, 108
 Schizophrenia, 55, 105
 Sclerosis, 51, 105
 Screening, 13, 27, 53, 62, 63, 65, 89, 101,
 105, 108
 Second Messenger Systems, 105
 Sediment, 105, 108
 Sequencing, 70, 105

Side effect, 69, 72, 84, 105, 106, 107
 Signs and Symptoms, 4, 56, 57, 62, 105
 Skin Pigmentation, 5, 22, 105
 Small intestine, 96, 105
 Social Work, 59, 105
 Soft tissue, 87, 105
 Solid tumor, 87, 105
 Soma, 105
 Somatic, 45, 48, 59, 98, 99, 105
 Somatic cells, 45, 48, 59, 98, 99, 105
 Somatic mutations, 48, 105
 Specialist, 63, 80, 105
 Species, 72, 92, 97, 98, 99, 103, 105, 106, 107, 109
 Spectroscopic, 9, 106
 Sperm, 44, 45, 48, 57, 58, 59, 62, 69, 88, 94, 95, 104, 105, 106
 Spinal cord, 88, 100, 106
 Sporadic, 104, 106
 Stenosis, 10, 11, 20, 106
 Stillbirth, 60, 106
 Stimulant, 106, 108
 Stomach, 93, 96, 105, 106
 Stool, 89, 106
 Strand, 35, 106
 Stricture, 106
 Stroke, 53, 74, 87, 106
 Subspecies, 105, 106
 Supportive care, 101, 106
 Sweat, 88, 106
 Sweat Glands, 88, 106
 Synapses, 106
 Systemic, 85, 87, 92, 96, 97, 104, 106

T
 Tendon, 28, 106
 Terminator, 89, 106
 Thoracic, 86, 106, 109
 Thrombosis, 103, 106
 Thyroid, 62, 106, 107
 Thyroid Gland, 62, 107
 Thyroid Hormones, 107
 Thyroxine, 102, 107
 Tone, 101, 107
 Toxic, iv, 34, 100, 107, 108
 Toxicity, 68, 104, 107
 Toxicology, 75, 107
 Trachea, 87, 106, 107
 Transcription Factors, 43, 107

Transfection, 87, 94, 107
 Translation, 42, 43, 94, 107
 Transplantation, 22, 97, 107
 Tremor, 101, 107
 Trinucleotide Repeat Expansion, 57, 107
 Trinucleotide Repeats, 107
 Trisomy, 48, 85, 107
 Tyrosine, 4, 6, 8, 29, 92, 107

U
 Ultraviolet radiation, 45, 107
 Uremia, 97, 104, 108
 Urethra, 108
 Uric, 26, 103, 108
 Urinalysis, 21, 82, 108
 Urinary, 11, 26, 86, 100, 108
 Urinary tract, 86, 108
 Urine, 3, 4, 6, 12, 21, 23, 24, 85, 86, 87, 97, 101, 108
 Urography, 20, 108
 Urolithiasis, 23, 108
 Uterus, 62, 101, 108

V
 Vaccine, 103, 108
 Vacuoles, 101, 108
 Valine, 6, 108
 Vancomycin, 9, 108
 Vascular, 96, 107, 108
 Vector, 67, 68, 108
 Vein, 97, 100, 104, 108
 Ventricle, 85, 103, 108
 Venules, 87, 109
 Veterinary Medicine, 75, 109
 Viral, 67, 94, 109
 Virulence, 107, 109
 Virus, 67, 104, 109
 Viscera, 105, 109
 Vitro, 62, 109
 Vivo, 109

W
 White blood cell, 45, 85, 98, 109
 Windpipe, 106, 109
 Womb, 108, 109

X
 X-ray, 9, 97, 100, 104, 109

Y
 Yeasts, 93, 102, 109

Z
 Zygote, 90, 99, 109