



THE WORLD BANK

WORLD BANK OPERATIONS EVALUATION DEPARTMENT



Maintaining Momentum to 2015?

An Impact Evaluation of Interventions to Improve Maternal and Child Health and Nutrition in Bangladesh





Maintaining Momentum to 2015?

An Impact Evaluation
of Interventions to Improve
Maternal and Child Health
and Nutrition in Bangladesh



© 2005 The International Bank for Reconstruction and Development / The World Bank
1818 H Street, NW
Washington, DC 20433
Telephone: 202-473-1000
Internet: www.worldbank.org
E-mail: feedback@worldbank.org

All rights reserved
Manufactured in the United States of America
First edition August 2005

The findings, interpretations, and conclusions expressed here are those of the author(s) and do not necessarily reflect the views of the Board of Executive Directors of the World Bank or the governments they represent.

The World Bank cannot guarantee the accuracy of the data included in this work. The boundaries, colors, denominations, and other information shown on any map in the work do not imply on the part of the World Bank any judgment of the legal status of any territory or the endorsement or acceptance of such boundaries.

Rights and Permissions

The material in this publication is copyrighted. Copying and/or transmitting portions or all of this work without permission may be a violation of applicable law. The World Bank encourages dissemination of its work and will normally grant permission promptly.

For permission to photocopy or reprint any part of this work, please send a request with complete information to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, USA, telephone 978-750-8400, fax 978-750-4470, www.copyright.com.

All other queries on rights and licenses, including subsidiary rights, should be addressed to the Office of the Publisher, World Bank, 1818 H Street NW, Washington, DC 20433, USA, fax 202-522-2422, e-mail pubrights@worldbank.org.

Cover photo courtesy of the World Bank Photo Library.

ISBN 0-8213-6376-X
e-ISBN 0-8213-6377-8

Library of Congress Cataloging-in-Publication Data is in process

World Bank InfoShop
E-mail: pic@worldbank.org
Telephone: 202-458-5454
Facsimile: 202-522-1500

Operations Evaluation Department
Knowledge Programs and Evaluation Capacity
Development Group (OEDKE)
Room # 19-352
MSN # 19-903
(202) 458-0382
Facsimile: (202) 522-3125



Printed on Recycled Paper

Contents

vii	Acknowledgments
ix	Executive Summary
xiii	Résumé analytique
xix	Resumen
xxiii	Acronyms and Abbreviations
1	1. Maternal and Child Health in Bangladesh: A Record of Success
3	Scope of the Study
5	Evaluation Approach
5	Overview of the Report
7	2. Health, Family Planning, and Nutrition Services in Bangladesh: An Overview
7	Family Planning Programs
10	Health Services
12	Nutrition
13	3. Trends in Under-Five Mortality, Nutrition, and Fertility
13	Patterns of Mortality Decline
13	Anthropometric Outcomes
14	What Has Been Happening to Fertility?
17	4. Impact of Specific Interventions on Child Health and Fertility
17	Income Growth Accounts for Some, But Not All, Improvement in Outcomes
19	Under-Five Mortality
26	Fertility Reduction
28	Nutrition

29	5. A Closer Look at Nutrition: The Bangladesh Integrated Nutrition Project
30	Overview of the Project
30	Project Coverage and Targeting
33	Acquiring Knowledge
34	Turning Knowledge into Practice: The Knowledge-Practice Gap
36	The Nutritional Impact of BINP Interventions
38	Testing the Theory in Practice—How Well Did the Causal Chain Operate?
41	6. Lessons Learned
45	Annexes
45	A. Trends in Maternal and Child Health Outcomes
49	B. Cross-Country Analysis of Child Health and Nutrition Outcomes
65	C. Neonatal, Postnatal, and Child Mortality in the 1990s
109	D. Child Malnutrition during the 1990s
125	E. Women's Agency, Household Structure, and Health Outcomes
139	F. Fertility
155	G. Analysis of BINP's Community-Based Nutrition Component
187	H. DFID and World Bank Programs in Bangladesh
193	I. Agricultural Production, Natural Disasters, Seasonality, and Nutritional Outcomes
199	J. Approach Paper
207	Endnotes
219	References
	Boxes
2	1.1 Measures of Welfare Outcomes
22	4.1 Which Children Get Immunized?
23	4.2 Polio Eradication in Bangladesh
36	5.1 Qualitative Perspectives of the Knowledge-Practice Gap: The PPS-BD Study
	Figures
3	1.1 Both Under-Five Mortality and Fertility Have Fallen Rapidly
11	2.1 Immunization Coverage of Children Aged 12–23 Months
14	3.1 Nutritional Status Improved in the 1990s
14	3.2 Data from Different Sources Present a Consistent Picture
16	3.3 Fertility Decline Continued in the 1990s according to a Range of Indirect Measures
16	3.4 Knowledge of Modern Contraceptives Is Universal and Use Continues to Rise
18	4.1 Bangladesh's Improvement in Social Outcomes Is Greater than Can Be Explained by Economic Growth Alone
25	4.2 Secondary Enrollments Have Risen Rapidly in the 1990s: Educational Attainment of Women 17 to 24 Years Old
32	5.1 Various Factors Affect Women's Participation, but Restrictions on Women's Mobility in More Conservative Areas Are the Most Important

34	5.2 Women Living in Project Areas Are More Likely to Have Nutritional Information, Especially if They Participate in Project Activities
35	5.3 The Knowledge-Practice Gap in Project Areas: More Women Say They Know Good Behavior than Actually Practice It
38	5.4 Rice Production and Daily Energy Supply Grew Rapidly in the Late 1990s

Tables

8	2.1 World Bank Credits for Health, Population, and Nutrition
15	3.1 Fertility Decline Has Always Been Erratic Based on Direct Estimates, but Continued into the 1990s Using Indirect Ones
19	4.1 Growth in GNP per Capita Accounts for at Most One-Third of the Reduction in Mortality . . . and Less than a Fifth of Lower Fertility
20	4.2 Significant Determinants of Infant and Child Mortality
23	4.3 Neonatal Mortality Has Fallen More Rapidly for Births Attended by Trained TBAs
35	5.1 Many Factors Prevent Women from Putting Nutritional Advice into Practice, Though the Project Partially Overcomes Some of These
37	5.2 Cost of Nutrition Improvements and Mortality Reduction
39	5.3 Links in the Causal Chain
42	6.1 Cost-Effectiveness of Interventions in Reducing Under-Fiver Mortality



Acknowledgments

This study is the second in a series of three impact studies being conducted by the Operations Evaluation Department (OED) under a Department for International Development (DFID)–OED partnership agreement. Financial support was also provided through Danish Consultant Trust Funds. In these studies, OED is exploring different ways of carrying out ex post impact analysis in cases that did not have an evaluation framework in place at the start of the intervention. For the first study of Ghana basic education (OED 2004b), a survey was commissioned to follow on from a nationally representative household and school survey conducted 15 years earlier. This second study is based on existing data sets: the Bangladesh Demographic and Health Surveys, the Household Income and Expenditure Survey, two data sets related to the Bangladesh Integrated Nutrition Project—the evaluation data set and that collected by Save the Children—and data from Helen Keller International’s (HKI) Nutritional Surveillance Project. Thanks are due to Meera Shekar of the World Bank’s Nutrition Hub, Dora Panagides of HKI, and Anna Taylor and Dr. Arabella Duffield of Save the Children for facilitating access to the latter three data sets.

This report was prepared by Howard White, with assistance from Nina Blöndal, Edoardo Mas-

set, and Hugh Waddington, and inputs provided by Professor Kabir and Dr. Sayed Haider of Research Evaluation Associates for Development (READ) and Dr. Sharifa Begum of the Bangladesh Institute of Development Studies (BIDS). In addition, a background paper was commissioned from the Participatory Practitioner’s Society–Bangladesh (PPS-BD). Preparation of the report was assisted by Alain Barbu, Martha Ainsworth, and Denise Vaillancourt. Professor Nicholas Mascie-Taylor (University of Cambridge) and Dr. Arabella Duffield (Save the Children U.K.) acted as external reviewers. Comments were also received from Harold Alderman, Farial Mahmud, Md. Abdul Maleque (of the Implementation, Monitoring and Evaluation Department of the Ministry of Planning in Bangladesh), Bina Valaydon, the Planning Commission of the Government of Bangladesh, and Bernabé Sánchez, Nick York, and Shona Wynd of the DFID Evaluation Department.

Thanks are due to all those in the government of Bangladesh, nongovernmental organizations (NGOs), and donor agencies who provided their time and support to the undertaking of this study. In the World Bank office in Dhaka, Rafael Cortez, Farzana Ishrat, and Shirin Jahangeer provided valuable inputs, and excellent logistical assistance was provided by the

support staff of the Health Project Support Office, including arranging a presentation of preliminary findings. From DFID, Dinesh Nair and Neil Squires provided useful guidance. Thanks are due to Kaosar Afsana of Bangladesh Rural Advancement Committee (BRAC) and Mohammed

Azizul Hoque of Thengamara Mohila Sabuj Sangha for their assistance in trips to the field.

Administrative support was provided by Soon Woon-Pak. The report was edited by Bill Hurlbut, Michele Spring, and Caroline McEuen.

Acting Director-General, Operations Evaluation: *Ajay Chhibber*
Acting Director, Operations Evaluation Department: *Kyle Peters*
Manager, Sector, Thematic, and Global Evaluation Unit: *Alain Barbu*
Task Manager: *Howard Nial White*



Executive Summary

Improving maternal and child health and nutrition is central to development goals. The importance of these objectives is reflected by their inclusion in poverty-reduction targets such as the Millennium Development Goals (MDGs) and Bangladesh's Interim Poverty Reduction Strategy Paper, supported by major development partners, including the World Bank and the U.K. Department for International Development (DFID).

This report addresses the issue of what publicly supported programs and external assistance from the Bank and other agencies can do to accelerate attainment of such targets as reducing infant mortality by two-thirds. The evidence presented here relates to Bangladesh, a country that has made spectacular progress, but needs to maintain momentum in order to achieve its own poverty-reduction goals.

The report addresses the following issues: (1) What has happened to child health and nutrition outcomes and fertility in Bangladesh since 1990? Are the poor sharing in the progress being made? (2) What have been the main determinants of maternal and child health (MCH) outcomes in Bangladesh over this period? (3) Given these determinants, what can be said about the impact of publicly and externally supported programs—notably those of the World Bank and DFID—to improve health and nutrition? (4) To the extent that interventions have brought about positive impacts, have they done so in a cost-effective manner?

Trends in Under-Five Mortality, Fertility, and Nutrition

Despite an inauspicious start coming out of war and famine, Bangladesh has achieved spectacular rates of progress in the past two decades, most notably with respect to fertility decline. Contrary to common perceptions, fertility continued to decline during the 1990s. Under-five mortality has also been reduced at a substantial rate—Bangladesh is one of the few countries to achieve a sufficient rate of reduction to achieve the MDG of a two-thirds decline by 2015. The exception to these successes has been nutrition. Physical measures of nutritional status only began to show some improvement in the 1990s, and malnutrition remains at high levels.

Improvements in these outcomes have been spread across all Bangladeshis. Although children of the poor are more likely to suffer premature death than their better-off counterparts, this gap is narrowing, with mortality rates falling faster among the poor than the non-poor. Con-

traceptive use and low fertility are also common among the poor.

Sources of Under-Five Mortality Decline

Analysis of the determinants of mortality using both cross-country and Demographic and Health Survey (DHS) data shows that a variety of factors have influenced the decline in under-five deaths. Improved economic well-being is the most important reason for lower child mortality, but plays less of a role for infants. Social sector interventions—both health and education—are also found to matter, with expanded immunization coverage and greater female enrollment in primary and secondary education both playing a substantial part in mortality reduction. The results also show a pronounced sex bias in mortality against girls, especially in the Sylhet and Chittagong divisions. Analysis of selected interventions reveals the following:

- Immunization coverage was at less than 2 percent in the early 1980s, but grew in the latter part of the decade (largely with the support of the United Nations Children's Fund [UNICEF], but later also with the support of other donors, including the World Bank), so that by 1990, close to half of all children were fully vaccinated in their first 12 months. Immunization has averted more than two million child deaths in the last two decades, at a cost of \$100–\$300 per life saved.
- The World Bank financed the training of approximately 14,000 traditional birth attendants (TBAs) through the late 1990s, when training of TBAs was abandoned following a shift in international opinion toward a policy of all births being attended by skilled birth attendants. However, the evidence presented in this report shows that training TBAs saved infant lives, at a cost of \$220–\$800 per life saved.
- Female secondary schooling expanded rapidly in the 1990s, especially in rural areas, partly as a result of the stipend paid to all female students in grades 6–10 in rural areas. This stipend was supported by Norwegian aid, the Asian Development Bank, the World Bank, and the Bangladesh government. Among the benefits of the increase in female secondary schooling is lower mortality, at a cost of \$1,080–\$5,400 per life saved.
- Rural electrification, supported through three World Bank programs in the 1980s and 1990s, reduces mortality through income effects, improved health services, easier water sterilization, and improved access to health information, especially from TV. Taking these various channels into account means that children in households receiving electrification have an under-five mortality rate of 25 per 1,000 lower than that of children in non-electrified households. Based on historic costs, this amounts to \$20,000 per life saved, and \$40,000 based on current connection costs.

Nutrition

In order to address the poor state of nutrition, the government implemented, with World Bank assistance, the pilot Bangladesh Integrated Nutrition Project (BINP). The core of BINP is the Community-Based Nutrition Component (CBNC), which promotes nutrition counseling to bring about behavior change, complemented by supplementary feeding for pregnant women and young children.

Analysis of the causal chain from BINP inputs to child anthropometric outcomes shows the following:

- There is a weak link in the chain, as behavioral change communication has been excessively focused on mothers, who are often not the main decisionmakers for all nutrition-related practices.
- Program coverage is generally high in project areas, but notably lower in more conservative thanas (subdistricts), especially among women who live with their mothers-in-law.
- There are some deficiencies in targeting: (a) too strict a criterion was applied in admitting malnourished children to supplementary feeding, while admitting children who were growth-faltering but probably well-nourished, and (b) feeding of pregnant women excluded many who were eligible, and included those who were not.
- A large proportion of mothers of children receiving supplementary feeding claim not to have received nutritional counseling.

- There is a substantial knowledge-practice gap, whereby women do not turn the advice they receive into practice (economic resource and time constraints are a major reason for this).
- The impact on pregnancy weight gain is too small to have a substantial impact on birth weight. This situation is common in other programs; the mother's pre-pregnancy nutritional status is a more important factor in low birth weight than pregnancy weight gain and might therefore have been a better focus for the project.

The list above may be read as problems to be fixed in the project. To some extent they have been addressed under the expanded National Nutrition Project: the targeting criteria for children's supplementary feeding have been revised and another attempt made to reach men with nutritional counseling. But the program has not been a very cost effective means of improving nutritional status—which has improved generally with the acceleration in food availability associated with the yield-driven increase in rice production since the late 1990s, and consequent reduction in the real price of rice. Simulations show that simply giving food to families with children would have had a larger nutritional impact. The cost per life saved from the hypothetical rice ration is just over \$2,000, half the cost of lives saved by BINP.

Fertility Reduction

The rate of fertility reduction in Bangladesh is shown to exceed that which may be expected from other socioeconomic developments, such as income growth and expanding female education. While socioeconomic developments, including the demographic transition, explain a part of Bangladesh's rapid fall in fertility, a large part is attributable to the country's family planning service, built up with substantial external support in the years following liberation in 1971. The continued decline of fertility in the 1990s, driven by rising contraceptive prevalence, demonstrates the continued effectiveness of this program.

The government's health, nutrition, and population (HNP) Strategic Investment Plan highlights the role of increasing the age at marriage as a means of reducing fertility, and several programs, including the counseling provided under BINP, promote getting married later. It is a condition of the female secondary school stipend program, supported by the Bank among others, that recipients remain unmarried. It is true that the age at marriage in Bangladesh is low, with half of all women marrying by age 14. It is also true that there is a well-established international pattern whereby increasing the age at marriage drives down fertility. But this pattern should not be expected to be observed in Bangladesh for two reasons: (1) raising the age at marriage of girls aged 13 or less has no effect on the age at which they have their first child (as the age at marriage has risen, the gap from marriage to first birth has fallen); and (2) if a woman plans to have only three to four children, as the majority of Bangladeshi women do, then this can be accomplished whether childbearing begins at 15 or 20. The direct effect of expanding secondary education will be muted, as Bangladesh has already attained fertility levels comparable to those in countries with higher education levels. Hence, raising the age at marriage, while desirable for both maternal and child health (children born to young mothers have a greater chance of premature death), will have little impact on the number of children borne by each woman during her reproductive years—though there would be a temporary tempo effect on the total fertility rate and a second-order effect as the mortality-reducing effect of later births will reduce the desired number of births. Instead, high-fertility households should be targeted, partly by an attempt to restore the use of permanent contraceptive measures to their previous levels. Efforts should also be made to tackle son preference, which creates a barrier to fertility decline. And continued success in reducing mortality will also help reduce fertility.

Lessons Learned

The following general lessons follow from the analysis in this report:

- Externally supported interventions have had a notable impact on MCH-related outcomes in Bangladesh. Immunization has proved particularly cost effective, and has saved the lives of up to two million children under the age of five.
- World Bank support to sectors outside of health has contributed to better child health outcomes.
- Small amounts of money save lives . . . although the amount varies significantly by intervention.
- Although interventions from many sectors affect maternal and child health outcomes, this fact need not imply that multisectoral interventions are always needed.
- World Bank support for training traditional birth attendants has reduced neonatal mortality . . . but this program has now been abandoned following the international trend toward support for skilled birth attendants.
- Programs should be based on local evidence, rather than on general conventional wisdom.
- Gender issues are central to health strategies in Bangladesh. More attention is needed to redressing gender biases to maintain momentum in mortality decline and fertility reduction. But traditional attitudes are not the absolute constraint on service provision, as is sometimes suggested.
- The Bank's BINP has improved nutritional status, but not by as much as planned. Serious attention needs to be given to ways of improving both the efficacy and efficiency of the program—or, if this is not possible, then to consideration of alternatives to scaling up.
- Rigorous impact evaluation can show which government programs and external support are contributing the most to meeting poverty-reduction goals.
- National surveys can be used for evaluation purposes, but some adaptations would make them more powerful, notably a more detailed community questionnaire.



Résumé analytique

L'amélioration de la nutrition et de la santé maternelle et infantile tient une place centrale dans les objectifs de développement. L'importance de ces objectifs ressort du fait qu'ils sont inclus dans les objectifs de réduction de la pauvreté, tels que les objectifs de développement pour le Millénaire (ODM) ou ceux établis dans le document intérimaire de stratégie pour la réduction de la pauvreté du Bangladesh, auxquels les principaux partenaires de développement, y compris la Banque mondiale et le DFID, apportent leur concours.

Ce rapport examine ce que les programmes menés sur financement public et l'aide extérieure fournie par la Banque et les autres organisations peuvent faire pour accélérer la réalisation des objectifs tels que la réduction de deux tiers de la mortalité infantile. Les éléments d'appréciation qu'il présente concernent le Bangladesh, un pays qui a accompli des progrès spectaculaires mais qui doit maintenir l'élan acquis afin d'atteindre ses propres objectifs de réduction de la pauvreté.

Les questions ici traitées sont les suivantes: 1) Comment ont évolué la situation sanitaire et nutritionnelle des enfants et la fécondité au Bangladesh depuis 1990? Les pauvres prennent-ils part aux progrès réalisés? 2) Quels ont été les principaux déterminants de la situation du Bangladesh en matière de santé maternelle et infantile (SMI) durant cette période? 3) Compte tenu de ces facteurs déterminants, que peut-on dire

de l'incidence des programmes exécutés sur financements publics et extérieurs—tout particulièrement ceux de la Banque mondiale et du DFID—pour ce qui est d'améliorer la santé et la nutrition? 4) Dans la mesure où les interventions ont eu des incidences positives, cela s'est-il fait dans de bonnes conditions de coût-efficacité?

Tendances de la mortalité des moins de 5 ans, de la fécondité et de la nutrition

En dépit d'une situation de départ peu prometteuse, puisqu'il sortait d'une période de guerre et de famine, le Bangladesh a réalisé ces 20 dernières années des progrès spectaculaires, surtout pour ce qui est de la baisse de la fécondité. Contrairement à ce que l'on croit en général, la fécondité a continué de baisser dans les années 90. La mortalité des moins de 5 ans a elle aussi diminué sensiblement, le Bangladesh étant l'un des rares pays à connaître à un taux de diminu-

tion suffisant pour être en mesure de parvenir à la réduction de deux tiers fixée par les ODM à l'horizon 2015. Le domaine qui a fait exception à ces résultats positifs est la nutrition. Les indicateurs matériels de l'état nutritionnel ont commencé à donner des signes de progrès seulement dans les années 90, et les niveaux de malnutrition restent élevés.

Pour ce qui est des progrès réalisés, ils ont concerné l'ensemble de la population. S'il est vrai que les risques de décès prématuré sont plus marqués pour les enfants des milieux défavorisés, le décalage qui existe à cet égard diminue progressivement, les taux mortalité baissant à un rythme plus rapide chez les pauvres que chez les non-pauvres. L'emploi de contraceptifs et un faible taux de fécondité sont également fréquents parmi les pauvres.

Facteurs à l'origine de la baisse de la mortalité des moins de 5 ans

L'analyse des facteurs de mortalité basée sur les données couvrant différents pays ainsi que sur les données de l'enquête démographique et sanitaire (EDS) révèle qu'une diversité de facteurs sous-tend la réduction observée du nombre de décès. L'amélioration du bien-être économique est la principale raison expliquant la baisse de la mortalité juvénile, mais elle joue moins dans le cas de la mortalité infantile. On constate par ailleurs que les interventions dans les secteurs sociaux—santé et éducation—ont de l'importance, l'augmentation de la couverture vaccinale et l'accroissement des taux de scolarisation féminine dans le primaire et le secondaire contribuant l'un et l'autre pour une bonne part à la baisse de la mortalité. Les données recueillies font par ailleurs état d'une tendance particulièrement défavorable de la mortalité à l'égard des filles, surtout dans les divisions de Sylhet et de Chittagong. L'analyse de certaines interventions fait ressortir les éléments suivants:

- La couverture vaccinale était de moins de 2 % au début des années 80, mais elle a augmenté dans la seconde partie de cette décennie (en grande partie avec l'appui de l'UNICEF mais,

par la suite, avec celui d'autres bailleurs de fonds également, y compris la Banque mondiale), si bien que près de la moitié des enfants en 1990 étaient totalement vaccinés en atteignant 12 mois. La vaccination a permis d'éviter plus de 2 millions de décès d'enfants au cours des 20 dernières années, pour un coût oscillant entre 100 et 300 dollars par vie sauvée.

- La Banque mondiale a financé la formation d'environ 14 000 accoucheuses traditionnelles jusqu'à la fin des années 90, après quoi ce type de formation a été abandonné à la suite d'un revirement de l'opinion internationale dans le sens d'une politique tendant à ce que toutes les naissances soient assistées par des professionnels qualifiés. Mais les éléments d'appréciation présentés dans ce rapport montrent que la formation des accoucheuses traditionnelles a permis de sauver des vies d'enfants, pour un coût de 220 à 800 dollars par décès évité.
- La scolarisation féminine dans le secondaire a progressé rapidement dans les années 90, et ce surtout en milieu rural, du fait en partie des allocations d'études accordées à toutes les élèves de la classe de 6e à la 3e avec l'appui de l'aide norvégienne, de la Banque asiatique de développement, de la Banque mondiale et du gouvernement. Parmi les avantages qui en découlent figure une réduction de la mortalité, pour un coût de 1 080 à 5 400 dollars par décès évité.
- L'électrification des zones rurales, qui a reçu l'appui de trois programmes de la Banque mondiale dans les années 80 et 90, réduit la mortalité grâce aux effets de revenu qui en découlent, à l'amélioration des services de santé, à la possibilité de stériliser plus facilement l'eau et à la plus grande accessibilité de l'information sanitaire, du fait principalement de la télévision. La prise en compte de ces divers facteurs aboutit à un taux de mortalité des moins de 5 ans inférieur de 25 pour 1 000 chez les enfants des ménages ayant accès à l'électricité, comparé à ceux qui n'y ont pas accès. Sur la base des coûts historiques, cela représente 20 000 dollars par vie sauvée; aux coûts de raccordement actuels, cela équivaut à 40 000 dollars.

Nutrition

Pour faire face au mauvais état nutritionnel de sa population, le pays a mis en œuvre avec l'aide de la Banque mondiale une opération pilote, le Projet intégré de nutrition du Bangladesh (BINP). Son élément central est la composante de nutrition communautaire, consistant à promouvoir les services de conseils nutritionnels dans le but de susciter une modification des comportements, un volet complété par la fourniture d'une alimentation d'appoint aux femmes enceintes et aux jeunes enfants.

Une analyse de la chaîne de causalité mettant en rapport, d'un côté, les apports du BINP et, de l'autre, les données anthropométriques qui se dégagent au niveau des enfants révèle les éléments suivants: 1) la chaîne présente un maillon faible en ce sens que la communication en matière de modification des comportements a trop privilégié les mères, qui ne sont pas, dans bien des cas, le principal décideur pour l'ensemble des pratiques adoptées en matière nutritionnelle; 2) le niveau de couverture du programme est généralement élevé dans les zones visées mais notablement moindre dans les thanas (sous-districts) moins en pointe, surtout parmi les femmes vivant avec leur belle-mère; 3) le ciblage présente quelques lacunes: a) un critère trop strict a été appliqué lorsqu'il s'est agi d'admettre les enfants souffrant de malnutrition à bénéficier d'une alimentation complémentaire, alors qu'on a admis des enfants qui avaient des problèmes de croissance mais qui étaient probablement bien nourris; b) le volet d'alimentation des femmes enceintes a exclu bon nombre de femmes qui y avaient droit, tout en admettant un certain pourcentage de femmes qui n'y avaient pas droit; 4) parmi les mères dont les enfants reçoivent une alimentation complémentaire, une forte proportion a indiqué ne pas avoir reçu de conseils nutritionnels; 5) il existe un net écart entre connaissances et pratiques, en ce sens que les femmes ne mettant pas en pratique les conseils qui leur sont dispensés, situation due pour beaucoup aux contraintes de temps et de ressources économiques; et 6) l'impact du programme sur la prise de poids en cours de grossesse est trop limité pour avoir une réelle incidence sur le poids à la naissance, état de fait

qui se retrouve fréquemment dans des programmes analogues entrepris dans d'autres pays; l'état nutritionnel de la mère avant la grossesse joue davantage dans l'insuffisance pondérale à la naissance que la prise de poids en cours de grossesse, et aurait donc pu être un meilleur axe de focalisation pour le projet.

La liste figurant dans le paragraphe qui précède peut se lire comme une liste de problèmes à régler dans le cadre du projet, comme ils l'ont été dans une certaine mesure au titre de l'opération élargie qu'est le Projet national de nutrition: les critères de ciblage pour l'alimentation complémentaire destinée aux enfants ont été révisés, et une autre tentative effectuée pour atteindre la population masculine dans le cadre des services de conseils nutritionnels. Mais le programme n'a pas été un moyen particulièrement rentable pour l'amélioration de l'état nutritionnel de la population, lequel s'est amélioré d'une manière générale grâce à l'augmentation rapide des ressources alimentaires disponibles qui a accompagné l'accroissement, sous l'effet des rendements, de la production rizicole depuis la fin des années 90, et la réduction du prix réel du riz qui en a résulté. Les simulations montrent que le simple fait de distribuer de la nourriture aux familles ayant des enfants aurait eu plus d'impact au plan nutritionnel. Le coût par vie sauvée au moyen de la ration hypothétique de riz s'élève à un peu plus de 2 000 dollars seulement, soit la moitié du coût des vies sauvées grâce au BINP.

Réduction de la fécondité

Le taux de réduction de la fécondité observé au Bangladesh est supérieur à celui auquel on peut s'attendre sous l'effet d'autres facteurs socioéconomiques, comme l'accroissement des revenus ou de la scolarisation féminine. Bien que les phénomènes socioéconomiques, et notamment la transition démographique, expliquent en partie la baisse rapide de la fécondité dans ce pays, cette baisse tient pour beaucoup au service de planning familial que celui-ci a établi, avec d'importants appuis extérieurs, dans les années ayant suivi l'indépendance en 1971. Le fait que le recul de la fécondité se soit poursuivi dans les années 90, sous l'effet de la prévalence accrue de la contra-

ception, démontre l'efficacité que continue d'avoir ce programme.

Le programme d'investissement stratégique en santé, nutrition et population établi par le gouvernement met en avant le recul de l'âge au mariage comme moyen de réduire la fécondité, et plusieurs programmes, dont les services de conseils fournis dans le cadre du BINP, encouragent les Bangladais à se marier plus tard. Une condition du programme d'allocations d'études destinées aux filles inscrites dans l'enseignement secondaire, programme appuyé notamment par la Banque, est que les bénéficiaires de ces allocations restent célibataires. Il est vrai que l'âge au mariage est peu élevé au Bangladesh: à 14 ans, la moitié des femmes sont mariées. Il est également vrai qu'il existe une tendance bien établie au plan international, selon laquelle le relèvement de l'âge au mariage s'accompagne d'une baisse de la fécondité. Mais on ne doit pas s'attendre à observer cette tendance au Bangladesh, et ce pour deux raisons: 1) le relèvement de l'âge au mariage pour les filles âgées de 13 ans ou moins n'a pas d'effet sur l'âge auquel elles ont leur premier enfant (ainsi, à mesure que l'âge au mariage a augmenté, le délai entre le mariage et la première naissance s'est réduit), et 2) si une femme compte avoir trois ou quatre enfants seulement, ce qui est le cas de la majorité des femmes au Bangladesh, elle pourra y parvenir qu'elle ait son premier enfant à 15 ans ou à 20 ans. L'effet direct de l'augmentation de la scolarisation secondaire sera estompé du fait que le Bangladesh a déjà atteint des taux de fécondité comparables à ceux observés dans les pays qui ont des taux de scolarisation plus élevés. Ainsi, le relèvement de l'âge au mariage, quoique souhaitable du point de vue de la santé maternelle et infantile (les enfants nés de mères jeunes sont plus exposés au risque de décès prématuré), n'aura guère d'impact sur le nombre d'enfants qu'aura chaque femme durant sa période de reproduction—étant entendu qu'il y aurait un effet de tempo provisoire sur l'indice synthétique de fécondité, et qu'il y aura un effet de second ordre à mesure que la baisse de la mortalité liée aux naissances plus tardives entraînera une réduction du nombre de naissances souhaité. Au lieu de cela, il conviendrait de faire un effort ciblé sur les ménages présentant de forts

taux de fécondité, en essayant notamment de ramener à ses niveaux antérieurs l'utilisation de mesures contraceptives permanentes. Des efforts devraient également être entrepris à l'égard de la préférence pour les fils, qui crée un obstacle au recul de la fécondité. Enfin, la poursuite des succès obtenus en matière de réduction de la mortalité contribuera également à faire baisser la fécondité.

Enseignements tirés

On peut tirer les enseignements généraux suivants de l'analyse contenue dans ce rapport:

- Les opérations entreprises avec un appui extérieur ont eu un impact notable sur la situation de la santé maternelle et infantile au Bangladesh. L'effort de vaccination s'est révélé particulièrement rentable et a sauvé la vie à quelque deux millions d'enfants âgés de moins de cinq ans.
- L'appui fourni par la Banque mondiale dans les secteurs autres que la santé a contribué à améliorer la situation en termes de santé infantile.
- De faibles sommes d'argent sauvent des vies... quoique le montant en jeu varie sensiblement selon l'opération considérée.
- Bien que des opérations dans plusieurs secteurs influent sur la situation de la santé maternelle et infantile, cela ne signifie pas pour autant que des opérations plurisectorielles s'imposent en permanence.
- L'appui de la Banque mondiale aux activités de formation des accoucheuses traditionnelles a permis une réduction de la mortalité néonatale... mais ce programme a désormais été abandonné, l'opinion internationale ayant évolué dans le sens d'un recours à des professionnels qualifiés pour les accouchements.
- Les programmes doivent reposer sur les réalités locales, plutôt que sur les idées généralement admises.
- La problématique hommes-femmes est un élément déterminant pour les stratégies adoptées en matière de santé au Bangladesh. Il convient de veiller davantage à rectifier les partis pris sexistes pour maintenir l'élan acquis sur le front de la baisse de la mortalité et du recul de la fécondité. Mais les comportements traditionnels

ne sont pas l'entrave absolue aux prestations de services que certains suggèrent parfois.

- Le BINP entrepris avec l'aide de la Banque a amélioré l'état nutritionnel de la population, mais bien moins que prévu. Il y a lieu de prêter sérieusement attention aux moyens d'améliorer ce programme en termes d'efficacité et d'efficience—ou, si cela n'est pas possible, d'envisager alors des solutions de rechange au passage à plus grande échelle.
- Une étude d'impact rigoureuse peut mettre en évidence les programmes gouvernementaux et appuis extérieurs qui contribuent le plus à la réalisation des objectifs de réduction de la pauvreté.
- Les enquêtes nationales peuvent être utilisées pour les besoins de l'évaluation, mais leurs potentialités pourraient être renforcées moyennant certains ajustements, notamment un questionnaire communautaire plus détaillé.



Resumen

El mejoramiento de la salud y la nutrición maternoinfantiles es indispensable para las metas de desarrollo. La importancia de esos objetivos se refleja en su inclusión en metas de reducción de la pobreza como, por ejemplo, los objetivos de desarrollo del milenio (ODM) y el documento provisional de estrategia de lucha contra la pobreza de Bangladesh, que cuentan con el respaldo de los principales asociados para el desarrollo, incluidos el Banco Mundial y el Departamento para el Desarrollo Internacional del Reino Unido (DFID, por sus siglas en inglés).

En el presente informe se examina qué pueden hacer los programas que reciben ayuda pública y asistencia externa del Banco y otros organismos para acelerar el logro de metas como la reducción de la mortalidad infantil en dos terceras partes. La información facilitada en el informe se refiere a Bangladesh, un país que ha avanzado de forma espectacular, pero que necesita conservar el impulso para alcanzar sus propias metas de reducción de la pobreza.

En el informe se tratan las siguientes cuestiones: 1) ¿Cómo han evolucionado los resultados en materia de salud y nutrición infantiles y de fecundidad en Bangladesh desde 1990? ¿Comparten los pobres los progresos realizados?; 2) ¿Cuáles han sido los principales factores determinantes de los resultados en salud maternoinfantil en Bangladesh a lo largo de ese período?; 3) En vista de esos factores, ¿qué puede decirse de los efectos de los programas que reciben apoyo público y exterior

—especialmente del Banco Mundial y del DFID— para mejorar la salud y la nutrición?; 4) En la medida que las intervenciones han propiciado efectos positivos, ¿lo han hecho de forma rentable?

Tendencias en la mortalidad de niños menores de cinco años, la fecundidad y la nutrición

A pesar de un inicio poco prometedor como consecuencia de la guerra y la hambruna, Bangladesh ha progresado espectacularmente en las dos últimas décadas, sobre todo en lo concerniente a la disminución de la fecundidad. Contrariamente a la idea más extendida, la fecundidad mantuvo su tendencia descendente en el decenio de 1990. La mortalidad de los niños menores de cinco años se ha reducido también a buen ritmo, siendo Bangladesh uno de los pocos países que ha alcanzado un nivel suficiente para cumplir el ODM de una reducción en dos terceras partes para 2015. La ex-

cepción a estos éxitos ha sido la nutrición. Las medidas físicas del estado nutricional sólo empezaron a registrar ligeras mejoras en los años noventa y la desnutrición permanece en niveles elevados.

Estos resultados más favorables han beneficiados a todos los ciudadanos. Aunque los niños de las familias pobres tienen más probabilidades de morir prematuramente, la brecha se va colmando, con tasas de mortalidad que descienden más rápido entre los pobres que en los demás sectores de la sociedad. El uso de anticonceptivos y los bajos niveles de fecundidad son también comunes entre los pobres.

Causas de la disminución de la mortalidad de niños menores de cinco años

El análisis de los factores determinantes de la mortalidad utilizando tanto datos de comparaciones entre países como de las Encuestas de Demografía y Salud (EDS) muestra que son diversos los factores que han influido en el descenso del número de muertes. Destaca el mayor bienestar económico como razón de la disminución de la mortalidad infantil, pero su influencia es menor en lo referente a lactantes. Se ha constatado que las intervenciones en el sector social—tanto en salud como en educación—desempeñan también un papel importante, y la mayor cobertura de inmunización, junto con un aumento en la matrícula femenina en la enseñanza primaria y secundaria, han repercutido sustancialmente en la reducción de la mortalidad. Los resultados revelan, asimismo, un pronunciado sesgo de orden sexual, pues la mortalidad femenina es más elevada, sobre todo en las divisiones de Sylhet y Chittagong. El análisis de algunas intervenciones indica lo siguiente:

- La cobertura de inmunización era inferior al 2% a principios de los años ochenta, pero aumentó en la segunda mitad del decenio (en gran medida gracias al apoyo del UNICEF, aunque posteriormente también de otros donantes como el Banco Mundial), de modo que para 1990 casi la mitad de los niños habían recibido todas las vacunas en sus primeros 12 meses de vida. La inmunización ha evitado más de dos millones de muertes infantiles en

los dos últimos decenios, con un costo por vida salvada situado entre US\$100 y US\$300.

- El Banco Mundial financió la capacitación de aproximadamente 14.000 parteras tradicionales hasta fines de los años noventa, momento en que se suspendió la capacitación a raíz de un cambio en la opinión internacional en favor de que todos los partos fueran asistidos por profesionales. Sin embargo, los datos presentados en este informe muestran que la formación dispensada a parteras tradicionales ha salvado la vida de muchos lactantes, con un coste de entre US\$220 y US\$800 por fallecimiento evitado.
- La matrícula femenina en la enseñanza secundaria experimentó un rápido aumento en el decenio de 1990, especialmente en zonas rurales, debido en parte a los subsidios para becas concedidos a todas las alumnas inscritas en los niveles 6 a 10 en dichas zonas con ayuda de Noruega, el Banco Asiático de Desarrollo, el Banco Mundial y las autoridades nacionales. Una de las ventajas de la mayor escolarización femenina en secundaria es el descenso de la mortalidad, con un coste de entre US\$1.080 y US\$5.400 por muerte evitada.
- La electrificación rural, respaldada mediante tres programas del Banco Mundial en los años ochenta y noventa, reduce la mortalidad mediante efectos de ingreso, mejorando los servicios sanitarios, facilitando la esterilización del agua y favoreciendo el acceso a información sobre la salud, especialmente por televisión. Si se tienen en cuenta estos diversos cauces, se observa que la tasa de mortalidad de los niños menores de cinco años en los hogares con suministro eléctrico es un 25 por mil menor que en los hogares que carecen de electricidad. Sobre la base de costos históricos, el total asciende a US\$20.000 por vida salvada y US\$40.000 en función de los costos de conexión actuales.

Nutrición

A fin de hacer frente a la deficiente situación nutricional, las autoridades pusieron en marcha a modo experimental, con ayuda del Banco Mundial, el Proyecto de Nutrición Integrado de Ban-

gladesh (PNIB). El eje central de dicho proyecto es el componente de nutrición de base comunitaria (CNBC), que fomenta el asesoramiento nutricional para instaurar un cambio de comportamiento, completado con alimentación suplementaria a mujeres embarazadas y niños de corta edad.

El análisis de la cadena causal desde los insumos del PNIB hasta los resultados antropométricos infantiles muestra lo siguiente: 1) existe un eslabón débil en la cadena, ya que la comunicación en pro de un cambio de comportamiento se ha centrado excesivamente en las madres, que a menudo no son quienes toman las decisiones relacionadas con todas las prácticas alimentarias; 2) la cobertura del programa es, por lo general, elevada en las zonas del proyecto, pero mucho menor en *thanas* (subdistritos) más conservadores, en especial entre las mujeres que viven con su suegra; 3) se observan algunas deficiencias en la definición de los grupos de beneficiarios: a) se aplicaron criterios demasiado estrictos para la admisión de niños desnutridos en la alimentación suplementaria, mientras que se admitió a niños con retraso de crecimiento pero probablemente bien alimentados; b) en la alimentación de mujeres embarazadas se excluyó a muchas que cumplían las condiciones de admisión mientras que se aceptó a otras que no los cumplían; 4) un porcentaje elevado de madres cuyos hijos recibían alimentación suplementaria señalaron que nadie les daba consejos sobre nutrición; 5) existe una brecha considerable entre los conocimientos y la práctica que hace que las mujeres no apliquen los consejos que reciben: ello obedece, principalmente, a la falta de recursos económicos y tiempo, y 6) los efectos sobre el aumento de peso durante el embarazo son demasiado reducidos para que repercutan en el peso al nacer, lo cual sucede frecuentemente en programas similares en otros países; el estado nutricional de la mujer antes del embarazo incide de forma más acentuada en el bajo peso al nacimiento que el aumento de peso durante el embarazo y, por ende, podría haber sido más acertado ocuparse de ese aspecto en el proyecto.

La lista del párrafo anterior puede considerarse como una lista de problemas que deben solucionarse en el proyecto, pues hasta cierto

punto incumben al Proyecto Nacional de Nutrición ampliado; se han revisado los criterios de selección de niños para la alimentación complementaria y se ha intentado de nuevo transmitir consejos sobre nutrición a la población masculina. Ahora bien, el programa no ha sido un medio muy rentable de mejorar el estado nutricional, que ha mejorado en general con la aceleración de la disponibilidad de alimentos asociada al aumento del rendimiento de la producción arroceras desde finales de los años noventa y la consiguiente reducción del precio efectivo del arroz. Las simulaciones demuestran que repartiendo simplemente alimentos a las familias con niños se habría logrado un mayor impacto nutricional. El costo por vida salvada con la ración de arroz hipotética supera en poco los US\$2.000, la mitad del costo de las vidas salvadas por el PNIB.

Reducción de la fecundidad

La tasa de reducción de la fecundidad en Bangladesh es superior a la que cabría esperar de otros avances socioeconómicos, como el aumento del ingreso y la expansión de la educación femenina. Si bien los avances socioeconómicos, incluida la transición demográfica, explican parte de la rápida caída de la fecundidad en Bangladesh, una parte importante es atribuible al servicio nacional de planificación familiar, creado con considerable ayuda del exterior en los años que siguieron a la liberación en 1971. La disminución persistente de la fecundidad en los años noventa, movida por la creciente prevalencia de la anticoncepción, demuestra la eficacia constante de este programa.

El Plan Estratégico de Inversión en Salud, Nutrición y Población lanzado por las autoridades nacionales destaca la utilidad de aumentar la edad de matrimonio como medio para reducir la fecundidad, y diversos programas, incluido el asesoramiento brindado en el marco del PNIB, promueven un matrimonio más tardío. Una de las condiciones del programa de subsidios para que las niñas y jóvenes cursen estudios de secundaria, apoyado por el Banco y otras entidades, es que las beneficiarias permanezcan solteras. Es cierto que la edad de matrimonio en Bangladesh es baja y que la mitad de las jóvenes

se casan a los 14 años o antes. También es cierto que existe una pauta internacional bien establecida según la cual el aumento de la edad de matrimonio reduce la fecundidad. Ahora bien, no puede esperarse que esa pauta se siga en Bangladesh por dos razones: 1) el aumento de la edad de matrimonio de niñas de 13 años o menos no incide en la edad a la éstas que tienen su primer hijo (la edad en el momento del matrimonio aumenta, pero el lapso entre el momento del matrimonio y el nacimiento del primer hijo disminuye), y 2) si una mujer ha previsto no tener más de tres ó cuatro hijos, como es el caso de la mayoría de las mujeres en Bangladesh, puede lograrlo aunque los embarazos comiencen a la edad de 15 ó 20 años. El efecto directo de extender la educación secundaria será escaso, puesto que Bangladesh ha alcanzado ya niveles de fecundidad equiparables a los de países con niveles de educación más elevados. Por lo tanto, el aumento de la edad de matrimonio, aunque es deseable tanto para la salud materna como para la infantil (los niños nacidos de madres adolescentes tienen mayor probabilidad de muerte prematura), influirá poco en el número de niños nacidos de cada mujer en sus años reproductivos; de todos modos, habría un efecto temporal de ritmo en la tasa total de fecundidad y un efecto de segundo orden, ya que la reducción de la mortalidad en nacimientos más tardíos reduciría el número deseado de hijos. Sería, pues, conveniente centrarse en las familias que exhiben una alta fertilidad para, entre otras cosas, intentar restablecer el uso permanente de medidas anticonceptivas en sus niveles anteriores. Asimismo, deberían desplegarse esfuerzos para abordar la preferencia por hijos varones, la cual dificulta la disminución de la fecundidad. Si prosigue la reducción de la mortalidad, disminuirá también la fecundidad.

Enseñanzas aprendidas

Del análisis efectuado en el presente informe se desprenden las siguientes enseñanzas generales:

- Las intervenciones que cuentan con apoyo exterior han tenido un notable impacto en los resultados de salud materno-infantil en Ban-

gladesh. La vacunación ha resultado especialmente eficaz en función de los costos y salvado la vida de hasta dos millones de niños menores de cinco años.

- El apoyo del Banco Mundial a sectores distintos del sanitario ha permitido obtener mejores resultados en el ámbito infantil.
- Con pequeñas cantidades de dinero se salvan vidas... aunque los montos varían significativamente de una intervención a otra.
- Aunque las intervenciones en muchos sectores afectan a los resultados en materia de salud materno-infantil, ello no implica necesariamente que se requieran siempre intervenciones multisectoriales.
- El apoyo del Banco Mundial a la capacitación de parteras tradicionales ha reducido la mortalidad neonatal... pero el programa ya se ha suspendido siguiendo la tendencia internacional a apoyar la formación de parteras profesionales.
- Los programas deberían basarse en datos locales, en lugar de la sabiduría popular general.
- Las cuestiones de género son fundamentales en las estrategias de salud en Bangladesh. Es necesario concentrarse más en corregir los sesgos de género para mantener el impulso cobrado en la disminución de la mortalidad y la reducción de la fecundidad. Pero las actitudes tradicionales no son la máxima cortapisa en la prestación de servicios, como a veces se sugiere.
- El PNIB del Banco ha mejorado el estado nutricional, pero no por mucho menos de lo previsto. Es preciso prestar especial atención a las formas de mejorar tanto la eficacia como la eficiencia del programa o, en caso de que ello no sea posible, considerar las alternativas a la ampliación.
- Si se evaluarán las consecuencias rigurosamente se podría conocer qué programas públicos y asistencia externa contribuyen en mayor medida a alcanzar los objetivos de reducción de la pobreza.
- Es posible recurrir a encuestas nacionales con fines de evaluación, pero algunos ajustes las harían más útiles, en particular un cuestionario comunitario más detallado.

Acronyms and Abbreviations

ANC	Antenatal care
BCG	Bacillus Calmette-Guerin
BDHS	Bangladesh Health Survey
BFS	Bangladesh Family Survey
BINP	Bangladesh Integrated Nutrition Project
BMI	Body-mass index
BRAC	Bangladesh Rural Advancement Committee
BRSFM	Bangladesh Retrospective Survey of Fertility and Mortality
CAS	Country Assistance Strategy
CBNC	Community-Based Nutrition Component
CNP	Community Nutrition Promoter
CPS	Contraceptive Prevalence Survey
DFID	Department for International Development (U.K.)
DHS	Demographic and Health Survey
DPT	Diphtheria, pertussis, and tetanus
EOC	Emergency obstetric care
EPI	Expanded Program of Immunization
FHV	Family health visitor
FSSAP	Female Secondary School Stipend Program
FWA	Family welfare assistant
FWC	Family Welfare Center
GoB	Government of Bangladesh
GRR	Gross Reproduction Rate
HA	Health assistant
HAZ	Height-for-age Z score
HIES	Household income and expenditure survey
HKI	Helen Keller International
HPSP	Health and population sector project
HPNSP	Health, population, and nutrition sector project
IEC	Information, education, and communication

IPRSP	Interim Poverty Reduction Strategy Paper
IUD	Intra-uterine device
JFS	Joint Finance Scheme
KP	Knowledge and practice
MCH	Maternal and child health
MDGs	Millennium Development Goals
MOFHW	Ministry of Family, Health and Welfare
NGO	Nongovernmental organization
NIDS	National Immunization Days
NNP	National Nutrition Program
NORAD	Norwegian Agency for Development
NSP	Nutritional Surveillance Project
OED	Operations Evaluation Department (World Bank)
PCR	Project Completion Report
PHP	Population and Health Project
PPS-BD	Participatory Practitioner's Society–Bangladesh
READ	Research Evaluation Associates for Development
SIDA	Swedish International Development Agency
TBA	Traditional birth attendant
TFR	Total fertility rate
TMSS	Thengamara Mohila Sabuj Sangha
TTBA	Trained traditional birth attendant
UHZ	Upazilla Health Complex
UNFPA	United Nations Population Fund
UNICEF	United Nations Children's Fund
USAID	United States Agency for International Development
SAR	Staff Appraisal Report
WAZ	Weight-for-age Z score
WDI	World Development Indicator
WHO	World Health Organization
WHZ	Weight-for-height Z score

Note: Dollars amounts cited are in U.S. dollars unless stated otherwise.



Maternal and Child Health in Bangladesh: A Record of Success

Improving maternal and child health is central to the development challenge. Bangladesh has a remarkable record in the reduction of fertility and under-five mortality, but has less impressive achievements with respect to nutrition. This study examines the impact of interventions from various sectors—health, population, nutrition, education, and electrification—on these outcomes. Will existing interventions be adequate to maintain momentum toward the achievement of poverty-reduction goals, or are changes required?

Improving maternal and child health (MCH) and nutrition is central to meeting the development challenge. This is reflected in the incorporation of MCH outcomes in development goals. Two of the eight Millennium Development Goals (MDGs) refer to MCH outcomes—reducing under-five and maternal mortality—and child malnutrition is an indicator for the first MDG. Four of the 10 goals in Bangladesh's Interim Poverty Reduction Strategy Paper relate to MCH. These four are to (1) reduce infant and under-five mortality rates by 65 percent and eliminate gender disparity in child mortality; (2) reduce the proportion of malnourished children under five by 50 percent and eliminate gender disparity in child malnutrition; (3) reduce the maternal mortality rate by 75 percent; and (4) ensure reproductive health services for all (Bangladesh 2003, pp. 7–8). These goals are reflected in the Bank's Country Assistance Strategy (CAS): human de-

velopment is listed as the first development priority, and the second of the four aims in the CAS is to “consolidate gains in human development, addressing development challenges in education, health, and nutrition” (World Bank 2002b).

Adoption of a results-based approach requires understanding the main drivers behind changes in target outcomes. Which publicly supported interventions can accelerate the pace of improvement, and so secure the achievement of development goals? Have the interventions supported by the Bank and the Department for International Development (DFID) helped them meet the goals of their respective country strategies? This report addresses these questions in the context of Bangladesh, a country that has made notable progress but needs to

Improving maternal and child health and nutrition is central to meeting the development challenge.

The decline in infant and child mortality will enable Bangladesh to be one of the few countries to meet the MDG of a two-thirds reduction by 2015. Achievements in fertility reduction are even more remarkable.

maintain momentum in order to achieve its own poverty-reduction goals. The following outcomes are analyzed: infant and child mortality, child nutrition, nutritional status of pregnant women and low birth weight, and fertility (see box 1.1). Maternal mortality is excluded on account of lack of

data.¹ However, fertility is included as a known correlate of maternal mortality, as well as a correlate to child health and nutrition.

The years immediately following Bangladesh's liberation in 1971 were inauspicious ones for the

prospects for development. Emerging from a violent struggle that cost more than one million lives, the country was beset by a famine that was responsible for at least another quarter of a million deaths. Nearly one in four children was dying before reaching their fifth birthday, reflecting the highest under-five mortality rate in the region, and one of the highest in the world.²

But by the start of the 1980s the situation had changed. The decline in infant and child mortality, which began slowly in the 1960s, accelerated (figure 1.1).³ By 2000, under-five mortality had fallen to 82 per 1,000 live births. This rate is now below that of neighboring countries, and the rate of decline is sufficient for Bangladesh to be one of the few countries on track to meet the Millennium Development Goal of a two-thirds

BOX 1.1

Measures of Welfare Outcomes

Mortality

The three mortality measures are:

- **Infant mortality rate (IMR):** the probability of death before an infant's first birthday, usually expressed per 1,000 live births
- **Child mortality rate (CMR):** the probability of death between a child's first and fifth birthdays, expressed per 1,000 live births
- **Under-five mortality rate (U5M):** the probability of death before a child's fifth birthday.

Anthropometric measures

The nutritional status of children under five is often monitored through measurements of their height, weight, and age. These three pieces of information are used to calculate three ratios:


- **Height for age (stunting):** a measure of long-run nutritional status
- **Weight for height (wasting):** a measure of short-run nutritional status
- **Weight for age (underweight):** a combination of the above two measures.

Nutritional status is determined by converting a measure to a z score—that is, subtracting the mean and dividing by the standard deviation (SD) from a reference population. The corresponding statistics are called the height for age z score (HAZ), weight for height z score (WHZ), and weight for age z score (WAZ). Values of less than -2 (2 SDs below the reference mean) are taken to indicate malnutrition; less than -3, severe malnutrition.

For adults, the appropriate measure is the body mass index (BMI), which is equal to a person's weight (in kilograms) divided by the square of their height in meters. A BMI of less than 18.5 indicates that a person is underweight.

Fertility

The most common measure, the total fertility rate (TFR), is the number of children a woman would have if her childbearing equaled the age-specific fertility rates of women currently in those age groups. That is, TFR is a cohort measure. In the presence of declining fertility, a woman entering her childbearing years is expected to have fewer children than the TFR, whereas the average number of children born to women who have completed their reproductive years will exceed the TFR.

FIGURE 1.1**Both Under-Five Mortality and Fertility Have Fallen Rapidly**

Source: Annex A.

reduction between 1990 and 2015. The country's achievements with respect to fertility are even more remarkable. Fertility was very high, and possibly even rising, in the 1960s and early 1970s, with each woman bearing an average of seven live births. A rapid decline in fertility starting in the mid-1970s has more than halved the total fertility rate.⁴ There was less progress regarding nutrition, with improvements in anthropometric indicators occurring only in the 1990s. Maternal mortality also remains very high.

This report analyzes the factors behind the success story, focusing on the 1990s. The main questions addressed are:

- What has happened to child health and nutrition indicators and fertility in Bangladesh since 1990? Are the poor sharing in the progress being made?
- What have been the main determinants of MCH and nutrition outcomes in Bangladesh over this period?
- Given these determinants, what can be said about the impact of publicly supported programs to improve health and nutrition?
- To the extent that interventions have brought about positive impacts, have they done so in a cost-effective manner?

Scope of the Study

Maternal and child health and nutrition outcomes result from public policies and private actions in many sectors—not only health provision and individual choices, but also income generation, education, and infrastructure. This study focuses on a few selected interventions supported by the World Bank, DFID, and other external agencies from both within and outside the population and health sector. The interventions have been selected on the grounds of (1) their demonstrated importance for health and nutrition outcomes and (2) the involvement of external partners in supporting service provision.

Health, nutrition, and population programs are clear candidates for inclusion in the evaluation on grounds of both an expected impact on MCH-related outcomes and the level of externally supported public provision of services. Bangladesh has benefited from a high degree of donor coordination in the health and population sector under the auspices of five World Bank credits since 1975, with substantial levels of cofinancing from bilateral agencies, including those in the United Kingdom.⁵ The first three of these projects focused on establishing the country's reproductive health system, but the fourth and fifth projects expanded coverage to all aspects of

Bangladesh has benefited from a high degree of donor coordination in the health and population sector under the auspices of five World Bank credits since 1975, with substantial levels of cofinancing from bilateral agencies.

example, DFID health spending in this period was dominated by UK£25 million denoted as “World Bank time-slice financing.”

Preparations are under way for the follow-up Health, Population and Nutrition Sector Project (HPNSP), with further expansion in coverage. While nutrition activities were partly included in earlier health projects, a stand-alone pilot project, the Bangladesh Integrated Nutrition Project (BINP) was initiated in 1996. BINP, and the follow-on National Nutrition Programs (NNP), embody a behavior-change approach to improving nutritional status, an approach of growing importance in various areas of health care. This study re-examines the data from recent evaluations of BINP to determine the effectiveness of this approach.

Evidence of impact on MCH outcomes from two other subsectors is also assessed: female secondary schooling and rural electrification. Female education is a well-established correlate of child health and nutrition, a link confirmed by the empirical analysis in this study (Chapter 4 and Annexes C and D). The World Bank has been one of the agencies supporting the female secondary school stipend program (FSSAP), which is shown to have made a substantial contribution to the rapid growth in secondary enrollments. Electrification appears as a determinant of mortality in many studies, although the channels for this effect are not so well-documented. Together with other agencies, the World Bank supported rural electrification in Bangladesh through three projects in the 1980s and 1990s,⁶ and this link thus warrants further investigation.

health care delivery. This study is mainly concerned with the period covered by the Fourth Population and Health Project (PHP4, 1992–98) and the Health and Population Sector Project (HPSP, 1998–2004). HPSP marked the transition from coordinated project support to sector program financing (sector budget support). For ex-

It is worth mentioning some areas that have been excluded from the study, despite their importance to MCH outcomes: income-generating activities, disaster prevention and relief, and economic infrastructure. The many interventions whose main impact on health is through their income-generation effects have been excluded from the analysis on two grounds. The first is that estimating income effects of interventions is a sizeable task in itself, and not one to which the data strategy adopted for this study is well-suited. The second reason is that while economic growth explains part of Bangladesh’s remarkable progress in improving social outcomes, it is not, as is shown in Chapter 4, the whole explanation.

A second area excluded from this study is that of disaster prevention. Bangladesh is subject to repeated disasters, most notably flooding. A substantial amount of external support has gone into measures to prevent flooding and to protect people from its consequences, as well as to fund relief and rehabilitation in the wake of such events.⁷ The success of these efforts, and similar programs funded from other sources, is shown by the reduced death toll from natural disasters.⁸ These flood control measures should be important to preserving livelihoods (though some early Bank-supported projects had the opposite effect) and saving lives. But the number of deaths directly attributable to flooding, at less than 2,000 in each disaster year, is relatively small compared with the approximately 300,000 under-five deaths each year.⁹

Another area of great importance, but not considered in this study partly for the reason income-generating activities are excluded (difficulty in tracing the causal chain with the data at hand), is economic infrastructure. Large-scale infrastructure, such as the DFID-supported renovation of the Chittagong port, have helped spur the rapid expansion of the garment industry, which has brought both economic growth and changes in the position of women. Other large projects, such as the Jamuna Bridge, have greatly reduced travel time across the country, promoting social and economic integration. Smaller-scale rural roads, and particularly bridges, foster

the integration of remote areas, facilitating access to markets and other services.¹⁰

Evaluation Approach

This study draws on the analysis of a number of existing data sets. Demographic and Health Survey (DHS) data are used to model the determinants of child health and nutrition outcomes, and of fertility. Bangladesh has had three such surveys, in 1992, 1996, and 1999,¹¹ allowing an analysis of the relative importance of the various determinants in explaining improved outcomes in the 1990s. This analysis informed the choice of sectors to be included in the evaluation.

The impact of the interventions is estimated through a structural modeling approach. That is, multivariate estimates are made of the outcomes of interest (mortality, nutrition, and fertility), drawing on the state of the art in the literature. The endogeneity of behavioral factors, such as antenatal care and immunization, is controlled for through the use of appropriate instruments, and the selection bias from children who have died not being in the sample for the nutrition equation allowed for by a two-part sample selection model. Further details on methodology can be found in the relevant Annexes (notably C, D, and G) and in the Approach Paper (Annex J). Combining the marginal impact of different interventions on welfare outcomes with cost data facilitates the cost-effectiveness analysis.

The analysis of nutrition data is a partial exception to the above approach. The Bangladesh Integrated Nutrition Project (BINP) conducted a survey for evaluation purposes. This study presents a reanalysis of these data, modifying the control group using propensity score-matching by drawing on the nationally representative Nutritional Surveillance Project carried out by Helen Keller International. A theory-based approach was adopted so that the causal chain by which inputs are intended to affect outcomes could be examined in detail.

The main thrust of this approach is heavily quantitative. But any impact evaluation needs to

place itself in context. Qualitative information is relevant to the analysis in various ways. First is the historical political context. Bangladesh has historically had a strong commitment to basic health, introducing an essential drugs policy in the face of substantial opposition; but in recent years there has been a poor record of delivering essential health at the field level (see Chapter 2). Second is the changing role of women, and the way in which *purdah* (social restrictions on women) may restrict their ability to access health services. A third factor is the importance of the private sector, although this study focuses on services that have been provided by government with external support.

The findings from this study can be related to OED's evaluation criteria of relevance, efficacy, and efficiency. This chapter has already illustrated the relevance of interventions intended to improve maternal and child health and nutrition to Bangladesh's poverty-reduction goals. The efficacy of the interventions depends on establishing a link from supported activities to welfare outcomes, as is done in Chapters 4 and 5. Efficiency is determined using cost-effectiveness analysis, which is also reported in Chapters 4 and 5.

Overview of the Report

Chapter 2 presents background information on the evolution of health and family planning services in Bangladesh and the role of external agencies in supporting these programs. The record with respect to mortality, fertility, and nutrition is presented in Chapter 3. Chapters 4 and 5 examine the impact of selected interventions on health, nutrition, and fertility outcomes. Chapter 6 concludes with lessons learned.

Bangladesh has historically had a strong commitment to basic health, introducing an essential drugs policy in the face of substantial opposition; but in recent years there has been a poor record of delivering essential health at the field level.



Health, Family Planning, and Nutrition Services in Bangladesh: An Overview

Following independence, Bangladesh established a substantial network of health and family planning facilities. This network, including staff costs, was largely financed by donor assistance during the first decade. Although facilities were created for both health and family planning, the focus of service delivery until the 1990s was on family planning, with the exception of immunization, for which a successful campaign was launched in the mid-1980s. The family planning services were built on a home visit system; similarly, immunization operates through 8,000 outreach centers around the country. Plans to move to a fixed-site system, for which construction of community clinics has been undertaken at donor expense, have met with very limited success. A pilot Integrated Nutrition Project was launched in 1996, which is now being rolled out nationwide.

Family Planning Programs

Promotion of family planning began with the volunteer Family Planning Association in what was then East Pakistan in 1952. A national family planning program was adopted in 1965, setting up a network of family planning clinics run by the Family Planning Board. The program used incentives for both personnel and clients, promoting mainly sterilization and intrauterine devices (IUDs). Success was limited, with fewer than 10 percent of couples using contraception in 1969, though awareness of modern methods increased markedly. As a result of limited acceptance, fertility remained high, with a total fertility rate (TFR) of

seven. Planned improvements in the program were interrupted by the liberation war in 1971. The newly independent government recognized population as one of the country's most pressing problems: by that time the country already had the highest population density in the world.

The Bangladesh government's efforts to reduce population growth cannot be separated from those of external agencies, since

The newly independent government recognized population as one of the country's most pressing problems: by that time the country already had the highest population density in the world.

This project was the first cofinanced social sector project undertaken by the World Bank anywhere in the world.

these agencies were central from the outset, paying the majority of program costs, including salaries, in the early years. The country's first Five-Year Plan proposed a health and family planning system with a hospital in each district,¹ which in some cases was supplemented by a maternal and child welfare center at the subdistrict level. Each thana was to have an upazilla health complex (UHC), and each union a family welfare center (FWC). Each FWC is headed by a medical assistant, working with a family welfare visitor (FWV) and a pharmacist. The FWC's staff consists of a female family welfare assistant (FWA) and a male health assistant (HA). The FWV conducts satellite clinics, the FWA makes home visits, and the HA has responsibility for malaria and epidemic control and environmental sanitation.

The World Bank was invited to participate in implementation of the plan and first fielded a mission in May 1973. Donor coordination was envisaged from the outset. However, disagreement arose with the U.S. Agency for Inter-

national Development (USAID) over whether family planning should fall under a separate organization or be integrated with health services, and with the United Nations Population Fund (UNFPA) over donor leadership. Hence, USAID and UNFPA signed their own agreements with the government; but six bilateral agencies (Australia, Canada, Germany, Norway, Sweden, and the United Kingdom) cofinanced the First Population Project (1975–82, table 2.1), with the World Bank, with the objective of assisting the development of a comprehensive fertility-control program. This project was the first cofinanced social sector project undertaken by the World Bank anywhere in the world, and began at a time when the Bank was new to population issues. At the outset, there was little consensus—for example, on the importance of tackling supply or demand and the appropriateness of sterilization encouraged with cash payments.

The First Population Project set about creating the infrastructure needed for service delivery. Sixty-nine percent (US\$31.7 million) of project costs were for civil works—half of that amount for the health complexes (UHCs), and

TABLE 2.1

World Bank Credits for Health, Population, and Nutrition

Project	Years	Project costs (\$ million) ^a	Loan amount (\$ million) ^a	Cofinanciers
First Population Project	1975–82	45.70	15.0	Australia, Canada, Germany, Norway, Sweden, United Kingdom
Second Population and Family Health Project	1979–85	89.90	30.8	Canada, Germany, Norway, Sweden ^b
Third Population and Family Health Project	1986–92	246.40	100.9	Australia, Canada, Germany, Netherlands, Norway, United Kingdom
Fourth Population and Health Project (PHP4)	1992–98	756.30	188.4	Australia, Canada, European Union, Germany, Netherlands, Norway, Sweden, United Kingdom ^c
Bangladesh Integrated Nutrition Project (BINP)	1996–02	65.74	58.6	None
Health and Population Sector Program (HPSP)	1998–04	2815.90	250.0	Canada, European Union, Netherlands, Sweden, United Kingdom
National Nutrition Project (NNP)	2000–05	124.50	92.0	Canada, Netherlands

a. Actual amounts other than HPSP and NNP, which are appraisal estimates.

b. The United Kingdom was to cofinance but decided to not participate.

c. Belgium was to cofinance but decided to not participate.

the bulk of the remainder for Family Health Visitor (FHV) training schools. On the software side, most of the funds were used for FWA training and for salaries for 11,700 FWAs (project savings realized through the depreciation of the taka were largely used for this purpose). By 1978, 16,700 FWAs had been posted, with UNFPA paying salaries for most of those not supported by Bank finance. Although the bulk of funds went for health and family planning purposes, the First Population Project had a complex design with several multisectoral components. The Ministry of Health and Population Control was responsible for the largest part of the project, but another six ministries were also involved: Local Government, Rural Development and Cooperatives; Labor and Social Welfare; Agriculture; Education; and Information and Broadcasting. In part, these ministries implemented programs for spreading contraceptive knowledge—for example, through agricultural extension workers. However, they also carried out other activities such as income generation for women, since increasing the status of women was recognized from the outset as important in reducing fertility.

During preparations for the second Five-Year Plan, the government invited all donors to participate as technical advisors in drawing up the family planning strategy, and prepared for a follow-on population project, subsequently named the Second Population and Family Health Project. Once again, about half of project costs went to civil works, mainly UHCs and family welfare centers. The third population project continued this pattern. Although the projects paid the salaries of field staff, government was assuming a larger share. Under the first two projects, government had paid only 10 percent of project costs, but its share rose to 17 percent by the third project, and would be 28 percent in the fourth project.

The first three population projects established the planned health infrastructure in much of the country. By 1990 there were more than 2,700 upazilla health centers and family welfare centers, compared with 147 in 1975, when the first project began. There were more than 21,000 family welfare assistants in place, and the distribution of contraceptives had grown dramatically. However,

even toward the end of the 1980s, the program was still not widely recognized as a success. While the achievements in output delivery were recognized (although with project delays due to civil works problems), fertility outcomes were perceived as insufficient. An OED project assessment report of the first two projects, produced in 1989, did not give the impression that there was any great success story in Bangladesh's experience. Problems were attributed to both design (too great a focus on permanent rather than temporary contraceptive methods) and implementation, in particular the poor quality of training. The report also noted continuing problems with the integration of health and family planning, and reported that managerial capability had been overstretched by the complex multisectoral design.

However, a series of surveys started telling a different story. The Bangladesh Family Survey (BFS) of 1989 showed a steep decline in the TFR, from 6.8 in 1979 to 4.6 in 1988. The CPSs of 1989 and 1991, as well as the registration scheme of the Bangladesh Bureau of Statistics, revealed a similar trend. World Bank documents started proclaiming the success. An OED report from 1991 noted that there was incontrovertible evidence of fertility decline and that Bank-financed projects had contributed to that trend.² The project completion report (PCR) for the third population project noted that the progress was remarkable and the contribution of the Bank's projects both substantial and undeniable. Most notably, Cleland and others (1994) argued that fertility reduction had been achieved as a result of the program, despite an inhospitable socioeconomic setting. Others have challenged this conclusion, arguing that there has been socioeconomic change that can account for falling fertility.³

Increasing the status of women was recognized from the outset as important in reducing fertility.

Even toward the end of the 1980s, the population projects were not widely recognized as a success—but a series of surveys started telling a different story. World Bank documents started proclaiming the success.

Subsequent projects have supported the continued expansion and implementation of the family planning program, although the focus on family planning has declined as greater emphasis has been placed on health services. This changing focus was reflected in a broadening of objectives; for example, the Third Population and Family Health Project was to assist government in achieving not only its fertility goals, but also those for reducing infant and maternal mortality. Two policy issues have remained contentious between the government and donors. The first is the functional relationship of health and family planning.

Two policy issues have remained contentious between the government and donors: the functional relationship of health and family planning and provision of fixed services versus a system of home visits.

The second is provision of fixed services versus a system of home visits. Government has been more inclined to keep health and family planning separate, and to maintain home visits, which donor critics see as reflecting the power of vested interests rather than a dedication to improved service delivery.

Health Services

The general view is that, given the emphasis on family planning, other health services in Bangladesh have been relatively neglected.⁴ While there are serious shortcomings in the delivery of public health services today, two points should be noted. First, Bangladesh led the world in establishing an essential drugs policy in the early 1980s (National Drugs Policy of 1982), despite opposition from the United States and, for a time, the World Bank (Chowdhury 1996). This policy helped create the large pharmaceutical sector in the country and to make essential medicines available at low cost. At that time, the Bangladesh government clearly had

Bangladesh led the world in establishing an essential drugs policy in the early 1980s, despite opposition from the United States and, for a time, the World Bank.

both the capacity and will to implement an imaginative and difficult health policy. Second, the immunization program, discussed below, is an example of a program successfully conducted using government staff.


These apparent successes, and that of the family planning program, stand in contrast to the failure of current efforts to deliver the essential service package through fixed sites. Community clinics have been constructed under HPSP, but usage rates are extremely low, and there has been no rise in the proportion of people utilizing clinics.⁵ This low rate may be partly attributed to problems of accessibility and restrictions on women's movement, but the main reason is the poor quality of service offered at the clinics, with drugs being in short supply. In union health centers many doctors simply do not take up their posts.⁶ These problems are manifestations of the governance issues that plague service delivery in Bangladesh.⁷

As with reproductive health, other health services have been assisted by a very broad array of donor interventions. Each World Bank project has had a large and growing number of subcomponents, reaching 66 subcomponents under HPSP. Similarly, DFID has simultaneously supported a large number of health-related interventions, reaching a maximum of 41 separate activities in 1999–00.⁸ Hence, any evaluation cannot possibly hope to meaningfully cover the full range of funded activities; indeed, it is difficult even to imagine how they can be supervised. This study focuses its attention on a small number of selected interventions—immunization and training of traditional birth attendants.

Immunization

The government immunization program began in 1979, providing immunization services from fixed centers. Recommended vaccinations were one shot of bacillus Calmette-Guérin (BCG) for tuberculosis; three shots of diphtheria, pertussis, and tetanus (DPT) vaccine; one shot of measles vaccine; and three oral doses of polio vaccine, all to be administered by 12 months of age. Funding was provided mainly by the United Nations Children's Fund (UNICEF), but it was insufficient and vaccines were frequently not available; in consequence, very limited progress was made in raising coverage, which remained well below 5 percent until the mid-1980s (figure 2.1).

The Expanded Program of Immunization (EPI) was launched in 1985 with financial sup-

FIGURE 2.1 Immunization Coverage of Children Aged 12–23 Months

Sources: Annual data from WDI; bars from DHS.

port from several donors, mainly the Swedish International Development Agency (SIDA) and UNICEF. The World Health Organization (WHO) provided substantial technical inputs. The program was revised to include community-level delivery through outreach centers, bringing about very rapid increases in immunization rates toward the end of the decade (figure 2.1). Continued improvements in coverage were made until 1995. During the second part of the 1990s, just over half of all children were fully vaccinated by 12 months. While the percentage of children fully vaccinated has not increased in recent years, the percentage of those having no vaccination at all has continued to decline, reaching less than 10 percent by the end of the 1990s.

In rural areas, immunization continues to be provided through outreach centers, complemented by services at district and subdistrict hospitals, union-level clinics, annual campaigns, and catch-up days to reach children who have missed doses. National immunization days were instituted during the 1990s to deliver two doses of polio vaccine a year. Immunization has been included under the essential services package developed for HPSP, according to which immunization will be made available from community

clinics rather than outreach centers, but this change has not taken place.

Support for immunization has become one of the largest externally funded health activities in the country. During the 1990s, external agencies paid about half the costs of the immunization program. The World Bank and UNICEF were the main supporters; others were USAID, WHO, Japan, and SIDA. From 1997 to 1998, SIDA stopped supporting immunization, instead paying the funds into the HPSP finance pool. Donors giving money to the pool, which includes

the UK£25 million from DFID, are indirectly contributing to the immunization program, which consumes just under 10 percent of Ministry of Family Health and Welfare (MOFHW) expenditure. Despite this, the government was concerned that the move to pool financing would reduce the funds available for the immunization program, and so the World Bank assured them that it would make up any shortfall. In addition, USAID and the United Kingdom capitalized the

Immunization coverage remained well below 5 percent until the mid-1980s . . . But by the end of the 1990s less than 10 percent of children had received no vaccination.

rotating fund for Bangladesh to utilize UNICEF's Vaccine Independence Initiative, which is used to purchase DPT vaccine.⁹ Finally, DFID has become the main agency financing the polio program post-eradication (see box 4.2).

Training TBAs

During the 1990s, traditional birth attendants (TBAs) attended over 60 percent of all births in Bangladesh.¹⁰ Training of TBAs was, until recently, a central element of community-based health programs and an important part of strategies for safe motherhood. The World Bank, under HFP III and IV, and UNFPA trained approximately 14,000 TBAs in Bangladesh. However, in its decennial year (1997), the Safe Motherhood Initiative¹¹ disavowed the approach. TBAs are mostly illiterate, and it was argued that they are able to understand little of the training they receive. Drawing on international evidence, the Initiative claimed that there was little or no evidence that training TBAs had any impact on maternal mortality—although the possible im-

impact on neonatal mortality (which is demonstrated below) received less attention. Hence, the Safe Motherhood Initiative is now built around the objective of all births being attended by skilled birth attendants—explicitly excluding trained TBAs—an

objective that has been included as an indicator for the Millennium Development Goals. Bangladesh has followed these international recommendations. Previous programs to train TBAs have been largely abandoned, with a new emphasis on making skilled attendants available.

Programs to train traditional birth attendants have been largely abandoned, with a new emphasis on making skilled attendants available.

Nutrition

While considerable progress had been made by the end of the 1980s in putting in place a family planning system, and areas of primary health were being developed, nutrition continued to suffer from neglect—despite high levels of malnutrition. The Bangladesh National Nutrition Council was founded in 1975 to oversee nutrition policy and coordinate the various activities being undertaken by the different ministries. These activities included (1) subsidized food supplements to vulnerable population groups under the Ministry of Relief and Rehabilitation; (2) homestead garden production implemented by the Ministry of Agriculture; (3) distribution of vitamin A capsules twice a year to children aged six months to six years; and (4) iron and folic acid tablets for pregnant women distributed through satellite clinics. Several NGOs have been active in nutrition, notably the Bangladesh Rural Advancement Committee (BRAC), which implemented a community-based nutrition project similar to that proposed by the Bank.

In place of this fragmented set of programs, the World Bank proposed a comprehensive approach to nutrition, at the heart of which was community-level nutritional counseling to encourage good nutritional practices among pregnant women and mothers of young children. A pilot program, the Bangladesh Integrated Nutrition Project (BINP), began in six thanas in 1996, expanding to 59 thanas by the time the project was completed in 2002. By that time the National Nutrition Program had begun, which is rolling the program out nationally. As a pilot project, BINP was subject to an intensive evaluation. These data are reexamined in Chapter 5.



Trends in Under-Five Mortality, Nutrition, and Fertility

This chapter discusses trends in under-five mortality, nutrition, and fertility. There is no dispute that mortality has fallen, and it is shown that the decline has benefited all population groups, although disparities remain. Nutrition has also improved, although, contrary to what is suggested by some data sources, reductions in malnutrition did not occur in the 1980s. Fertility continued to decline during the 1990s, which is contrary to the picture outlined by the direct measures of the total fertility rate, which suggest a stagnation during the past decade.

Patterns of Mortality Decline

The decline in under-five mortality has benefited all groups of the population, by age, wealth, gender, and location. But disparities between boys and girls, and different regions of the country, remain high. This section examines these patterns using data from the three DHSs.

During 1985–89 and 1995–99, under-five mortality fell by close to 40 percent, from over 150 deaths per 1,000 live births to less than 100 (figure 1.1). This decline took place for all age groups—neonates as well as children aged one to five¹—and for all income groups. Indeed, during the 1990s under-five mortality fell marginally faster for the poor than for the non-poor.²

However, there is a sex bias in Bangladeshi mortality rates. In most countries, male children are more likely to die between their first and fifth birthdays than are females. Bangladesh is an exception to this pattern: female children aged one

to four years are 33 percent more likely to die than are male children. This is one of the highest female-male child mortality ratios in the world.³

Marked regional variations in life chances also persist. A child is twice as likely to die before reaching his or her fifth birthday in Sylhet as he or she—but particularly she—is in Khulna.

Anthropometric Outcomes

Until the end of the 1980s there was scant improvement in nutritional status in Bangladesh: close to 70 percent of children were both stunted (HAZ) and wasted (WHZ) (see box 1.1 for definitions). However, during the 1990s

The decline in under-five mortality has benefited all groups of the population, by age, wealth, gender, and location. But disparities between boys and girls, and different regions of the country, remain high.

Malnutrition rates have fallen from extremely high levels of 70 percent in the early 1980s to still high levels of around 50 percent, with the bulk of the improvement taking place during the 1990s.

for age or weight for age (figure 3.1). Data from the Helen Keller International (HKI) Nutritional Surveillance Project show a slightly different picture of steady improvement since the early

1980s. But the HKI data cannot be used to construct a trend in this way, since they are only nationally representative since 1998. Before that date, the sample was biased toward disaster-prone areas. Hence, as expected, the malnutrition rates in those areas were higher than the national average given by the World Bank data (figure 3.2).⁴ From 1998, HKI and World Bank data coincide. The conclusion is thus that malnutrition rates have fallen from extremely high levels of 70 percent in the early 1980s to still high levels of around 50 percent, with the bulk of the improvement taking place during the 1990s.

This picture is clear from data reported in the World Bank's World Development Indicators, with little change in nutritional status during the 1980s in either height


for age or weight for age (figure 3.1). Data from the Helen Keller International (HKI) Nutritional Surveillance Project show a slightly different picture of steady improvement since the early

1980s. But the HKI data cannot be used to construct a trend in this way, since they are only nationally representative since 1998. Before that date, the sample was biased toward disaster-prone areas. Hence, as expected, the malnutrition rates in those areas were higher than the national average given by the World Bank data (figure 3.2).⁴ From 1998, HKI and World Bank data coincide. The conclusion is thus that malnutrition rates have fallen from extremely high levels of 70 percent in the early 1980s to still high levels of around 50 percent, with the bulk of the improvement taking place during the 1990s.


What Has Been Happening to Fertility?

Following the rapid decline in fertility during the 1980s, it was feared that there was a fertility plateau in the 1990s. Direct estimates of the total fertility rate (TFR) from DHS show it to have declined only slightly, from 3.4 to 3.3, between 1993 and 1996, and then remained at that level until 1999.⁵ If there is indeed a fertility plateau, it seems that current family planning efforts have stalled. While the strategy was sufficient to achieve a certain level of fertility decline (presumably by reaching "easy acceptors"), additional efforts have to be made to maintain the momentum toward the government's target of fertility at the replacement rate.

However, three arguments can be advanced to suggest that fertility decline was not halted in the 1990s.⁶ First, direct estimates of fertility are less reliable than indirect ones, and indirect estimates give a different picture. Second, even if the direct estimates are used, there are reasons to believe



Source: WDI (2004).



Sources: WDI (2004), HKI.

that fertility was underestimated in 1993–94 because of displacement in birth reporting. Third, if indirect estimates are used, they show that the direct estimates underestimate fertility (particularly in 1993–94, reinforcing the previous point), and that fertility did continue to decline in the 1990s. These arguments are considered in turn.

Direct Versus Indirect Estimates

Table 3.1 summarizes direct fertility estimates from various surveys, together with indirect estimates made using data from these surveys. Two points can be noted. First, the direct estimates are systematically lower than the indirect ones. Second, the trend in fertility decline is far less from the direct estimates than from the indirect ones. For example, both CPS 1983 and 1989 yielded direct TFR estimates of 4.9. But these results have not been used to argue that there was a fertility plateau in the 1980s. To the contrary, this is seen as the period of most rapid decline.

Underestimation of TFR in DHS, 1993–94

The argument that the direct estimate of TFR in the 1993–94 DHS was an underestimate is based on an apparent displacement of births—births reported as taking place more than three years ago were actually more recent. The evidence for this phenomenon comes from comparing the number of births reported in the three years preceding the survey with the number reported for the three years prior to that (that is, four to six years before the survey). Since they cover similar time periods, the ratio of these two amounts will be one, except that mortality will tend to force it below one, and declining fertility or postponed births will raise it above one.

But this ratio for the 1993–94 survey is greatly in excess of both one and the value of the ratio observed in the other surveys. In the earlier survey, 26 percent more births were reported for four to six years before the survey, compared with the period within three years before the survey. This ratio compares with just 4 percent for 1996–97 and a 5 percent reduction in 1999–00. Yet comparing the ratio for five-year periods, there is little difference between the 1993–94 and 1996–97 surveys. These data thus show that in 1993–94, children under three

were being reported as being older. The effect of this birth displacement is exactly the same as for the postponement of births. *Three arguments can be advanced to suggest that fertility decline was not halted in the 1990s.*

That is, it will reduce the reported total fertility rate. But since, in this case, the reduction comes from incorrectly attributing births to a later period, there is a downward bias of the direct fertility estimate for 1993–94.

Indirect Fertility Measures


Figure 3.3 presents indirect estimates of fertility using three different approaches, using BFS 1989 and the three DHSs. One set of estimates was also made using 1991 and 2001 census data (see Annex F).

The data shown here confirm that direct methods underestimate fertility. This gap is larger for 1993–94 than the surveys in the succeeding or preceding years, which *The data shown here confirm that direct methods underestimate fertility.*

TABLE 3.1		Fertility Decline Has Always Been Erratic Based on Direct Estimates, but Continued into the 1990s Using Indirect Ones	
Year	Survey	TFR estimates	
		Direct ^a	Indirect ^b
1974	Bangladesh Retrospective Survey of Fertility and Mortality (BRSFM)	4.8	7.3
1975	Bangladesh Family Survey (BFS)	5.4	7.4
1983	Contraceptive Prevalence Survey (CPS)	4.9	7.0
1985	CPS	4.6	6.5
1989	CPS	4.9	5.9
1989	BFS	4.6	5.4
1991	CPS	4.2	n.a.
1993/94	Bangladesh Demographic and Health Survey (BDHS)	3.4	4.3
1996/97	BDHS	3.3	4.0
1999/00	BDHS	3.3	3.7

a. Direct estimates are from the various survey reports.

b. 1974–89 from Cleland and others (1994), and 1993–99 from OED analysis.


FIGURE 3.3
Fertility Decline Continued in the 1990s According to a Range of Indirect Measures


Note: See Annex F for explanation.

supports the argument that birth displacement exacerbated underestimation in that year. The results also show that fertility continued to decline during the 1990s, albeit at a slower rate than had been the case from 1989 to 1993.

The evidence of continuing fertility decline is consistent with patterns of contraceptive knowl-

edge and use (figure 3.4). By the mid-1980s, virtually all women in Bangladesh had heard of modern contraception methods, although just under a third had ever used them. The proportion who have both ever used them, and are currently doing so, has continued to rise since the mid-1980s.

FIGURE 3.4
Knowledge of Modern Contraceptives Is Universal and Use Continues to Rise




Impact of Specific Interventions on Child Health and Fertility

Analysis of both cross-country and Bangladeshi household data shows the clear contribution of interventions from several sectors to improvements in maternal and child health outcomes. Immunization has averted up to two million child deaths since the launch of the Expanded Program of Immunization, and done so in a cost-effective manner. Training traditional birth attendants, which has been largely abandoned, is also shown to have been a cost-effective way of reducing neonatal deaths. From other sectors, female secondary schooling and rural electrification are shown to have had a significant impact on under-five mortality and fertility, although electrification is a much less cost-effective means for achieving these objectives.


Income Growth Accounts for Some, But Not All, Improvement in Outcomes

There is a well-established relationship between income and social indicators. Higher income is generally associated with reduced mortality, higher educational achievement, better nutrition, and so on. However, the rapid improvements in social outcomes achieved in Bangladesh are greatly in excess of what can be explained from income growth alone. The four graphs in figure 4.1 show this link between income and four social outcomes—under-five mortality, the total fertility rate, and the prevalence of stunting and underweight among under-fives—using data from a cross-section of 78 developing countries. Each data point represents the decade averages of income and the social outcome shown, using values

from the 1970s to the current decade, so that there are up to four observations for each country. Analysis using data for a specific country across time reveals a similar pattern (see, for example, Haddad and others 2003).

The solid line in each figure is the average relationship between income and the social outcome; that is “the fitted line.” In the 1980s, Bangladesh (indicated by the triangular data points, each labeled by its decade) lay above the average for under-five mortality and fertility, meaning that those indicators were

Immunization has averted up to two million child deaths since the launch of the Expanded Program of Immunization, and done so in a cost-effective manner.

FIGURE 4.1**Bangladesh's Improvement in Social Outcomes Is Greater than Can Be Explained by Economic Growth Alone**

Note: See discussion in the text for explanations.

This finding suggests that there have been important, non-income-related factors behind the improvement in mortality and fertility in Bangladesh.

worse than should be expected for a country at its income level. If these indicators had improved following the internationally established relationship with income, then subsequent observations for Bangladesh would have lain along the dashed line.¹ But these later observations lie below the fitted line, showing that Bangladesh now does better than expected for a country at its income level. This finding suggests that there

have been important, non-income-related factors behind the improvement in mortality and fertility in Bangladesh.

In the case of the nutrition indicators, the observations from all three decades lie above the fitted line. Bangladesh continues to have worse anthropometric outcomes than the average for countries at a similar income level. But this discrepancy has narrowed over time. The improvement in all four outcomes greatly exceeds that in gross national product (GNP) per capita, and exceeds the expected improvement derived for growth in national income.

The numbers behind these graphs provide an upper estimate of the extent to which growth in gross domestic product (GDP) per capita has contributed to improved social outcomes in Bangladesh (table 4.1; see Annex B for more details).² For example, under-five mortality was 205 per 1,000 live births in the 1980s. Income growth alone would have reduced it to 163 by 2000, but by then the actual rate was 78. Hence, at most, just under one-third of the improvement comes from higher average income. For the other outcomes shown in figure 4.1, the share of income is even less, explaining, for example, at most, 16 percent of the reduction in fertility.

Further analysis (contained in Annex B) of the cross-country data examines the underlying factors behind changing outcomes in Bangladesh other than GNP per capita.³ This analysis shows that:⁴

- The largest single factor is immunization, which accounts for about one-third of the reduction in under-five mortality.
- Increasing female literacy is an explanatory factor for all four variables, accounting for between 5 and 15 percent of the observed change.
- Improved daily energy supply accounts for 10 to 15 percent of improved nutrition.
- Safe water has also contributed to improved nutrition, accounting for close to one-quarter of the reduction in stunting.

Two general conclusions emerge from these results. First, interventions from different sectors have all contributed to improved MCH outcomes. Not only health interventions (immunization) but also safe water, education, and agricultural output have all played a part. Analysis of additional data (see Annex B) also shows

electrification to be a fairly robust determinant of mortality outcomes. Second, there is a residual (unexplained) element in these figures, most notably for weight for age and the fertility rate. The remainder of this chapter builds on these

results to further analyze the factors behind improved MCH outcomes in Bangladesh and the extent to which public interventions have brought about these improvements.

Under-Five Mortality

The reasons for premature death vary by age.

For very young children, factors related to maternal health during pregnancy and the birth itself are important, with other aspects of health care having an impact throughout infancy.

But for children over one year, the general socioeconomic environment matters more. It is thus common to analyze mortality determinants separately for the different age groups. Such an analysis was conducted by OED using DHS data from the three surveys carried out in the 1990s (see Annex C for detailed results).

Table 4.2 summarizes the significant determinants of the mortality for

The largest single factor is immunization, which accounts for about one-third of the reduction in under-five mortality, but safe water, education, and agricultural output have also played a part.

For very young children, factors related to maternal health during pregnancy and the birth itself are important, with other aspects of health care having an impact throughout infancy. But for children over one year, the general socioeconomic environment matters more.

TABLE 4.1		Growth in GNP Per Capita Accounts for at Most One-Third of the Reduction in Mortality . . . and Less than a Fifth of Lower Fertility		
	1980 actual	2000 actual	2000 income-based estimate	Percent reduction explained by income
Under-five mortality	205.0	77.5	163.1	32.9
Total fertility rate	5.6	3.0	5.2	16.0
Stunting	67.6	44.7	62.3	23.1
Underweight	69.5	47.7	64.6	22.4

Source: Calculated from data used for figure 4.1 (see Annex B).

TABLE 4.2

Significant Determinants of Infant and Child Mortality

	Child characteristics	Household characteristics	Community characteristics
Neonatal	Public: Birth order, preceding birth interval. Private: Multiple birth, born in April/May (lower mortality).	Public: Electricity, water and sanitation, mother secondary education, maternal contraceptive knowledge, mother's age at birth. Private: Mother's mobility	Antenatal visits (public).
Postnatal	Public: Birth order, preceding birth interval. Private: Higher birth order female, born in October/November in rural areas (higher mortality).	Public: Mother's secondary education, maternal contraceptive knowledge, mother's age at birth. Private: Wealth.	Lower in Khulna, Rajshahi, and rural areas.
Child	Public: Immunization, vitamin A supplements. Private: Female, stronger for higher birth order and in Chittagong.	Public: Mother's primary and secondary education, mother's age at birth. Private: Wealth, mother remarried (higher mortality).	Lower in Khulna, Rajshahi.

Note: "Public" indicates a characteristic that is a function of publicly provided services; "Private" indicates otherwise.

Source: Annex C.

the different age groups as revealed by OED's analysis. The main findings are:

- Mother's education matters, though it must be to at least secondary level to affect infant mortality. Mother's contraceptive knowledge is included as a proxy for maternal health knowledge, and exerts a significant impact for infants but not children. Other measures of women's agency have some effect, notably the greater mortality risk for children of women who have remarried.
- Money matters, though not for neonatal mortality.
- Various factors correlated to fertility influence mortality: there is a nonlinear relationship between mother's age and infant mortality, with the probability of premature death decreasing until close to age 30, after which it increases again. For children, there is a simpler relationship: children born to older mothers have a lesser chance of dying. A short birth interval adversely affects mortality. The pattern for birth order is more complex: lower-order births, particularly the first-born child, are at greater risk, though this risk is mitigated slightly if the mother attends antenatal care. The data suggest that

the risk of mortality for infants declines with birth order until a birth order of six, after which it rises again.

- Sex bias manifests itself in various ways. Female children are more likely to die than male, especially those in Chittagong⁵ and those of a higher birth order.
- The health- and nutrition-related variables have the expected positive impact.⁶ Immunization is dealt with in detail below. Antenatal visits reduce neonatal mortality, but only if provided by trained medical personnel, either doctor or nurse. Training of both doctors and nurses has been a major area of DFID support over the last decade, and so a route through which they are contributing to reduced mortality.
- There are seasonal patterns in mortality, related to both mother's nutritional status during pregnancy and seasonal factors affecting illness.

The relative importance of these different factors can be explored in three ways. The first is to look at the extent to which they have contributed to mortality decline during the 1990s. To do so, the explanatory variable must both be significant and have changed notably during the

decade. The results of this analysis may be misleading, as there are some determinants, such as antenatal care or immunization, that have not changed much over the decade, but which would adversely affect mortality if not provided. A second approach, importance analysis, overcomes this difficulty, being based on the product of the coefficient and the standard error of the variable. Finally, the sources that explain the discrepancy between high and low mortality areas within the country are examined.

The first approach shows that improvements in economic well-being were the main driving force behind changes in child mortality in the 1990s,⁷ but this is closely followed by the role of health provision and maternal education (see Annex C). Expanded vaccination coverage alone accounts for over half of the mortality reduction achieved through higher wealth. But for infants the roles are reversed: health and education are more important than economic growth. The fertility-related variables have a mixed effect: survival chances are higher in smaller families with larger birth intervals, but the reduction in birth order tends to increase mortality.⁸ The importance analysis bears out these results, showing in particular the role of health and education. Two factors mainly explain why mortality outcomes are so much better in Khulna than Chittagong: (1) differing rates of accessing health services, and (2) pronounced excess female mortality in Chittagong, which is not present in Khulna.⁹

The above analysis of determinants is now used to quantify the impact and cost-effectiveness of the following interventions: immunization, training TBAs, female secondary schooling, and rural electrification.

Immunization

Immunization in Bangladesh has expanded greatly in the last two decades (see Chapter 2 and box 4.1). Improvements in both socioeconomic conditions and immunization reduce disease. A disease such as polio spreads more easily where access to safe water is limited, so that improving water supply can reduce incidence. But eradication of the disease requires a coordinated public health strategy, as has been implemented in Bangladesh with external assistance (see box 4.2).

More generally, there are various ways in which the deaths averted by immunization may be calculated (see Annex C for details on the calculations):

- The cross-country regression analysis undertaken for this study (Annex B) suggests that increased immunization coverage accounts for close to one-third of the reduction in under-five mortality over the past two decades. Immunization began to rise from the mid-1980s. If mortality had remained at its 1987 level, there would have been 3.28 million additional under-five deaths from 1987 to 2002—just under one million of these lives saved is attributable to immunization.
- A study conducted in the late 1990s considered the life-saving effects of each vaccination separately, producing an estimate of 1.15 million deaths prevented in the period 1987–98 (Khan and Yoder 1998).
- The child mortality regressions show that children with no vaccinations are between 50 percent to twice as likely to die as those children who have some coverage.¹⁰ Applying this ratio to observed mortality data suggests that more than two million child deaths were averted by immunization between 1981 and 2002 (see Annex C for details).

These varying approaches give somewhat different estimates of the number of lives saved by the Expanded Program of Immunization (EPI), but provide a range of one to two million deaths averted, which can be used as a basis for cost-effectiveness analysis. There are also wide estimates of the full cost of immunizing a child. A study on immunization in Bangladesh in the late 1990s reported a figure of close to \$12 per child, a little less than international norms.¹¹ The study also presents the direct costs

The first approach shows that improvements in economic well-being were the main driving force behind changes in child mortality in the 1990s, but this is closely followed by the role of health provision and maternal education.

But for infants the roles are reversed: health and education are more important than economic growth.

BOX 4.1**Which Children Get Immunized?**

Bangladesh's immunization program took off in the second half of the 1980s, rapidly expanding coverage. Despite this expansion, close to half of all children do not receive the full course of recommended vaccinations, and a significant number remain unvaccinated.

Multivariate analysis of DHS data shows the following:

- Several variables related to women's position (see Annex E for discussion of measures and correlates of women's agency) have a significant impact on the likelihood of a child being vaccinated: mother's age, mobility, agency, and education. Children of mothers who have remarried are less likely to be vaccinated.
- The divisional disadvantage of Dhaka, Sylhet, and Chittagong remains significant even once the other factors mentioned are controlled for. And the advantage of Khulna and Rajshahi is also significant. Distance from thana headquarters (HQ) reduces the likelihood of a child being immunized.
- Electricity has a significant positive impact. Two indirect channels for electrification—wealth and access to information (TV)—are already included, as is a locational variable (distance to thana HQ), which electrification may proxy for. Hence the result most likely reflects, at least in part, the impact of electrification on preserving the cold chain (keeping vaccines at the required temperature).
- Wealth has a positive but statistically insignificant impact on the probability of being immunized.

These results point to the importance of information campaigns in overcoming remaining barriers to immunization. As discussed in Chapter 5, the message at the heart of behavior-change communication—that all actors in the decisionmaking process need be reached—must be borne in mind.

The OED analysis thus confirms the relatively low cost of immunization as a means of saving lives.

There have been cases of substantial reductions in maternal mortality in systems relying on trained TBAs, when backed up by a good referral system for emergency obstetric care.

of the EPI program and estimates of total costs—that is, including staff and equipment costs not reflected in the EPI bud-

get—for a single year (\$18.3 million in 1997–98). Applying these figures gives a range of the cost per life saved from just under \$100 to just under \$300.¹² The OED analysis thus confirms the relatively low cost of immunization as a means of saving lives.¹³ These figures are likely to be an underestimate since they ignore the benefits

of herd immunity, which is yet to be realized in Bangladesh.¹⁴

The number of deaths averted can be attributed to the agencies supporting the immunization program on a pro rata basis. The World Bank

has contributed approximately 10 percent of program costs over the last 15 years, thus saving the lives of between 100,000 and 200,000 Bangladeshi children. DFID's contribution to immunization has been mainly to the polio program since eradication. Before the polio eradication campaign more than 10,000 children a year were developing the disease, which gives an indication of the benefits from keeping the disease from reappearing.

Trained TBAs

As described in Chapter 2, Bangladesh followed international trends in abandoning training of TBAs. But the international evidence is more nuanced than this wholesale shift in focus suggests. First, studies are clear that training TBAs *does* make a difference to their behavior. For example, a study in Bangladesh found that 45 percent of trained TBAs practiced the “three cleans” (hand washing with soap, clean cord care, clean surface) compared with 19 percent of untrained

BOX 4.2**Polio Eradication in Bangladesh**

Polio is a viral disease that affects the nervous system and muscles and can cause paralysis. Children under three are most at risk. The disease is transmitted by physical contact, and is highly contagious in areas with poor sanitation and high population density—making areas of Bangladesh very vulnerable. But polio is easily preventable using inexpensive vaccines.

In 1988 the World Health Assembly of the WHO set the goal of the global eradication of polio by year 2000. There has been considerable progress toward this goal. In 1988 there were 350,000 cases every year worldwide. In 2003 the number had dropped to 1,000. Only six countries still have endemic polio: Afghanistan, Egypt, India, Niger, Nigeria, and Pakistan.

Bangladesh has tackled polio through the four strategies identified by WHO: routine immunization as a part of EPI, national immunization days (NIDs), a surveillance system to identify outbreaks, and a mopping up strategy in case polio reappears in any location. This strategy in Bangladesh has been very successful. Before 1986 an estimated 11,500 children developed polio every year. This number was reduced to 2,300 in 1994, and 324 in 1999. Since 2000 no polio cases have been identified.

Funding efforts have been coordinated internationally by the Global Polio Eradication Initiative, which includes WHO, UNICEF, Rotary International, USAID, and other donors. In Bangladesh nearly 50 percent of the funding has been provided by the government of Japan, which spent \$40–50 million on NIDs up to 2003. Other large donors are the United States' CDC and Rotary (10 percent each); other minor donors follow: UNICEF (6 percent), WHO (3 percent), USAID (3 percent), DFID (2 percent), and so on. However, since eradication, DFID has expanded its support with a £7 million project, Support to Polio Eradication, under which coverage through NIDs is approaching 100 percent (97 percent in 2004).

*Sources: WHO/CDC/UNICEF (2004), Polio eradication Web site, <http://www.polioeradication.org/> and CDC, *Morbidity and Mortality Weekly Report* (various issues).*

TBAs (Goodburn and others 2000, p. 396; see Sibley and Sipe 2004 for results of a meta-analysis). Moreover, there have been cases of substantial reductions in maternal mortality in systems relying on trained TBAs, when backed up by a good referral system for emergency obstetric care (EOC) China is the best-known case;¹⁵ and Bangladesh has been cited as one in which there was no such system of EOC and, consequently, no impact on maternal deaths.¹⁶ Finally, while there is no evidence of an impact on maternal mortality, there *is* evidence of an impact on infant mortality. The meta-analysis of the impact of training TBAs reported that 14 out of 17 studies found a reduction of peri-neonatal mortality from births being attended by trained, rather than untrained, TBAs (Sibley and Sipe 2004).

Turning to the Bangladeshi data, training TBAs appears to have a beneficial impact on neonatal mortality, though there are complications in interpreting these figures. Table 4.3

shows the share of births attended by trained and untrained TBAs from each DHS, and the neonatal mortality rate for births so attended. The existence of training programs for TBAs is evident from the expansion of the share of births attended by trained TBAs (TTBAs) from 4 to 10 percent between 1993 and 1999. At the same time the neonatal mortality rate among births attended by trained TBAs has more than halved—

TABLE 4.3**Neonatal Mortality Has Fallen More Rapidly for Births Attended by Trained TBAs**

	Percent of deliveries		Neonatal mortality (per 1,000 live births)	
	Untrained	Trained	Untrained	Trained
1993	55.0	4.3	49	72
1996	56.7	7.6	48	49
1999	53.2	9.5	38	31

a much larger decline than that observed in general or among births attended by untrained TBAs.

But various factors complicate the picture.¹⁷ Trained TBAs are likely to attend more difficult births than untrained TBAs, as is evident from the higher mortality rate for these births in 1993.¹⁸ As training for TBAs expands, so that trained TBAs are more widespread, this effect may be diluted, thus driving down mortality associated with TBA-attended birth. Second, the training encourages TBAs to refer difficult births to EOC. Partly in consequence, the share of births attended by doctors also increased during this period, though this was mainly in urban areas (that is, only for a minority of births). This effect will also reduce the mortality associated with trained TBAs. Finally, training may reduce mortality owing to greater skills of TBAs who are trained, which is what we hope to observe, but which is difficult to disentangle from the other factors described above. However, it can be observed that neonatal mortality among babies whose birth was attended by a trained TBA was, in 1999, lower than that for untrained TBAs, despite the continued presence of the greater likelihood of difficult births for trained TBAs, which strongly suggests that training TBAs had a beneficial impact on neonatal mortality.

An estimate of the advantage of training is given by the single difference impact of 7 deaths averted per 1,000 live births.¹⁹ The coefficients from the multivariate analysis suggest a double

In a setting where 60 to 75 percent of births are attended by TBAs, it is not prudent to sideline them in any strategy for improving health outcomes. Doing so in Bangladesh was the result of faddism on the part of the international community, rather than a decision based on solid local evidence.

difference impact of 5 per 1,000. These figures are consistent with the meta-analysis mentioned earlier, which found that training TBAs reduced peri-neonatal mortality by 7 per 1,000 (Sibley and Sipe 2004). The cost of training a single TBA is in the range \$350–\$400, and one TBA attends between 10 and 15 births a year.²⁰ Hence, training one TBA saves 0.05–0.105 lives per year, suggesting a cost of between


US\$3,300–\$8,000 per life saved each year. The TBA may continue to practice using this knowledge for 10–15 years after training, bringing the cost per life saved down to US\$220–800. These figures show training TBAs to be a cost-effective strategy for tackling infant mortality.

In a setting where 60 to 75 percent of births are attended by TBAs, it is not prudent to sideline them in any strategy for improving health outcomes. Doing so in Bangladesh was the result of faddism on the part of the international community, rather than a decision based on solid local evidence. Such evidence suggests that training these women does change practices and that doing so makes a notable and cost-effective impact on infant mortality. However, there will not be an impact on maternal mortality unless EOC is also available. Hence, the stress needs to be on increasing availability of EOC services, and ensuring that training of TBAs results in speedy referrals where appropriate. Any renewed focus on training TBAs, which the above analysis suggests should certainly be considered, needs to pay attention to the need for a good supervision system for these workers and making periodic refresher courses available.

Female Secondary Schooling

Female secondary enrollments grew rapidly in the 1990s. DHS data show that in 1993, 20 percent of 17–24 year olds had secondary education, compared to 35 percent just 6 years later. Various factors lay behind this rapid increase. One of these is the stipend program, supported by the Asian Development Bank, NORAD, the World Bank, and the Government of Bangladesh. Introduced in 1994, this program pays a stipend to all girls in rural areas as long as they attend school on at least 75 percent of school days, maintain a passing grade, and remain unmarried (see Annex I for further discussion of the stipend program).

The beneficial impact of female education on infant and child mortality is shown above. Figure 4.2 confirms the bivariate relationship, with under-five mortality decreasing monotonically as education increases. Multivariate analysis generally supports this relationship, even though the effect may be underestimated, since the equation

FIGURE 4.2**Secondary Enrollments Have Risen Rapidly in the 1990s: Educational Attainment of Women 17 to 24 Years Old**

also includes some behavioral factors that may be affected by education (for example, number and timing of birth), as well as household income (proxied by the wealth index). The results are somewhat mixed for primary, showing no education effect for neonatal and a beneficial impact of primary on postnatal, but the regression for infant mortality suggests that lower secondary education compared with no education saves around 15 lives per 1,000 live births. For child deaths, both primary and secondary education are robust in reducing mortality. A child born to a mother with primary education is around 20 percent less likely to die than a child born to a mother with no education; for children of mothers with secondary education, this figure is 80 percent.²¹ Child mortality for children of women with no education stood at 42 per 1,000 by the end of the 1990s. Educating a mother to primary level would reduce the probability of premature death from 0.042 to 0.034; and educating her to secondary, to just 0.007, reductions of 8 and 35 per 1,000 respectively. The gain from secondary compared with primary is thus 27 child lives saved per 1,000 live births.

Some approximate figures illustrate the method used to calculate the cost-effectiveness of female secondary education in reducing under-

five mortality. The total fertility rate is around 3.3, so that 300 women will, over their lifetimes, give birth to 1,000 children. If these 300 women receive secondary education (rather than primary), then the number of child deaths averted is around 25 (using a conservative estimate based on child deaths only). The stipend cost per girl for the five years of secondary, including administrative costs, is US\$90. Therefore, the cost of getting the 300 girls to school is US\$27,000 ($300 \times \90). The cost per death averted is thus US\$1,080 ($=\$27,000/25$). This figure is an underestimate, since infant deaths have been excluded from the calculation, and various indirect behavioral channels are not considered. But it is also an overestimate for the reasons explained in the next paragraph.

The above calculation assumes that the presence of the program is the reason the girls go to school, though in fact some would have gone to school in the absence

For child deaths, both maternal primary and secondary education are robust in reducing mortality. A child born to a mother with primary education is around 20 percent less likely to die than a child born to a mother with no education, increasing to 80 percent for women with secondary education.

of the stipend. Determining the marginal impact of the stipend on girls' enrollment is not that straightforward. Best estimates suggest that of every 10 girls receiving the stipend, 2 of those would not have gone to school in the absence of the stipend (see Annex I). This raises the cost per death averted by a factor of five, to \$5,400.²² While, perhaps unsurprisingly, more expensive than the cost of direct health interventions discussed above, this figure is not high by international standards.

From 1993 to 1999 an additional 1.4 million girls attended secondary school, of which one million were from rural areas. Over this period,

Electrification can affect health outcomes through several channels:

(1) income effects;
(2) quality of health care, notably the cold chain;
(3) sanitary environment, including boiled water;
and (4) access to media, and thus to health messages.

the World Bank provided 0.8 million stipend years, which, assuming the stipend was paid to each girl for four years, means that 0.2 million additional girls were paid to go to school. These 0.2 million girls will go on to have 660,000 children. Since 25 deaths are averted per 1,000 live births, then 16,500 under-five deaths will be averted by this increased enrollment.

As above, it is assumed that the whole enrollment increase is on account of stipends. Adjusting for the likely marginal impact of the program on enrollments reduces the number of deaths averted by the World Bank's support of the stipend program to 3,300 (that is, just over 1 percent of the actual number of under-five deaths each year).

Rural Electrification and Access to Media

Electrification can affect health outcomes through several channels: (1) income effects; (2) quality of health care, notably the cold chain;²³ (3) sanitary environment, including boiled water; and (4) access to media, and thus to health messages. There is evidence of the importance of each of these effects in the results, partly evident in a direct effect from the electrification variable in multivariate analysis.

A study of the impact of rural electrification conducted for USAID found electrification to

cause an increase in income of 15 percent,²⁴ which translates into approximately an 11 percent increase in the wealth index used to measure economic well-being in DHS data.²⁵ Given the regression coefficients, electrification-induced income effects have reduced infant mortality by around 5 per 1,000, and for children at around 10 per 1,000. Assigning the whole "TV effect" adds another 8 per 1,000, and direct effects another 2 per 1,000. Electrification thus reduces the probability of under-five death by 0.025 (25 per thousand). During the 1990s a household connection cost US\$500,²⁶ so that connecting 1,000 households cost US\$0.5 million. Connecting these 1,000 households will avert 25 deaths, at a cost of US\$20,000 each.²⁷ As the program rolls out, connections become more expensive, currently reaching around US\$1,000 per household, bringing the cost per life saved to US\$40,000. Rural electrification is having a measurable impact on child health outcomes. Since there are many benefits to electrification other than reduced mortality, it is unsurprising that the cost per life saved is high relative to that of other interventions.

Access to media variables—particularly TV and radio—are significant in the outcome regressions for mortality (as well as those for fertility and nutrition). These results provide evidence of the importance and effectiveness of information, education, and communication (IEC) campaigns. Given the diverse nature of these campaigns (see Chapter 2 on the different ministries involved), and different channels through which households access information, it is not possible to quantify their impact and cost-effectiveness.

Fertility Reduction

The cross-country evidence presented at the beginning of this chapter feeds directly into a debate as to the causes of fertility decline in Bangladesh. Some have claimed that the usual socioeconomic factors explaining lower fertility have been absent and that the country's internationally renowned family planning program explains the success in fertility reduction. Others argue that there has been progress—notably, re-

duced poverty and a measure of women's empowerment—that can explain lower fertility. The evidence presented above shows that both explanations have played a part. Socioeconomic changes, including the usual demographic transition effect of mortality reductions driving down fertility, do explain some of Bangladesh's fertility reduction in recent decades—but not all of it. There is a substantial residual, which is most plausibly explained by the presence of a successful family planning program.

Why can the unexplained fertility reduction most plausibly be attributed to family planning? Evidence to support this argument comes from a number of ways in which Bangladesh is exceptional. First, figure 4.1(b) already showed the large reduction in fertility to be greatly in excess of what would have been expected from the country's income growth over this period. Second, figure 3.4 showed that contraceptive knowledge was almost universal in Bangladesh by the early 1980s. Bangladesh is an outlier in this respect, meaning that contraceptive knowledge is much higher than expected for a country of Bangladesh's income and female education levels. Furthermore, as discussed below, Bangladesh is also an exception in the low level of its fertility given the low age at which women get married. These discrepancies are best explained by the presence of the family planning program. Finally, multivariate analysis confirms the links between family planning acceptance and fertility in Bangladesh (Annex F). Less directly, the analysis shows the importance of access to media in lowering family size, indicating the contribution of IEC campaigns to spreading awareness of small family size and contraceptive knowledge.

The argument as to the impact of the family planning program in the past has significance to the present day, since the claimed fertility plateau in the 1990s has cast new doubt on the program's effectiveness. However, it was shown in the last chapter that fertility decline did indeed continue throughout the 1990s, with contraceptive usage continuing to grow. There are valid concerns among policymakers about contraceptive discontinuation, but it would be a mistake to conclude that the program has stalled.

Nonetheless, fertility decline has slowed, so it is legitimate to ask how momentum may be maintained. The government's new HNP Strategic Investment Plan highlights the role of increasing the age at marriage as a means of reducing fertility, and several programs, including the counseling provided under the World Bank–

supported Bangladesh Integrated Nutrition Project (BINP), promote getting married later. It is a condition of the female secondary school stipend program that recipients remain unmarried. It is true that the age at marriage in Bangladesh is low, with half of all women marrying by age 14. It is also true that there is a well-established international pattern whereby increasing the age at marriage drives down fertility. But this pattern should not be expected to be observed in Bangladesh for two reasons: (1) raising the age at marriage of girls aged 13 or less has no effect on the age at which they have their first child (as the age at marriage has risen, the gap from marriage to first birth has fallen: see Annex F); and (2) if a woman plans to have only three to four children, as the majority of Bangladeshi women do, then this can be accomplished whether childbearing begins at 15 or 20. The effect of expanding education is muted since Bangladesh has already attained levels of fertility comparable to those in countries with higher levels of education. Hence, raising the age at marriage, while desirable for both maternal and child health (children born to young mothers have a greater chance of premature death),²⁸ will have little impact on fertility. At the same time, there remains a substantial proportion of high-fertility households. Reducing fertility among these would make an important contribution to continued fertility decline. One means of doing this would be to attempt to restore the use of permanent contraception to its previous level—

Socioeconomic changes do explain some of Bangladesh's fertility reduction in recent decades—but not all of it. There is a substantial residual, which is most plausibly explained by the presence of a successful family planning program.

Fertility decline has slowed, so it is legitimate to ask how momentum may be maintained.

Higher-fertility women are less likely to be utilizing health services, so that the case for home visits should be reconsidered as a part of the wider strategy for fertility reduction.

long-term methods accounted for 30 percent of contraceptive use in 1991, but only 12 percent by 2004.²⁹ Higher-fertility women are less likely to be utilizing health services, so that the case for home visits

should be reconsidered as a part of the wider strategy for fertility reduction. In addition, though it is difficult to suggest appropriate policy responses, attempts can be made to tackle son preference, which creates a barrier to fertility decline.

Nutrition

In recent years the main focus for improving nutritional outcomes has been through BINP, which is the subject of the next chapter. However, BINP has had a limited geographical focus, while, as shown in Chapter 3, nutritional status has been improving across the country. DHS collected anthropometric data in 1996 and 1999, which thus allows some analysis of the factors behind this improvement. OED analysis shows that:

- Wealth is a significant determinant of a child's nutritional status, and the increase in the wealth index between the two surveys ac-

counts for close to 10 percent of the observed reduction in stunting.

- Mother's education to secondary level improves child nutrition, whereas children of mothers with primary education alone have no nutritional advantage over those whose mothers have no education. The expansion of secondary education accounts for about 6 percent of the observed fall in stunting. These nutritional gains can be added to the beneficial impact of the Female Secondary School Stipend Program (FSSAP).
- A household having electricity reduces the probability of stunting; service expansion accounts for about 4 percent of the decline in stunting in the late 1990s.
- Children of lower birth order have better nutritional status, capturing the advantage of smaller family size in limiting resource competition among young children. The reduction in mean birth order, as a result of falling fertility, accounted for about 6 percent of the improvement in nutritional status.
- The regression results leave a fair amount of the reduction in stunting unexplained. Possible reasons for this decline are picked up in the next chapter, which discusses the Bangladesh Integrated Nutrition Project in greater depth.



A Closer Look at Nutrition: The Bangladesh Integrated Nutrition Project

The Community-Based Nutrition Component of BINP was to improve nutritional status through nutritional counseling and supplementary feeding for malnourished children and pregnant women. Both counseling and feeding suffered from problems of inappropriate targeting strategies and a failure to reach intended groups. While counseling has changed women's knowledge, it has had less of an impact on behavior. There is a knowledge-practice gap, which is explained by resource constraints faced by women, including lack of time, that prevent them from putting advice into practice. The impact of the project on anthropometric outcomes has not been large, so that the approach does not appear to be a cost-effective means of tackling poor nutrition.

While rapid strides were being made in reducing mortality and fertility in the 1980s, malnutrition showed no improvement; it affects close to 70 percent of all children under five. In order to address this remaining problem, the government undertook a pilot nutrition intervention supported by the World Bank, the Bangladesh Integrated Nutrition Project (BINP), which was to initiate a national program whose ultimate goal was to reduce malnutrition to the extent that it ceased to be a public health problem. More specific targets—such as the reduction of severe malnutrition by 40 percent, and reduction of moderate malnutrition by 25 percent in the project areas—were also set. The project is now being scaled up to the national

level. This chapter takes a look at the evidence regarding the effectiveness of the BINP's main component, the Community-Based Nutrition Component (CBNC).

BINP, and notably the CBNC, has been the subject of several studies: the independent evaluation of BINP for the project (Karim and others 2003), a study by Save the Children (2003), an evaluation commissioned by the government's evaluation department (Implementation, Monitoring and Evaluation Department, IMED); Haider and others

While rapid strides were being made in reducing mortality and fertility in the 1980s, malnutrition showed no improvement; it affects close to 70 percent of all children under five.

The key assumption behind the Community-Based Nutrition Component is that “bad practices” are responsible for malnutrition in Bangladesh.

mentioned reports through the application of a theory-based evaluation framework, and it improves on the quality of the control for the BINP evaluation data set by using a nationally representative sample survey to construct a new control using propensity score matching.²

Overview of the Project

BINP had three components: (1) national nutrition activities (US\$20.6 million), including institutional development, IEC, and monitoring and evaluation (M&E); (2) community-based nutrition (US\$39.1 million); and (3) intersectoral nutrition program development (US\$7.6 million), supporting schemes such as home gardening and poultry rearing.

CBNC was the main component, the one addressed here. In each project thana a number of community nutrition promoters (CNPs) were recruited—local women with children of their own who had achieved at least an eighth-grade education. The CNPs implemented activities at the community level: monthly growth monitoring for children under 24 months old, supplementary feeding for malnourished children and pregnant women, and nutritional counseling in a variety of settings. The CNPs were overseen by a community nutrition officer and supervisors at the thana level who were staff of the implementing nongovernmental organization (NGO).³

The Theory Underlying the CBNC

The key assumption behind CBNC is that “bad practices” are responsible for malnutrition in Bangladesh. This point of view was strongly argued in the BINP appraisal document: “behaviors related to feeding of young children have at least as much (if not more) to do with the serious problem of malnutrition in Bangladesh as poverty and the resultant household food insecurity

2004) and two PhD theses from the University of Cambridge. For this study, OED has obtained the BINP evaluation and Save the Children data sets.¹ The analysis is extended from that contained in the afore-

do” (BINP SAR, para. 1.13, p. 4; World Bank 1995). Therefore, changing bad practice to good will bring about nutritional improvements. There are a number of steps in the causal chain behind this approach:

- The right people (those making decisions regarding undernourished children) are targeted with nutritional messages.
- These people participate in project activities, and so are exposed to these messages.
- Exposure leads to acquisition of the desired knowledge.
- Acquisition of the knowledge leads to its adoption (that is, a change in practice).
- The new practices make a substantial impact on nutritional outcomes.

A feeding program for malnourished children and pregnant women was implemented alongside growth monitoring. For this program to work:

- The target groups have to enroll in the program.
- The criteria are correctly applied in selecting those to receive supplementary feeding.
- Those selected for supplementary feeding attend sessions to receive the food.
- There is no leakage (for example, selling of food supplements) or substitution (reducing other food intake).
- The food is of sufficient quantity and quality to have a noticeable impact on nutritional status.

If project design fails to take account of one of these steps, there is a missing link in the causal chain. If activities take place corresponding to a step but are ineffective, then there is a weak link.

Project Coverage and Targeting

BINP nutritional counseling activities target pregnant and lactating women and adolescent girls.⁴ Clearly strategies to affect health behavior need to influence the attitudes of all those involved in making health decisions. In Bangladesh, decisions regarding health and nutrition do not rest solely with the mother, but also the husband, and frequently the mother-in-law. Data from the Demographic and Health Survey show

that only one in five women is solely responsible for decisions regarding their own health and that of their children, falling to only one in 10 of women living with their mother-in-law (see Annex E). While women are most likely to have responsibility over deciding what to cook (two-thirds of women are solely responsible for this decision, though this is so for only 42 percent of women living with their mother-in-law), which clearly matters for child nutrition, the effectiveness of this decisionmaking power is constrained by the fact that it is men who do the shopping in Bangladesh. This is reflected by the fact that only 17 percent of married women in male-headed households are responsible for decisions about daily purchases, and only 10 percent of those women living with their mother-in-law have this responsibility. Hence, there appears to be a need for the project to broaden the target audience for its nutritional messages.⁵

Similarly, just over half (52 percent) of married women can visit the health center either alone or with their children. But this percentage falls to 39 percent for women living with their mother-in-law (Annex E), showing that this domestic arrangement is an important factor in restricting women's mobility.

Ideally all children in the project area should participate in growth monitoring, with a target set in the project appraisal document that 80 percent of 0-to-24-month-olds should be registered, and 80 percent of these (64 percent of all children) receive at least 18 out of 24 monthly weighings. The data show that over 90 percent of children were weighed at least sometimes, with 88 percent being weighed on a regular basis.⁶ Hence the project's coverage targets were met, although these allowed for some children to go unmonitored. If the children not attending have little risk of malnutrition, then their omission can be seen as an efficiency gain for the project rather than a shortcoming. Multivariate analysis shows no income bias in participation in growth monitoring, though the most educated mothers, who presumably need it least, were less likely to attend (figure 5.1 and Annex G). Fieldwork suggested that factors constraining participation include remoteness and

traditional constraints on women's mobility. The multivariate analysis shows that these factors do indeed play a role, with constraints on mobility, particularly for women living with their mothers-in-law, being especially important in the two more conservative thanas included in the survey (Rajnagar and Shahrasti).

Low growth, identified by growth monitoring, was to be addressed in two ways: nutritional counseling and supplementary feeding. The feeding is said to have been intended as an example to mothers, the heart of the strategy being counseling to achieve behavior change.⁷ The growth monitoring sessions themselves are too chaotic a setting to provide nutritional counseling. But the CNPs work full time in their position and provide advice through different forums, such as the various group counseling sessions or one-to-one meetings with parents. However, over one-third of women whose children were receiving supplementary feeding said that they had neither discussed nutrition directly with the CNP, nor sat in any meeting where it was being discussed.


Supplementary feeding was provided to children who were detected to be severely malnourished (less than -4 standard deviations from the reference median weight for age) or experiencing growth faltering⁸ for a period of three months, extended for a further month if they did not show the required weight gain. Analysis of register and field data from the Cambridge studies showed a reasonably low Type II error:⁹ only 16 percent of children receiving food supplementation should not have been doing so.¹⁰ How-

Strategies to affect health behavior need to influence the attitudes of all those involved in making health decisions. In Bangladesh, decisions regarding health and nutrition do not rest solely with the mother, but also the husband, and frequently the mother-in-law.

Just over half of married women can visit the health center either alone or with their children. But this percentage falls to 39 percent for women living with their mother-in-law, showing that this domestic arrangement is an important factor in restricting women's mobility.

FIGURE 5.1

Various Factors Affect Women's Participation, but Restrictions on Women's Mobility in More Conservative Areas Are the Most Important



ever, of those enrolled, only one-quarter (26 percent) received supplementation for the recommended three months, the majority dropping out sooner. And Type I error was very high: over two-thirds (69.8 percent) of eligible children were not being fed.¹¹

While an acceptably low percentage of ineligible children were being fed, the criteria themselves are open to question. Growth faltering is quite normal, so enrolling growth faltering children, regardless of their nutritional level, will mean enrolling perfectly well-nourished children: one study found that 37 percent of a group of U.S. children from well-off backgrounds would qualify for supplementary feeding under the criteria used in the Tamil Nadu Integrated

Nutrition Project in India (Martorell and Shekar 1992). The Save the Children register data show that over 40 percent of those enrolled in the feeding program were

not malnourished (that is, the children were above -2 SDs WAZ).

Turning to monitoring of pregnancy weight gain, close to three-quarters of pregnant women attended weighing sessions, and just under half received supplementary feeding, with both these percentages being a bit lower at endline than the baseline. There is no pattern between attendance at weighing sessions and the mother's nutritional status, which is to be expected. Supplementary food was meant to be received by women not having normal body mass index. However, by the endline, about 60 percent of eligible women were not receiving the supplement. At the same time, 40 percent of those who were receiving the supplement were not eligible.¹²

For both children and mothers, there is evidence of both leakage and substitution of the food supplement. For example, 32 percent of women said they had shared their food supplement with someone else. Many of those who were not sharing said they did not eat more during pregnancy than usual, indicating that the

For both children and mothers, there is evidence of both leakage and substitution of the food supplement.

BINP-provided food was substituting for other foodstuffs. This was possible, since many women and children, contrary to project design, consumed the food at home.

In summary, enrollment in growth-monitoring sessions has been at a reasonable level for both children and pregnant women. However, attendance at these sessions has not provided opportunities for nutritional counseling for a sizeable minority of women. There have been problems in the targeting of feeding programs, especially the exclusion of eligible participants among both children and pregnant women. In the case of pregnant women, a considerable number of feeding beneficiaries are in fact ineligible. Such Type II error is less of a problem for child feeding, though the entry criteria themselves appear inappropriate, and only a minority of enrolled children complete the full three months of feeding.

Acquiring Knowledge

The central thrust of the project design was to change nutritional behavior of child caretakers. There are a number of nutritional practices considered adverse to child nutrition. Some are simple differences in habit, such as cutting vegetables before washing them rather than vice versa, which is nutritionally disadvantageous. Others, such as eating less during pregnancy (“eating down”), result from different perceptions of health risks and benefits (mothers perceive the benefit of a lower-risk delivery of a smaller child, discounting the risks to low-birth-weight children).¹³ And others stem from traditional beliefs that appear to have no plausible health-related rationale, such as avoiding meat, fish, and eggs during pregnancy.

Data from both Save the Children and the BINP evaluation show that knowledge of good nutritional practices is indeed higher in project areas than in the control. For example, 63 percent of women in project areas say that the baby should be fed the colostrums (the first milk), compared with only 52 percent in the control area, a difference of 11 percent. The BINP dataset allows examination of the double difference for two variables, for both of which there has been an in-

crease in knowledge in the project area but not in the control. At the baseline, a slightly higher percentage of women in the control thought it advisable to eat more during pregnancy than was the case in the project area. This proportion had not changed in the control areas, whereas in the project area it rose by 28 percent (which is therefore the double-difference effect, since there was no change in the control).

Of course, other factors may be affecting knowledge, such as superior education in the project area. Multivariate regression analysis was used to control for these factors (Annex G). The probability of a mother knowing a specific practice was modeled as a function of mother’s age, education, education of the household head, whether the woman was a group member, and participation in project activities. In the Save the Children data, variables were included in the analysis of whether a woman simply lived in the project area, or whether she also participated in nutrition education activities. The BINP evaluation dataset allowed the inclusion of additional measures of socioeconomic status; however, the only available measure of participation in project activities related to supplementary feeding and attending an antenatal clinic.¹⁴

In summary, knowledge of the prescribed nutritional practices is higher in the project areas than the control, the difference being greatest for women who have participated in project activities. But this knowledge is not universal, which is consistent with the evidence of weak links in the causal chain through incomplete enrollment, partial mistargeting of supplementary feeding (and so the associated counseling), and failure to transmit counseling messages to all women whose children have nutritional problems.

There have been problems in the targeting of feeding programs, especially the exclusion of eligible participants among both children and pregnant women.

Data from both Save the Children and the BINP evaluation show that knowledge of good nutritional practices is indeed higher in project areas than in the control.

Turning Knowledge into Practice: The Knowledge-Practice Gap

Although the project has had success in promoting nutritional information, a considerable knowledge-practice gap remains: that is, women do not put into practice the “good behaviors” (figure 5.3). This gap exists for every practice, and is extremely large in the case of exclusive breastfeeding. The gap exists in both project and control areas, with little evidence that the gap is any less in project areas than control.

Although the project has had success in promoting nutritional information, a considerable knowledge-practice gap remains: that is, women do not put into practice the “good behaviors.”

The multivariate analysis shows that attending nutritional counseling indeed has a significant impact on a woman’s nutritional knowledge, though being in receipt of supplementary feeding does not.¹⁵ However, even when these participation variables are in-

cluded, the BINP project dummy is still significant. This means either that there are spillover effects (women who get the knowledge in nutrition sessions communicate it to others)

or that other project activities not captured in the participation variables, for example, women’s group meetings, are also channels for communication of nutrition education. According to these regression results, simply living in the project area raises a women’s probability of having a piece of nutrition knowledge by 7 percent, but full participation in project activities increases this probability to between 10 and 23 percent (figure 5.2). That is, the proportion of women aware of the importance of colostrums feeding is 23 percent greater for women participating in project activities than for women in the control area. Other determinants of a women’s nutritional knowledge are found not to vary much between the project and control, and so account for only a small amount of the difference in knowledge between the two areas (Annex G).

Multivariate regression analysis (Annex G) and the results of qualitative fieldwork commissioned by OED identify a common set of factors that explain the knowledge-practice gap. Resource and time constraints are foremost among these. Women who have work to do, including caring for children and elderly relatives, are less

FIGURE 5.2

Women Living in Project Areas Are More Likely to Have Nutritional Information, Especially if They Participate in Project Activities





FIGURE 5.3
The Knowledge-Practice Gap in Project Areas: More Women Say They Know Good Behavior than Actually Practice It


likely to be able to rest or avoid hard work during pregnancy (table 5.1). Women engaged in agricultural work may also not have time to breastfeed, or not be able to do so if they are with the child away from the home. Although the project has some effect in reducing the gap, the multivariate analysis shows the magnitude of this effect to be very small. There are other channels—such as increasing education

of both males and females—that will also close the gap.

In summary, the project does increase knowledge about nutritional practices. However, there is a gap between knowledge and practice, and the project does not have an impact in reducing the size of the gap. But since knowledge is more widespread, and the gap is the same in project and control areas, then the promoted practices are more

TABLE 5.1
Many Factors Prevent Women from Putting Nutritional Advice into Practice, Though the Project Partially Overcomes Some of These

	Main determinants	Project effect	Moderating project impacts
Rest during pregnancy ^a	Agricultural work, children, elderly male in household, poverty.	None	None
Colostrum	Children reduce gap.	None	None
Breastfeeding	Agricultural work.	None	Reduces effect of living with mother-in-law and being poor.
More food during pregnancy	Poverty, children; having a vegetable garden reduces the gap.	Reduces gap	Bigger effect in working season.
Avoid hard work during pregnancy	Children, agricultural work (including vegetable garden), but lower for female-headed households and women not in paid employment.	Reduces gap	Reduction of gap larger for poor, but smaller reduction during working season.

a. The same results are found in both the Save the Children and the BINP data.

Source: Annex G.

widespread in the project area. So, are these practices beneficial to nutritional outcomes?

The Nutritional Impact of BINP Interventions

Child Nutrition

Existing studies have found somewhat different results with respect to the impact of BINP, though where an impact has been found its magnitude is not that great. This study used the nationally representative Nutritional Surveillance Project (NSP) data to construct a control group using propensity score matching. These results find a significant project impact from BINP at midline on all three z-scores of just over 0.1. However, at endline the impacts are smaller and most are not significant. This figure means that the project reduced malnutrition in project areas by at most 5 percent—well short of the project target, so the project failed to achieve its central objective.

It is difficult to construct a suitable control for supplementary feeding. However, the best estimate possible suggests that a child receiving

feeding has a weight gain equal to 0.1 WAZ for those under 12 months old, and over 0.4 for those over one year of age. The NSP data show that WAZ falls by about 0.5 WAZ over a three-month period for children under 12 months, and then stabilizes: hence, supplementary feeding has a positive impact for children of this age of around 0.6 WAZ. The impact is greater still for children who entered the program more malnourished.

Low Birth Weight

BINP aimed to increase pregnancy weight gain both by encouraging women to eat more during pregnancy and by providing supplementary feeding to malnourished pregnant women. While there was a striking improvement in pregnancy weight gain in project areas, there was an even greater improvement in the control, suggesting that non-project factors played a large role in the observed improvements. Nonetheless, multivariate analysis suggests that there was some project effect, of about 200 grams per month, mainly through the likelihood that

BOX 5.1

Qualitative Perspectives of the Knowledge-Practice Gap: The PPS-BD Study

OED commissioned the Participatory Practitioner's Society–Bangladesh (PPS-BD) to conduct a qualitative study to examine the existence of traditional beliefs related to child and maternal nutrition and to detect the reasons for mothers' inability to transform health knowledge into practice. The research was carried out in two communities falling under BINP and two in non-BINP villages.

In addition to focus groups and key informant interviews, semi-structured interviews were conducted with 150 individuals. These interviews began with a structured questionnaire asking similar questions to those used in the BINP evaluation to identify the presence of a knowledge-practice gap for both health and nutritional practices. Where a discrepancy between knowledge and practice was identified, unstructured questioning was used to probe the reasons for the gap.

In general the study found that harmful health beliefs, such as eating less and working more during pregnancy, were on the decline, and were more prevalent among older women. In project areas, the reduction was mainly attributed to BINP initiatives; in the control areas, the diffusion of mass-media, national campaigns and health workers were considered the main reasons for the change. Regarding the knowledge-practice gap, women pointed to the lack of resources as the reason for their inability to eat more during pregnancy. Women also mentioned excessive workload and family opposition to the need for taking rest when pregnant. Cost, distance, and dissatisfaction with the services offered were the reasons women stated for not seeking antenatal care.

In addition, a short questionnaire was administered to a sample of 10 CNPs in each project thana in order to assess the quality of counseling and to test the CNPs' ability to interpret the growth charts used in the child-growth-monitoring sessions. These interviews revealed a modest understanding of the charts, and great variation as to the counseling messages given in each case.

women in project areas would eat more during pregnancy than those in the control.¹⁶ The project-related weight gain is thus not much more than 1 kilogram over six months, which, evidence suggests, increases birth weight by only 20 grams. That is, the amount of pregnancy weight gain appears insufficient to have a substantial impact on low birth weight.

This conclusion is supported by direct estimates of project impact on low birth rate, although these suggest a slightly larger impact of about 80 grams, again mainly as a result of the mother not eating down. However, there is a much stronger impact of around 270 grams for children born to women with poor nutritional status, which is reassuring, as women of low body mass index (BMI) need a larger pregnancy weight gain to avoid low birth weight than do women with a higher than average BMI. The project also seems to smooth out the seasonal effects that come from mothers being less well nourished in the lean season, which is most likely a result of supplementary feeding.

Cost-Effectiveness Analysis

The impact results can be used to calculate the cost to move a child out of malnutrition, the results of which are shown in table 5.2. The first two rows of the table show the average amount of funds to be invested on a child over a year in order to move the child out of malnutrition, measured by WAZ and HAZ. Based on estimates of the relationship between child malnutrition and mortality, the last row shows the amount of money to be spent in order to save one life. For the BINP project, these figures were obtained using data on project costs, population of project areas, and project effects on malnutrition rates at the midterm and the endline (see Annex G). The rice ration simulation uses the relationship between food intake and physical growth observed in other studies, and simulates the nutritional effects of investing BINP funds on the same areas and population covered by the BINP project in 1998. The unit cost of saving one life and a malnourished child is less under the rice ration simulation, because, given the assumptions used, this intervention would produce a larger reduction in


TABLE 5.2		Cost of Nutrition Improvements and Mortality Reduction (US\$)	
	BINP		Rice ration
	Midterm (1998)	Endline (2003)	
Weight-for-age	187	333	110
Height-for-age	241	490	NA
Per life saved	2,328	4,095	2,223

malnutrition. The cost (using CBNC costs associated with children—that is, excluding those for pregnant women) per life saved from improved nutritional status is also shown. For comparative purposes, the results of a simulation assuming that the project funds were instead used to purchase rice that was given to families with children is also shown;¹⁷ this proves to be a more cost-effective route to improving child nutrition.

Other Sources of Improved Nutritional Outcomes

Both the OED results and the BINP evaluation find a stronger impact on child nutrition at midterm than endline. Between midterm and endline, pregnancy weight gain was greater in control thanas than those falling under BINP. The impact of BINP appears to have waned over time. One reason for this may be the difficulty of maintaining good implementation over time, or as the program is rolled out. Unfortunately, the evaluation data set has few process variables to explore this hypothesis. A second explanation is that there were external factors that allowed non-project areas to catch up. As already seen in Chapter 4, there was already an improvement in anthropometric outcomes in the late 1990s not explained by

Both the OED results and the BINP evaluation find a stronger impact on child nutrition at midterm than endline. One reason for this may be the difficulty of maintaining good implementation over time, or as the program is rolled out. A second explanation is that there were external factors that allowed non-project areas to catch up.

FIGURE 5.4**Rice Production and Daily Energy Supply Grew Rapidly in the Late 1990s**

Greater food availability, combined with rising incomes and lower prices, presents a very plausible explanation for improved nutritional status across the country.

the determinants of wealth, maternal education, and other factors.

There is a clear candidate for such an external factor: daily energy supply grew rapidly in Bangladesh in the second part of the 1990s as a result of a yield-driven increase in

rice production (figure 5.4 and Annex J). Famine theory shows the importance of entitlements: it is not simply food availability that matters, but whether people can afford to buy that food. In Bangladesh real incomes have been growing throughout the 1990s, at over 5 percent a year, picking up in the second half of the 1990s. Although growth in Bangladesh has not been pro-poor, real income growth has been above 2 percent for the lowest income groups (World Bank 2004). Meanwhile the rice price has been stable in nominal terms, representing a real fall in the average rice price of 14 percent from the

early to late 1990s, and a further 15 percent from 1999 to 2002 (Annex J).

Greater food availability, combined with rising incomes and lower prices, presents a very plausible explanation for improved nutritional status across the country. It is possible that at the time of the midterm, these factors had not fully passed through to improved nutritional status, allowing some effect of BINP to emerge. But by the endline, project impacts had been dampened by improved energy intake in all areas, since the project is most effective for severely malnourished groups.

Testing the Theory in Practice—How Well Did the Causal Chain Operate?

BINP succeeded in achieving high levels of participation in its activities in project areas, enrolled large numbers in supplementary feeding, brought about significant increases in nutritional knowledge, and, to a lesser extent, effected changes in practice. However, nutritional outcomes in terms of low birth weight have been disappointing.

While child nutrition, especially for children participating in supplementary feeding, appears to have been better, the overall difference in performance compared to the control is not great.

The longer the causal chain, the more likely it is that final outcomes will not be realized on account of missing or weak links in the chain, since there are more opportunities for external factors to undermine the logical flow from inputs to outcomes. There were two missing links in the BINP chain: the first was the relative neglect of some key decisionmakers regarding nutritional choices (men and mothers-in-law), and the second was the focus on pregnancy weight gain rather than prepregnancy nutritional status. Participation levels of the target audience were high, but many women escaped exposure to nutritional messages, and there was a high Type I error in the

feeding programs (see table 5.3). A substantial knowledge-practice gap persisted, so many women did not put the advice they received into practice, especially if they were resource or time constrained. Those receiving supplementary feeding often shared it with others or substituted it for their regular foodstuffs. This list of weak links in the chain explains why project impact was muted by the time final outcomes were considered. While attention

BINP succeeded in achieving high levels of participation in its activities in project areas, enrolled large numbers in supplementary feeding, brought about significant increases in nutritional knowledge, and, to a lesser extent, effected changes in practice. However, nutritional outcomes have been disappointing.

TABLE 5.3**Links in the Causal Chain**

Assumption	Children	Mothers
Attend growth-monitoring sessions.	Over 90 percent of all children attend growth-monitoring sessions.	Over 70 percent participate in monitoring pregnancy weight gain.
Targeting criteria is correctly applied; participants stay in the program to receive food.	Nearly two-thirds of eligible children are not fed (reasons: don't attend growth monitoring in the first place, wrong application of targeting criteria, drop out of feeding).	60 percent of eligible women are not receiving supplementary feeding.
Acquire knowledge and put it into practice.	One-third of mothers of children receiving supplementary feeding do not receive nutritional counseling. There is a knowledge-practice gap (see mothers).	There is a knowledge-practice gap driven by material resource or time constraints.
No leakage or substitution.	One-quarter of children are fed at home, increasing possibility of both leakage and substitution.	One-third admit sharing food, and there is substitution for those who do not. At most, 40 percent of eligible women receive full supplementation.
Feeding and nutritional advice have an impact on nutritional status.	Supplementary feeding has a positive impact on child nutritional status, especially for the most malnourished children. There is only weak evidence of any impact from nutritional counseling.	Pregnancy weight gain is too little to have a notable impact on low birth weight, except for the most malnourished mothers. Moreover, mother's pre-pregnancy nutritional status is more important than pregnancy weight gain. Consequently, birth weight gains are slight, though they are greater for children of severely malnourished mothers. Eating more during pregnancy is the main channel for both pregnancy weight gain and higher birth weight.

can be paid to each of these weak links, the BINP experience does demonstrate the difficulty of implementing complex designs.

Given that the nutritional gains from the project are small, it does not appear to be a particularly cost-effective intervention. The cost for

reducing malnutrition rates appears high, apparently more costly than simply buying food and giving it away, even allowing for a 25 percent administrative cost to do so. The efficacy and efficiency of the program must be improved in order to justify its continued existence.



Lessons Learned

Externally supported interventions have had a notable impact on MCH-related outcomes in Bangladesh. Immunization has proved particularly cost-effective, and has saved the lives of up to two million children under the age of five.

Immunization has prevented up to two million children from dying prematurely since the launch of the Expanded Program of Immunization, at a cost of less than US\$200 per death averted. Among the benefits that may be attributed to World Bank support are 200,000 lives saved through immunization and 3,300 saved through support for the Female Secondary School Assistance Project.

World Bank support to sectors outside of health has contributed to better child health outcomes.

Interventions in different sectors affect maternal and child health. The World Bank has supported both female secondary schooling and rural electrification, both of which have reduced under-five mortality.

Small amounts of money save lives . . . though the amount varies significantly by intervention.

Immunization is the most cost-effective means of saving lives. But even interventions

outside the health sector can avert deaths at a relatively low cost (table 6.1). These reductions in mortality are on top of the other benefits from these interventions, such as higher income from rural electrification and lower fertility from female education.

Although interventions from many sectors affect MCH outcomes, this fact need not imply that multisectoral interventions are always needed.

This study has shown that interventions in one sector affect outcomes in other sectors. But they were not multisectoral interventions. The female stipend program is implemented under the auspices of the Ministry of Education, and electrification by rural electrification boards. There was no need for the Ministry of Health and Family Welfare to implement its own education stipend programs or construct electricity generators—nor even for them to be consulted on the implementation of these programs by the responsible line ministries—for these intersectoral impacts to be felt. It is important that the presence of intersectoral impacts not be confused with the need for multisectoral interventions or overcentralization of sectoral activities.

TABLE 6.1**Cost-Effectiveness of Interventions in Reducing Under-Five Mortality (US\$ per death averted)**

Intervention	Lower estimate	Upper estimate
Immunization	100	300
Training TBAs	220	800
Female Secondary School Stipend Program	1,080	5,400
BINP	2,300	4,100
Rural electrification	20,000	40,000

Source: OED analysis.

World Bank support for training traditional birth attendants has reduced neonatal mortality...but this program has now been abandoned following the international trend toward support for skilled birth attendants.

Analysis of DHS data shows that TBA training in the 1990s reduced infant mortality. However, support to training TBAs stopped as the international Safe Motherhood Initiative chose to emphasize instead the role of skilled birth attendants. While training TBAs has had limited impact on maternal mortality in Bangladesh, as a result of weak emergency obstetric care, TBAs continue to attend over half of all births. Given this situation, one that is likely to take time to change, it is unwise to make such a wholesale policy turnaround. In Bangladesh the Bank followed this trend and dropped the training of TBAs from its projects, even though there was no local evidence to support such a decision.

Programs should be based on local evidence, rather than general conventional wisdom.

The abandonment of training TBAs is an example of the general finding of the importance of taking context into account in program design.

A second example is the effect that raising the age at marriage can have on fertility. There is a well-established relationship by which an increase in the age at marriage drives fertility re-

duction. However, there is a simple argument why this effect will be considerably less in Bangladesh than elsewhere—that is, fertility has already been reduced to levels compatible with a far greater age at marriage than currently observed. Based on current international norms, the average age at marriage could rise by over two years and Bangladesh would still have lower fertility than other countries with similar marital ages. While increasing the age at marriage has benefits for child health, its impact on fertility is likely to be limited. Strategies to reduce fertility would better target activities toward the pockets of high fertility.

Gender issues are central to health strategies in Bangladesh...more attention is needed to redressing gender biases to maintain momentum in mortality decline and fertility reduction. But traditional attitudes are not the absolute constraint on service provision as is sometimes suggested.

The study finds strong evidence of son preference, which both acts as a barrier to fertility reduction and creates excess mortality among girls. This problem is most severe in the more conservative parts of the country. Tackling gender bias should be a central aspect of health sector strategies.

It is often argued that traditional attitudes toward women in Bangladesh (*pardab*, or *seclusion of women*) act as a constraint on women's access to health services. There is some evidence that this is the case, which underlines the importance of reaching all decision-makers with behavior-change communication. Men have limited exposure to existing health and nutritional counseling, so additional channels should be sought, such as through religious authorities. However, the extent of these restrictions can be overstated. They are strongest in the most conservative areas, which is where efforts are likely to be more effective.

The Bank's BINP has improved nutritional status, but by much less than planned. Either the efficacy and efficiency of the program must be

improved, or alternatives to scaling up should be considered.

The limited impact of BINP and its cost raises serious doubts as to the justification for scaling the project up to the national level in its current form. This limited impact points to either a failing in design, in implementation, or both, and hence shortcomings in Bank performance. In order for the project to be justified, its efficacy and efficiency must be improved. In seeking to do this, several lessons emerge from this analysis which should be borne in mind:

- Supplementary feeding for children does have a positive impact, especially for the most malnourished children. NNP has revised the eligibility criteria so that only growth-faltering children with $WAZ < -2SDs$ receive feeding, as well as those with $WAZ < -4SDs$. The latter group constitutes a very small percentage of children, so there appears to be scope for raising this threshold to, say, $-3SDs$, to capture more children for whom feeding seems most successful. There is also some mistargeting, which is likely to be best addressed by further training of community nutrition promoters in interpreting growth charts.
- Supplementary feeding for pregnant women appears to be a flawed approach on two grounds: (1) the pregnancy weight gain achieved is mostly too small to have a notable impact on birth weight; and (2) it is pre-pregnancy weight that evidence suggests to be the more important determinant of birth weight. The program would be more successful if it restricted its attention to the most malnourished of women, improved targeting to reduce Type II error, and made additional efforts to discourage leakage and substitution. However, the fact that it is pre-pregnancy weight that matters suggests that a different approach altogether ought perhaps be considered, such as school feeding programs or targeting adolescent females in poorer areas.
- For both types of feeding programs, there is evidence of a greater impact in the lean season. There are grounds for considering in-

creasing the size of the food supplement in this period, restricting it to those months, or adjusting the eligibility criteria by time of year.

- Discouraging women from eating down during pregnancy has some benefit for birth weight. But all forms of knowledge transmitted by the project suffer from a knowledge-practice gap, so attention needs to be paid to both the resource constraints that create this gap and transmitting knowledge to other key actors: mothers-in-law and husbands.

However, based on current performance, scaling up will prove very costly, with limited nutritional gain. If efficiency cannot be improved through better implementation, then there are other ways of spending the money that would have a greater nutritional impact, and other channels could be sought for conveying nutritional information, tailoring it to the differing needs of different socioeconomic groups.

Rigorous impact evaluation can show which government programs and external support are contributing most to meeting poverty-reduction goals.

This study has demonstrated the impact of selected interventions and strategies on maternal and child health outcomes. These results show that impact evaluation can be used to quantify the contribution of selected activities toward meeting the MDGs' and countries' own poverty-reduction goals. It can show which interventions work and which do not, or which are more cost-effective than others.

National surveys can be used for evaluation purposes, but some adaptations would make them more powerful—notably, a more detailed community questionnaire.

This study relied on the Demographic and Health Survey to evaluate the impact of selected interventions on maternal and child health outcomes. Was this a successful approach to impact evaluation? The results reported above show that this approach can be used for certain interventions. But it could have been used to exam-

ine a broader range of interventions, and to unpack the causal chain in greater detail, if more detailed information had been available. The contrast can be made with the analysis of BINP, for which the availability of a broader range of data allowed the discussion of process issues, and so examination of the whole causal chain.

The use of DHS to evaluate HPNSP is being discussed. The experience of undertaking this study has some lessons for doing this. The first is that a more elaborate community survey should be used, including a health facility survey.¹ The community survey should record information on the various sources of health care within the community, including private phar-

macies, traditional birth attendants, NGO programs, and government health workers and the frequency of their visits. These data would aid in the investigation of changing patterns of health usage. The addition of these instruments is the easiest, and probably least costly, change to make. However, some modifications of the women's questionnaire would also be helpful in analyzing patterns of health-seeking behavior, including use of nongovernment services and problems experienced in accessing services. Seeking modifications of an existing survey instrument in this way will prove a more cost-effective approach to evaluation than undertaking separate, stand-alone surveys, and build institutional ownership of data and results within Bangladesh.

ANNEX A: TRENDS IN MATERNAL AND CHILD HEALTH OUTCOMES

Since the early 1970s, fertility rates, infant and child mortality, and malnutrition rates have declined substantially, according to the World Development Indicators (WDI) of the World Bank. However, because social indicators are often unreliable as a result of weak data collection systems, it is necessary to ask about the quality of the data. In Bangladesh many surveys provide high quality, nationally representative data, including the Demographic Health Survey (DHS), Contraceptive Prevalence Survey (CPS), Bangladesh Fertility Survey (BFS), Bangladesh Retrospective Survey of Fertility and Mortality (BRSFM), World Fertility Survey Bangladesh (WFS), the Helen Keller International Health and Nutritional Survey (HNS), and Health and Demographic Survey (HDS), and Maternal Mortality Survey (reported in Streatfield and others 2003). This annex compares the WDI data with these sources to check data quality and consistency.

Fertility

The Total Fertility Rate (TFR) indicates the expected number of births per woman during her lifetime, if she lived out her childbearing years. Between 1970 and the early 1990s, the TFR in Bangladesh dropped by 50 percent (figure A.1). During the 1990s, however, the trend plateaued; the estimated TFR remained virtually unchanged between 1993/94 and 1999/2000. This stagnation has caused major concern among policymakers. However, this report suggests that fertility continued to fall in the 1990s (see Annex F).

The TFR reported in the WDI differs little from those in national surveys (figure A.1). The higher rate in the early 1970s reported by the latter data sources reflects the different rates reported in the BRSFM for 1974, and the WFS, which is an average of the TFR between 1971 and 1975. The CPS


(1979: 82–83) argued that the rate reported in the WFS is likely to be low, because of the under-reporting of births during or in the aftermath of the independence war in the early 1970s. Since the 1980s, when the largest reduction in the TFR occurred, the various sources largely agreed with one another. There is no doubt that a profound demographic change take place in Bangladesh

Malnutrition


Malnutrition, as measured by weight-for-age z-scores (WAZ) and height-for-age z-scores (HAZ), dropped from an estimated 70 percent in 1983 to about 45 percent in 2000 (figure A.2). These figures, reported by both WHO and WDI, are based on many sources, including DHS, child nutrition surveys by the Bangladesh Bureau of Statistics, and various surveys by Helen Keller International (HKI). Even so, comparing this with

FIGURE A.1

Total Fertility Rate,
1960–2000



Sources: WDI (2004), BRSFM, CPS.

FIGURE A.2 HAZ & WAZ 1983–2000

Source: WDI (2004).

HKI data for WAZ in the 1990s (figure A.3) reveals a discrepancy; the latter reports a malnutrition prevalence 5 to 10 percent higher between 1991 and 1998. However, the HKI survey was not nationally representative until 1998. Its orientation initially was disaster-prone areas, where people are more likely to suffer from malnutrition. Thus, the HKI data overstate the national level for this earlier period, and an “end-to-end” comparison would overstate the downward trend.

Mortality


Childhood mortality has dropped appreciably since the 1960s, though the decline has been

greatest since the 1980s. Figure A.4 shows that the rate of under-five mortality fell from 250 per 1,000 live births in 1960 to less than 75 per 1,000 in 2002. That is a reduction in infant mortality by more than two-thirds (from 150 per 1,000 to about 45 per 1,000 live births) and a reduction in child mortality by more than three-quarters (from about 115 per 1,000 to 25 per 1,000 children surviving to age 1 year). The mortality series are plotted alongside the change in fertility to show that the decline in mortality occurred about a decade earlier than that of fertility. These trends are consistent with the demographic transition model, which predicts that improvements in nutrition/health and mortality rates lead to reductions in fertility, though with a lag.


Figure A.5 plots infant, child, and under-five mortality rates presented in the DHS Reports (1994, 1997, 2001), along with OED’s calculations based on pooling all three sets of data. Each data point is plotted midway through the five-year period. Rather than estimate mortality for each round, OED’s calculations pool the data from the three survey rounds; pooling makes estimates more efficient by increasing sample size. (Otherwise, the number of deaths on which the mortality calculations are based is relatively low.)

Both methods calculate mortality using the synthetic cohort probability model, (Rutstein 1984), which examines a hypothetical cohort subject to the age-specific mortality rates of one particular period; unless mortality is constant (or the period of analysis is long), the same person will not experience age-specific rates observed in one period during his or her life. The advantage of the synthetic cohorts method is that mortality rates are more sensitive to period-specific events such as natural disaster and economic crisis, affecting all cohorts simultaneously.


Although the two data series are similar, the estimates based on the pooled sample (the dashed lines in the figure, denoted by $\hat{}$) on average overestimate mortality. This difference is probably because the calculation of mortality rates uses longer recall periods. The events therefore are recorded closer to the present than they occurred (that this, “forward telescoping”).

FIGURE A.3 WAZ, 1990s

Sources: WDI (2004), HKI.

FIGURE A.4**Fertility and Mortality: The Demographic Transition in Bangladesh**

Source: WDI (2004).

FIGURE A.5**Mortality Trends**

Note: ^Calculated from pooled DHS data.

Sources: Mitra and others (1994, 1997), NIPORT and others (2001); authors' own calculations based on DHS data.

ANNEX B: CROSS-COUNTRY ANALYSIS OF CHILD HEALTH AND NUTRITION OUTCOMES

Some initial insights regarding the main determinants of child health and nutritional outcomes can be sought from cross-country regressions. This is not a novel area of analysis. Work emanating from the World Bank (Reutlinger and Selowsky 1976; Berg 1983), and followed up more recently in a series of papers by Haddad and others (e.g., Smith and Haddad 2000), has shown that income growth translates into improved nutritional status, but at too low a rate to achieve international targets. Concerning mortality, there has also been debate about the respective role of economic growth versus direct support for lower mortality via health service provision (e.g., Filmer and Pritchett 1999 and the debate around the World Health Report 2000; see Pedersen 2002).

This annex presents cross-country regression results drawing on two data sources. The first set of data updates the work of Haddad and Smith, using the same sources and series, but including more recent data. Their main results are replicated and extended through the addition of further variables and a closer look at the non-income drivers of changes in nutritional outcomes, specifically for the case of Bangladesh. DHS data, customarily used for the sort of household modeling presented in Annexes C and D, can also be used for cross-country analysis using national averages of the various variables, which are provided on the Macro International Web site. This second set of estimates is provided to corroborate the first set of results, and as some additional variables can be included.

Drivers of Maternal and Child Health and Nutritional Outcomes in Bangladesh

Smith and Haddad (2000) estimate two sets of models for the determinants of the proportion

of children who are underweight, using a cross-country data set culled from various sources. The first model regresses nutritional status on what they call the “basic determinants” of per capita GDP and democracy (measured by the Freedom House political freedom index). The second model, of underlying determinants, includes access to safe water, female secondary school enrollments; the ratio of male to female life expectancy, as a measure of gender inequality; and per capita dietary energy supply. The logic of the approach is that the latter variables, which directly affect nutritional status, depend, in turn (though only in part), on the “basic determinants.” The basic determinants equation is thus a reduced-form equation, though one that appears to suffer from omitted variable bias, as there are other determinants of the underlying variables than GDP and democracy alone. Using the results from a fixed-effects estimate of their underlying determinants model, they examine the relative importance of the different variables in the explained change in nutrition. Female education comes out as the most important factor (43 percent), followed by food supply (21 percent), then health environment (safe water, 19 percent), and finally women’s status (11 percent).

The analysis for this report has updated their data set with five modifications: (1) the analysis is carried out for four dependent variables: HAZ, WAZ, under-five mortality, and the total fertility rate; (2) more recent data have been obtained; (3) rather than using data for specific years, decade averages are used for the 1970s, 1980s, 1990s, and the current decade; (4) female literacy is used rather than secondary enrollment, as the stock of female education appears more

appropriate than the new flow into that stock, which is what the enrollment rate measures; (5) income inequality has been added to the data set, although this variable restricts sample size so that estimates are presented with and without the variable. As in Smith and Haddad, results are presented from both ordinary least squares (OLS) and fixed effects (FE) models, using time dummies in the latter and regional and time dummies in the former.

The results from the “basic determinants” model demonstrate the extent to which income growth has been behind improvements in maternal and child health and nutrition outcomes. There are both direct and indirect channels from income to better health and nutrition. The direct channel is from the higher consumption of food and health services made possible by higher income. Indirect channels include increasing both the supply and demand for education, improved water supply, and other facilities. To the extent that factors positively affecting health and nutrition outcomes are positively correlated with income, but not wholly explained by income, then the regression of any one outcome on income will provide an upward biased estimate of the income effect, which may be regarded as an “upper limit” estimate of this effect.

Table B.1 shows the extent to which income growth in Bangladesh has determined the change in the four outcomes being considered here. The results are based on the model 1 OLS estimates for each variable, shown in tables B.2(a)–B.5(a). Multiplying the coefficient on income by the change in Bangladesh’s (log) income gives the change in the outcome—for

example, under-five mortality, attributable to the income effect. This effect is highest in the case of mortality, in which case it is just under one-third, and lowest in the case of fertility, where it is just 16 percent.

The same argument may be made graphically, as is done in figure B.1. The four graphs in figure B.1 show this clear link between income and four social outcomes—under-five mortality, the total fertility rate, and the prevalence of stunting and underweight among under-fives—using the cross-section data.

In the 1980s, Bangladesh lay above the fitted line (the solid line in each figure) for under-five mortality and fertility, meaning that those indicators were worse than should be expected for a country at its income level. If these indicators had improved following the internationally established relationship with income, then subsequent observations for Bangladesh would lie along the dashed line. But these later observations lie below the fitted line, showing that Bangladesh now does better than expected for a country at its income level.

In the case of the nutrition indicators, the observations from all three decades lie above the fitted line. Bangladesh continues to have worse nutritional outcomes than the average for countries at a similar income level. But this discrepancy has narrowed over time. As for mortality and fertility, the rate of progress in nutritional status exceeds that to be expected from income growth alone.

Estimation found that inequality affects nutritional outcomes once an interactive term is included. Table B.6 shows some simulations to

TABLE B.1


Income Growth Accounts for at Most One-Third of the Reduction in Mortality in Bangladesh . . . And Less Than a Fifth of Lower Fertility

	1980 actual	2000 actual	2000 income-based estimate	Percent reduction explained by income
Under-five mortality	205.0	77.5	163.1	32.9
Total fertility rate	5.6	3.0	5.2	16.0
Stunting	67.6	44.7	62.3	23.1
Underweight	69.5	47.7	64.6	22.4

Source: Calculated from data used for figure B.1.

FIGURE B.1

Bangladesh's Improvement in Social Outcomes Is Greater Than Can Be Explained by Economic Growth Alone



help interpret this result. First note, that for any given level of inequality, HAZ falls as income rises, which is to be expected. However, HAZ is also *decreasing* in inequality up to a certain level of income (log GDP of 7.7, or just over \$2,200 per person), after which level inequality is detrimental to nutrition. The most likely explanation for this result is that at lower income, the average energy supply is below the minimum required to avoid malnutrition, so that if resources are equally distributed, everyone is malnourished. With inequality, at least some escape mal-

nutrition, while others are pushed further into food deprivation.

The next step in the analysis is the estimation of the underlying determinants model. While Smith and Haddad look at the contribution of the four variables to the *estimated* change in the data set as a whole, the estimates here are for the contribution to the *actual* change just for the case of Bangladesh. The “unexplained change” is captured through the residual. In addition, a further explanatory variable (immunization) is added to the under-five mortality regressions

TABLE B.2(a)		Height for Age Z-Score (HAZ), OLS						TABLE B.2(b)		HAZ Fixed Effects		
	Coeff.	Coeff.	Coeff.	Coeff.	Coeff.	Coeff.	Coeff.	Coeff. t	Coeff. t	Coeff. t	Coeff. t	Coeff. t
Log GDP per capita	-13.16*** (-12.72)	-54.19*** (-5.23)		-7.28*** (-3.53)	-33.23*** (-2.88)	-23.40** (-2.16)	-9.88*** (-4.73)	-15.77*** -5.13	-55.77*** -3.15		-11.54*** -3.06	-35.11*** -2.24
Democracy	-0.84 (-1.54)	-0.18 (-0.20)		-0.59 (-0.91)	0.00 (0.00)	-0.38 (-0.40)	-0.67 (-1.10)	0.10 0.16	-1.54 -1.19		0.94 1.47	0.47 0.41
Adjusted Gini coefficient		-5.89*** (-3.75)		-0.08* (-1.65)	-4.38** (-2.63)	-2.16 (-1.33)			-4.93 -1.54			-3.89 -1.43
Gini*GDP		0.77*** (3.73)		-0.03 (-0.67)	0.56** (2.59)	0.28 (1.33)			0.61 1.50			0.44 1.29
Safe water			-0.18*** -3.76	-81.55*** (-3.01)	-0.18** (-2.48)	-0.19** (-2.46)	-0.10** (-2.13)			0.01 0.13	0.00 0.07	-0.11* -1.98
Female literacy			-0.10*** -2.66	-0.01*** (-3.23)	-0.10* (-1.65)	-0.14** (-2.37)	-0.01 (-0.17)			-0.30*** -3.91	-0.23*** -2.99	0.05 0.26
Male:female life expectancy			-99.72*** -3.64		-66.80 (-1.80)**	5.14 (0.13)	-26.33 (-0.97)			-63.03 -1.43	-30.87 -0.72	-41.88 -0.78
Daily energy supply			-0.01*** -5.46		-0.01** (-2.03)	-0.01 (-1.56)	-0.01*** (-3.36)			-0.02*** -4.15	-0.01*** -3.08	-0.02** -2.36
Sub-Saharan Africa						-10.55*** (-3.44)	-4.88** (-2.42)					
Middle East & N. Africa						-6.80* (-1.87)						
South Asia							9.79*** (3.43)					
1980s						-7.83** (-2.18)						
1990s						-4.47 (-1.21)						
Intercept	134.65*** 18.02	448.42 5.75	189.19*** 6.91	206.75*** 7.70	402.11*** 4.39	255.56*** 2.82	168.17 6.38	151.19*** 6.45	478.68*** 3.44	153.25*** 3.38	190.50*** 4.33	418.78** 3.37
No. of observations	196	96		152	84	84	152		96		152	84

Note: *, **, *** significant at greater than 10 percent, 5 percent, and 1 percent respectively.

and under-five mortality itself appears as an explanatory variable in the fertility regression. The results are summarized in table B.7.

In the case of Bangladesh, the model does best in explaining the reduction in under-five mortality and height for age, with the included variables accounting for about 80 percent of the observed improvement. Higher immunization coverage has been a major factor in lower mortality in Bangladesh, accounting for close to one-third of the reduction. For both variables, safe

water, female literacy, and greater gender equality have all contributed to the observed improvements. The included variables do less well at explaining weight for age and fertility, accounting for only about one-third of the observed decline. That means that there have been other factors, not included in the model, behind improved outcomes. In the case of fertility, this analysis feeds directly into a debate about the extent to which reduced fertility is a result of changing socioeconomic conditions. The results

TABLE B.3(a)		Weight for Age Z-Score (WAZ), OLS						TABLE B.3(b)		WAZ Fixed Effects		
	Coeff.	Coeff.	Coeff.	Coeff.	Coeff.	Coeff.	Coeff.	Coeff. t	Coeff. t	Coeff. t	Coeff. t	Coeff. t
Log GDP per capita	-12.57*** -11.73	-48.47*** -4.53		-5.37** -2.48	-28.68** -2.29	-7.40*** -3.83	-23.93** -2.17	-10.68*** -4.82	-36.99** -2.64		-5.60** -2.16	-17.79 -1.21
Democracy	-0.13 -0.25	1.42*** 1.45		0.28 0.42	0.84 0.80	-0.04 -0.06	0.27 0.29	-0.21 -0.45	-0.93 -0.94		0.73 1.56	-0.31 -0.31
Adjusted Gini coefficient		-5.26*** -3.26			-4.19** -2.32		-2.74* -1.68		-2.55 -1.01			-1.80 -0.71
Gini*GDP		0.65*** 3.05			0.49** 2.07		0.34 1.59		0.35 1.09			0.19 0.61
Safe water			-0.13*** -2.66	-0.07 -1.34	-0.09 -1.09	-0.09** -2.05	-0.11* -1.64			-0.02 -0.70	-0.02 -0.51	-0.02 -0.45
Female literacy			-0.10*** -2.63	-0.08* -1.69	-0.08 -1.17	-0.05 -1.18	-0.02 -0.37			-0.14** -2.32	-0.13** -2.02	-0.08 -0.44
Male:female life expectancy			-125.87*** -4.51	-100.89*** -3.59	-101.58** -2.51	-38.40 -1.42	-37.04 -0.90			-30.10 -0.92	-24.14 -0.78	-83.91 -1.69
Daily energy supply			-0.01*** -4.85	-0.01*** -3.11	-0.01*** -2.69	-0.01 -4.09	-0.01** -2.60			-0.02*** -5.58	-0.01*** -4.90	-0.01* -1.92
Sub-Saharan Africa						-4.53*** -2.26	-3.72 -1.15					
South Asia						17.13*** 6.28	16.09*** 4.34					
Europe and central Asia						9.12* 1.64	7.21 0.88					
Intercept	119.35*** 7.73	405.22*** 5.02	202.79	202.74*** 7.27	414.92	155.39*** 5.71	290.41*** 3.06	105.06*** 6.25	307.78 2.78	104.58*** 3.06	130.96*** 3.87	307.79** 2.65
No. of obs.	208	99	165	159	86	159	86		99	165	159	86

Note: *, **, *** significant at greater than 10 percent, 5 percent, and 1 percent respectively.

do not support the idea that changing conditions have played no role, but this role does appear to be relatively minor (with the main change coming from under-five mortality). There is a large unexplained residual, for which the most likely candidate is family planning programs.

Estimates Using DHS Data

Demographic Health Surveys (DHS) are conducted under the auspices of Macro International. The company's Web site allows data to be downloaded from 126 surveys in the form of national averages—for example, the percentage of women in the survey age range (usually 15-50) having primary education, in addition

to national estimates of infant and child mortality and nutrition. Although DHS data do not include income, it has become common practice to create an asset index from information on ownership of a list of consumer durables, and sometimes also housing quality and education. For this analysis, an asset index was made by taking a simple average of the percentage of households owning the assets included for that country. This simple measure appears to be a good proxy for income, as shown by the clear negative relationship with nutritional deprivation and mortality; see, for example, the scatter plot for the percentage of children stunted (figure B.1). The simple correlation coefficient between the asset index and the outcomes mea-

TABLE B.4(a)	Under-Five Mortality (U5M), OLS						TABLE B.4(b)	Under-Five Mortality (U5M), Fixed Effects				
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
Log GDP per capita	-0.71*** -14.45	-0.65*** -9.11		-32.38*** -4.45	-24.84*** -3.54	-16.57* -1.71	-1.00*** -6.37	-1.66*** -5.58		13.92 1.08	19.19 1.53	13.52 0.58
Democracy	-0.08*** -3.27	-0.11*** -3.00		4.98** 2.45	3.11 1.64	-3.17 -1.19	-0.05 -1.55	-0.07 -0.98		-2.12 -1.02	-2.62 -1.29	-3.60 -0.93
Adjusted Gini Coefficient		0.02*** 3.02				1.11** 2.52		0.01 0.31				-0.47 -0.39
Safe water			-0.92*** -5.04	-0.65*** -3.25	-0.40** -2.17	-0.10 -0.43			-0.33* -1.85	-0.30* -1.72	-0.18 -1.00	0.12 0.47
Female Literacy			-1.34*** -9.62	-1.06*** -6.46	-1.10*** -6.62	-1.05*** -5.34			-1.33*** -4.21	-1.27*** -3.61	-0.77* -1.98	-1.44 -1.04
Male:female life expectancy			-100.67 -1.13	25.34 0.29	-171.70** -2.05	-71.48 -0.70			-277.36** -2.02	-308.70** -2.31	-378.51*** -2.88	81.01 0.44
Daily energy supply			-0.03*** -4.00	-0.01 -0.90	0.00 -0.34	0.00 -0.40			-0.04*** -2.98	-0.04*** -3.07	-0.03** -2.12	-0.03 -0.79
Immunization			-0.20 -1.04	-0.40** -2.16	-0.35* -1.75	-0.83*** -3.39			-0.55*** -3.35	-0.65*** -3.87	-0.32 -1.54	-0.79 -2.34
Sub-Saharan Africa					23.37*** 3.57	18.94** 2.30						
South Asia					-29.05***	-14.28 -2.98	-1.34					
Middle East & North Africa					-15.87 -1.55	-16.40 -1.51						
1980s					-12.11* -1.78	4.09 0.51					-18.62** -2.50	-0.73 -0.06
1990s					-12.46 -1.65						-26.15*** -2.72	
Intercept	10.23*** 29.08	9.15*** 16.42	451.24*** 5.02	465.86*** 5.44	595.32*** 7.31	397.75*** 4.16	12.37*** 10.42	17.43*** 7.54	638.48*** 4.30	575.33*** 3.59	536.75*** 3.46	136.90 0.45
No. of obs.	212		149	145	145	76			149		145	76

Note: *, **, *** significant at greater than 10 percent, 5 percent, and 1 percent respectively.

asures used here ranges from -0.47 (weight for height) to -0.65 (weight for age).

DHS includes a large number of variables related to access to, and use of, health services. These measures are correlated, of course, so that including them individually in the regressions creates a multicollinearity problem. To get around this problem, an index of health services was constructed using four variables. The vari-

ables were identified using two criteria: (1) relatively low correlation with other health variables, and (2) being available for most or all of the 126 observations. The four selected were: percentage of women using modern contraceptives (which partly proxies for access to health supplies), the percentage of assisted deliveries (which gives some idea of the availability of trained health workers), and, as measures of

TABLE B.5 (a)	Fertility (OLS)								
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
GDP per capita (logged)	-1.01*** -7.28	-1.33*** -7.15	-2.88*** -2.75			0.12 0.48	-0.06 -0.23	-0.08 -0.20	-0.05 -0.22
Democracy	-0.06 -0.89	-0.13 -1.39	-0.12 -1.25			0.12 1.70	0.04 0.46	0.11 1.00	0.13* 1.78
Inequality		0.05*** 3.02	-0.19 -1.21				0.03** 2.30	0.03 1.28	
Inequality x GDP			0.03 .50						
Safe water				-0.02*** -3.84	-0.01 -0.93	-0.02*** -4.08	-0.02*** -2.81	-0.02** -2.44	-0.02** -2.53
Female literacy				-0.02*** -5.53	-0.01 -1.35	-0.03*** -5.27	-0.03*** -4.92	-0.03** -2.47	-0.02*** -3.53
Female: male life expectancy				0.78 0.26	1.10 0.24	1.62 0.52	-1.62 -0.49	-6.88 -1.28	1.32 0.38
Daily energy supply				0.00*** -3.29	0.00 -1.43	0.00*** -2.89	0.00** -2.24	0.00 -1.31	0.00 -2.38
Under-five mortality (lagged)					0.01* 1.77			0.00 0.63	
South Asia									-0.19 -0.51
Sub-Saharan Africa									0.75*** 2.85
Middle East and North Africa									0.66* 1.80
Europe and Central Asia									-1.42** -2.08
1980s									-0.43 -1.42
1990s									-0.84*** -2.64
2000									-1.41*** -4.19
Intercept	12.64*** 12.63	13.29*** 9.16	24.95*** 3.16	8.69*** 2.94	4.80 1.01	7.00** 2.28	10.17*** 3.23	14.83*** 3.02	7.69** 2.19

Note: *, **, *** significant at greater than 10 percent, 5 percent, and 1 percent respectively.

TABLE B.5 (b)	Fertility (Fixed Effects)							
	(10)	(11)	(12)	(13).	(14)	(15)	(16)	(17)
GDP per capita (logged)	-0.81 -1.39	-3.35*** -4.86	-5.05** -2.15			1.24* 1.91	-0.48 -0.91	-0.15 -0.26
Democracy	-0.19 -1.58	-0.14 -0.86	-0.17 -0.99			0.18 1.55	-0.04 -0.37	-0.02 -0.18
Inequality		0.06 1.42	-0.26 -0.61				0.02 0.62	0.02 0.80
Inequality x GDP			0.04 0.76					
Safe water				0.00 0.61	0.00 0.28	-0.01 1.05	0.00 -0.79	0.00 0.42
Female literacy				-0.07*** -5.26	-0.07*** -3.65	-0.09*** -5.86	-0.10*** -3.23	-0.07* -1.92
Female: male life expectancy				6.27 0.82	7.33 0.94	5.48 0.71	-4.07 -0.76	4.33 -0.78
Daily energy supply				0.00** 2.64	0.00* -1.78	0.00*** 3.24	0.00 1.16	0.00 0.85
Under-five mortality (lagged)					0.00 0.84		0.01 0.92	0.00 0.78
South Asia								
Sub-Saharan Africa								
Middle East and North Africa								
Europe and Central Asia								
1980s								-0.26 -0.88
1990s								0.60 -1.22
2000								
Intercept	11.53** 2.63	28.31*** 5.27	41.81** 2.24	7.06 0.89	3.85 0.43	-0.12 -0.01	15.43** 2.25	12.03 1.61

Note: *, **, *** significant at greater than 10 percent, 5 percent, and 1 percent respectively.

antenatal services, the proportion of women having no antenatal services at all and the proportion receiving no tetanus vaccination. The weights were derived from principal components analysis, the last two variables entering with negative weights as should be expected.

Choice of Regressors

The approach adopted for the nutrition regressions assesses the robustness of the coefficients of the various variables. Table B.8 shows the simple regression estimate for each of the variables (equations 1-4 in table B.8), and the results when all variables are included in a single multiple regression (equation 5 in the table). In addition, to pursue a robust regression approach, all possible pairs and triplets of these variables were estimated. The main findings are as follows:

- The asset index is always significant in the simple regression, but never so in the multiple regression. Indeed, assets are not significant in two of the four regressions with three regressors for both HAZ and WHZ, and once for WAZ. This result means that the effect of assets found in the simple regression is picking up the impact of the other omitted variables that are correlated with assets, which they all are, but particularly the education variables. Of course the correlation with assets is in this case likely to be at least in part since income (for which assets are proxying) is one determinant of education.
- Health service inputs remain significant in two of the three nutrition equations, the exception being wasting (weight for height), the latter being the only case when the health index is not significant in any of the other regressions (it is not so in two of the four triplets for this regressand).
- The education variable is significant in all regressions. In the case of HAZ, secondary education worked better than primary and so was used instead.¹
- The various access to media variables were each tried, retaining in each case that which best fit for that dependent variable: watching TV for HAZ and WHZ and reading the newspaper for WAZ. These variables are significant

TABLE B.6		HAZ for Different Combinations of Income and Inequality				
		Gini coefficient				
		30	40	50	60	70
Ln GDP	6.0	84.8	71.9	59.1	46.2	33.4
	6.5	69.1	60.1	51.1	42.1	33.1
	7.0	53.4	48.3	43.1	38.0	32.9
	7.5	37.6	36.4	35.2	33.9	32.7
	8.0	21.9	24.6	27.2	29.8	32.4
	8.5	6.2	12.7	19.2	25.7	32.2
	9.0	-9.5	0.9	11.2	21.6	32.0
	9.5	-25.2	-11.0	3.2	17.5	31.7

cant in the multiple regressions (for WHZ only when the South Asia dummy is added, though the dummy just removes the significance of the newspaper variable in the WAZ equation).

- The South Asia dummy (added in equation 6 of table B.8 for each regressand) is always significantly positive, showing the worse nutritional status of that region once a range of determinants is allowed for.

These results were used as point estimates to model the drivers of change in nutritional status. The calculations were performed as follows:

- For each variable, table B.9 records the quartile values of the regressors.
- These regressors are used to calculate the expected value of the dependent variable, for a country with the respective set of characteristics. Although they need not be exactly equal, the corresponding values do correspond quite closely to the quartile values for the appropriate outcome variable in each case.
- The absolute change is the change in the outcome that would be caused by a country moving from across the inter-quartile range. This figure is calculated as the product of the difference in the value of the independent variable and the estimated coefficient. These changes sum to the predicted change in the

TABLE B.7

Contribution to Welfare Outcomes in Bangladesh

	Coeff.	1980	2000	Change	Share of actual
(a) HAZ					
Safe water	-0.11	-4.84	-10.67	-5.8	25.5
Female literacy	-0.29	-5.9	-9.1	-3.2	13.8
Gender inequality	-90.99	-90.0	-92.6	-2.6	11.3
Daily energy supply	-0.01	-27.1	-29.5	-2.4	10.4
Constant	176.72	176.7	176.7	0.0	0.0
Predicted		53.6	34.8	-18.8	82.1
Actual		67.6	44.7	-22.9	100.0
Residual		14.0	9.9	-4.1	17.9
(b) WAZ					
Safe water	-0.02	-0.99015	-2.18282	-1.2	5.5
Female literacy	-0.14	-2.9	-4.4	-1.5	7.1
Gender inequality	-30.10	-29.8	-30.6	-0.9	3.9
Daily energy supply	-0.01	-33.4	-36.3	-2.9	13.5
Constant	104.58	104.6	104.6	0.0	0.0
Predicted		38.5	31.0	-7.5	34.5
Actual		69.5	47.7	-21.8	100.0
Residual		30.9	16.7	-14.2	65.5
(c) U5M					
Safe water	-0.33379	-14.6869	-32.378	-17.7	13.9
Female literacy	-1.32569	-26.6	-40.8	-14.2	11.1
Gender inequality	-277.356	-274.5	-282.3	-7.9	6.2
Daily energy supply	-0.04125	-83.0	-90.3	-7.3	5.7
Immunization	-0.54844	-5.3	-43.9	-38.6	30.3
Constant	638.4797	638.5	638.5	0.0	0.0
Predicted		249.1	148.8	-100.3	78.7
Actual		205.0	77.5	-127.5	100.0
Residual		-44.1	-71.3	-27.2	21.3
(d) TFR					
Safe water	0	0	0	0.0	0.0
Female literacy	-0.01192	-0.2	-0.4	-0.1	5.0
Gender inequality	0	0.0	0.0	0.0	0.0
Daily energy supply	-0.00051	-1.0	-1.1	-0.1	3.5
Lagged under-five mortality	0.007539	1.7	1.0	-0.7	26.6
Constant	5.093549	5.1	5.1	0.0	0.0
Predicted		5.5	4.6	-0.9	35.1
Actual		5.6	3.0	-2.6	100.0
Residual		0.1	-1.6	-1.7	64.9

outcome variable. Hence, the changes from each separate regressor can be expressed as a percentage of the total change.

The results show that the smallest effect usually comes from the change in the asset index (which recall was anyhow insignificant), with education and media accounting for the bulk of the difference for the first two outcomes. In these first two cases, health care accounts for a bit under 20 percent of the difference, and rather more for WAZ, for which assets also appear more important.

What these results show is that improvements in education and access to media (presumably as a medium of health and nutritional advice) can bring about substantial reductions in nutritional deprivation. It may not be accurate to say these reductions can be brought about “independent of income growth” since that growth may be needed to generate both supply and demand for education and media services. Having said that, neither of these variables is driven solely by income, so there is scope for autonomous improvements in nutrition.

Somewhat similar results emerge for the mortality regressions (table B.10), although only the multiple regression results are shown here. The dependent variables are neonatal mortality (first month), postnatal mortality (months 1–11), and child mortality. As with nutrition, assets are significant in a simple regression but never so in the multiple regression. In these regressions it was possible to include both primary and secondary education and get significant results, despite the very high level of correlation between these two regressors. For infants, secondary is significant and primary is not, having even the wrong sign for postnatal mortality (which is also true if secondary is omitted). But for child mortality, primary is significant and secondary not. A possible interpretation of this finding is that the child rearing skills needed to ensure child health are of a more rudimentary nature than those required for infants. The health index is significant in all three regressions. In addition to health, a variable is included on the percentage of women



FIGURE B.2**Contribution of Different Factors to Improved Welfare Outcomes in Bangladesh****FIGURE B.3****The Income-Outcome Relationship in Cross-Country DHS Data**

TABLE B.8	Regression-Based Estimates of Contribution to Welfare Changes					
	(1)	(2)	(3)	(4)	(5)	(6)
Height for age						
Asset index	-0.73*** -7.19				-0.07 -0.38	0.11 -0.75
Health service index		-0.23*** -6.49			-0.12* -2.01	-0.10** -2.12
Secondary education			-0.26*** -6.60		-0.18* -1.85	-0.14* -1.85
Watch TV				-0.21*** -5.78	-0.13** -2.17	-0.15*** -2.99
South Asia dummy						14.93*** 4.99
Intercept	48.20*** 17.32	35.21*** 26.85	36.22*** 24.14	37.27*** 18.95	43.37*** 14.23	41.96*** 17.12
Number of observations	81	86	104	79	47	47
R-squared	0.40	0.33	0.30	0.30	0.57	0.73
Weight for height						
Asset index	-0.27*** -4.85				0.01 0.12	0.00 0.03
Health service index		-0.10*** -5.65			-0.04 -1.45	-0.04 -1.63
Primary education			-0.12*** -8.58		-0.13*** -4.87	-0.12*** -4.31
Watch TV				-0.08*** -3.96	-0.04 -1.55	-0.04* -1.64
South Asia dummy						3.02*** 1.88
Intercept	14.88*** 9.93	9.88*** 15.43	15.05*** 15.02	10.51*** 10.12	18.58*** 10.85	17.74*** 10.29
Number of observations	81	86	104	79	47	47
R-squared	0.23	0.28	0.42	0.17	0.70	0.73
Weight for age						
Asset index	-0.87*** -7.52				-0.18 -1.18	-0.32*** -2.78
Health service index		-0.262*** -6.91			-0.15** -2.19	-0.16*** -3.24
Primary education			-0.3*** -8.58		-0.14* -1.98	-0.07 -1.44
Read newspaper				-0.36*** -5.92	-0.18** -2.26	-0.11* -1.83
South Asia dummy						18.93*** 6.50
Intercept	46.16*** 14.61	30.471*** 21.85	42*** 17.1	32.8*** 14.8	48.363*** 11.42	42.59*** 13.73
Number of observations	81	86	104	79	47	47
R-squared	0.42	0.36	0.42	0.29	0.66	0.84

Notes: (1) In the multiple regression for HAZ, primary education in fact has the wrong sign, though it is not significant.

*, **, *** significant at greater than 10 percent, 5 percent, and 1 percent respectively.

TABLE B.9**Regression-Based Estimates of Contribution to Welfare Changes**

					Percentage difference	
	Coefficient	First quartile	Median	Second quartile	Absolute	Share
Height for age						
Asset index	−0.11	18.8	23.1	31.6	−1.39	8.3
Health service index	−0.10	10.0	26.3	42.2	−3.16	18.9
Secondary education	−0.14	12.2	24.3	43.6	−4.55	27.2
Watch TV	−0.15	24.9	35.7	77.2	−7.63	45.6
South Asia dummy	14.93	0	0	0	—	—
Intercept	41.96	1	1	1	—	—
Predicted		33.5	28.1	16.8	−16.7	100.0
Weight for height						
Asset index	0.002	18.8	23.1	31.6	0.03	−0.3
Health service index	−0.04	10.0	26.3	42.2	−1.44	16.1
Primary education	−0.12	43.3	71.9	89.2	−5.35	60.1
Watch TV	−0.04	24.9	35.7	77.2	−2.15	24.1
South Asia dummy	3.02	0	0	0	—	—
Intercept	17.74	1	1	1	—	—
Predicted		11.3	6.8	2.4	−8.9	100.0
Weight for age						
Asset index	−0.32	18.8	23.1	31.6	−4.04	24.5
Health service index	−0.16	10.0	26.3	42.2	−5.16	31.3
Primary education	−0.07	43.3	71.9	89.2	−3.32	20.1
Read newspaper	−0.11	13.1	26.2	49.8	−3.97	24.1
South Asia dummy	18.93	0	0	0	—	—
Intercept	42.59	1	1	1	—	—
Predicted		30.5	23.0	14.0	−16.5	100.0

aged 15–19 having no children. This variable picks up two effects: (1) the direct adverse effect of younger mothers on child survival chances, and (2) as a proxy for fertility. Finally, electricity is used here instead of the media variable (there is a high correlation between electricity and each of the media variables). Although only significant in one of the three regressions (CMR) (and just not so for PNM), electricity proved powerful in many specifications estimated during model specification testing, usually being the variable that knocked assets out of the equation.

The simulations were calculated in the same way as were those for nutrition with one difference. In this case the coefficients were obtained from re-estimated equations including only the significant variables. This was done to avoid perverse effects, such as the positive coefficient on

assets. Both education and health (including reproductive health captured in the no-birth variable) account for between one-third and one-half of the decline in mortality.

Conclusions

This annex has presented cross-country evidence of the determinants of child health and nutritional outcomes. It has also analyzed the importance of the different drivers of changes in these outcomes, at both the general level and specifically in the case of Bangladesh. The motivation behind this analysis is to help identify the selection of sectors for study in this evaluation. To this end, the main findings are as follows:

- While health inputs do matter for child health outcomes, they are by no means the sole—

TABLE B.10	Mortality Regression Estimates Using DHS Cross-Country Data		
	Neonatal mortality	Postnatal mortality	Child mortality
Asset index	0.21 1.42	0.27 1.04	0.40 0.81
Primary education	−0.08 −1.40	0.17* 1.73	−0.58*** −3.02
Secondary education	−0.20** −2.39	−0.48*** −3.24	0.19 0.65
No births, 15–19	−0.22* −1.85	−0.57*** −2.84	−1.29*** −3.32
Health index	−0.11** −2.25	−0.20** −2.33	−0.38** −2.32
Electricity	−0.02 −0.36	−0.18 −1.53	−0.39* −1.75
Sub-Saharan Africa	5.19 1.55	0.78 0.13	8.04 0.72
South Asia	13.08*** 3.73	−14.57** −2.41	−42.52*** −3.64
Middle East and North Africa	2.33 0.68	4.64 0.78	−14.69 −1.28
Intercept	60.86*** 5.42	96.83*** 5.01	208.71 5.57
Number of observations	75	75	75
R-squared	0.77	0.71	0.78

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively

nor are they always the most important—determinant of these outcomes. It makes good sense to incorporate determinants from other sectors.

- Education emerges as a very robust determinant of child health and nutritional outcomes. The evidence is mixed on whether primary education is sufficient for there to be a beneficial effect, or whether secondary education is required. The quality of education is a factor underlying this distinction. If many children are still illiterate after primary school graduation, then its beneficial effects will be limited.
- Access to media is important for nutrition, presumably reflecting access to health and

nutrition information. For mortality, electrification provides a better fit. Although this variable is highly correlated with access to media, electrification may provide additional channels for better health than simply health information.

- Activities in support of economic growth are one way in which external assistance can improve health and nutrition outcomes. However, the evidence suggests that these channels have not been the dominant ones, especially in Bangladesh.
- In summary, these findings support the choice of sectors for this study—that is, health and nutrition, female secondary education, and rural electrification.

TABLE B.11**Mortality Simulations**

	Coefficient	Bottom quartile	Median	Upper quartile	Change	
					Absolute	Share
Neonatal mortality						
Primary education	−0.11	43.3	71.9	89.2	−5.3	27.0
Secondary education	−0.16	12.2	24.3	43.6	−5.0	25.4
No births, 15–19	−0.42	76.5	84.9	91.0	−6.2	31.5
Health index	−0.10	10.0	26.3	42.2	−3.2	16.1
Sub-Saharan Africa	0.99	0	0	0	0.0	
South Asia	8.57	0	0	0	0.0	
Middle East and North Africa	2.22	0	0	0	0.0	
Europe and Central Asia	3.33	0	0	0	0.0	
Intercept	85.69	1	1	1	0.0	
Predicted		45.3	35.0	25.8	−19.5	
Postnatal mortality						
Secondary education	−0.32	12.2	24.3	43.6	−10.2	30.2
No births, 15–19	−0.55	76.5	84.9	91.0	−8.0	23.7
Health index	−0.11	10.0	26.3	42.2	−3.4	10.1
Electricity	−0.22	14.5	33.0	70.6	−12.1	36.0
Sub-Saharan Africa	−4.29	0.0	0.0	0.0	0.0	
South Asia	−19.87	0.0	0.0	0.0	0.0	
Middle East and North Africa	2.62	0.0	0.0	0.0	0.0	
Intercept	111.05	1.0	1.0	1.0	0.0	
Predicted		60.9	46.6	27.2	−33.6	100.0
Infant mortality						
Primary education	−0.51	43.3	71.9	89.2	−23.4	34.7
No births, 15–19	−1.30	76.5	84.9	91.0	−18.9	28.1
Health index	−0.37	10.0	26.3	42.2	−11.8	17.6
Electricity	−0.24	14.5	33.0	70.6	−13.3	19.7
Sub-Saharan Africa	7.88	0.0	0.0	0.0	0.0	
South Asia	−41.81	0.0	0.0	0.0	0.0	
Middle East and North Africa	−14.13	0.0	0.0	0.0	0.0	
Intercept	214.42	1.0	1.0	1.0	0.0	
Predicted		85.6	49.8	18.3	−67.3	100.0

ANNEX C. NEONATAL, POSTNATAL, AND CHILD MORTALITY IN THE 1990S

Reducing mortality in childhood has been a major component of health and population sector strategy in Bangladesh, and is now included in the Poverty Reduction Strategy (Bangladesh Government/IMF 2003). Health and nutrition programs have received substantial donor support (Annex H). Infant and child survival prospects have improved dramatically in recent decades. Under-five mortality rates fell by more than one-third during the mid-1980s to late-1990s (figure C.1).¹ Analysis suggests that mortality differentials have narrowed between “poor” and “non-poor” socioeconomic groups (Appendix C.3).

Gender discrimination has important implications for child survival prospects in South Asia (Sen 1998; Croll 2001), and discrimination against girl children in Bangladesh is among the most severe in the world (see below). While mortality has declined among both sexes, post-neonatal mortality among girls (that is, girls aged between 1 and 60 months of age) remains higher than among boys, and the difference between the sexes was relatively stagnant during the 1980s and 1990s (figure C.2). Socioeconomic inequalities in mortality have narrowed over time, falling more clearly for male children and neonates than all other groups of under-fives (Appendix C.3); in contrast, socioeconomic inequality in post-neonatal mortality is higher among girls and has remained constant over time at best.

The general improvement in survival chances in Bangladesh has been well documented. Ten years ago, Kabir and Amin (1993, p. 10) stated that “the fall in infant mortality [since the early 1980s] may be attributed to the large-scale immunization of children from the early 1980s, development of the health infrastructure at the

grassroots level and increased awareness of health care issues among the population through the intervention of mass communications channels.” This paper analyzes the determinants of mortality during the 1990s using Demographic and Health Survey (DHS) data, and attempts to ascertain the weight of various factors in achieving the improvement in infant and child survival using decomposition methods.


Modeling Childhood Mortality Risk

The usual point of reference for analyzing mortality determinants, at least in the demographic literature, is Mosley and Chen’s (1984) framework, which models mortality outcomes as being determined proximately by factors such as nutrient availability, use of health services, injuries and maternal fertility, which in turn depend on the underlying biological conditions (child-specific), socioeconomic and behavioral conditions (mother or household-specific), and environmental conditions including service provision (community-specific).

The economic literature derives from the household’s utility maximization problem. Households have a utility function—determining their preference orderings over consumption of different goods, including child health and nutritional status and leisure—which is maximized subject to the household’s labor constraint and the health and nutrition production functions. The health and nutrition production functions are specified as:

$$H \text{ and } N = F(NUTR_{-1}, H_{-1}, INC, MOTHEd, \\ CHILD, HCARE, ENV)$$


H and N are, respectively, health and nutritional status produced by each child, which depend on the child’s nutrition and health histories, $NUTR_{-1}$

FIGURE C.1 Mortality Rates (per 1,000)


Source: Calculated from DHS data.

and H_{-1} , household income, INC , mother's education and knowledge, $MOTHEd$, child-specific factors, $CHILD$ (such as gender, age, birth order), utilization of health services, $HCARE$, and factors determining exposure to environmental risk, ENV (for example, water, sanitation and pollution) (see Chamarbagwala and others 2004).

The economic model is generally consistent with demographic interpretations in terms of identifying determinants of health status.

FIGURE C.2 Female-to-Male Mortality Ratio (percent)


Source: Calculated from DHS data.

However, it also emphasizes that decisions on health are taken jointly with other household decisions such as labor supply. Thus, current income is considered endogenous to health status, so that its inclusion in multiple regression analysis results in inconsistent parameter estimates. A solution is to use wealth as a proxy for income, as it is unlikely to be endogenous to current nutrition/health choices—thus producing consistent estimates.² Furthermore, determining the impact of household decision variables—that is, variables determined in part behaviorally, such as utilization of health care—is problematic because of unobserved heterogeneity. In other words, where heterogeneous household behavioral characteristics cannot be fully accounted for in the set of explanatory variables, they are subsumed into the error term. Hence, if these characteristics are correlated with other explanatory variables, inconsistent estimates result (Rosenzweig and Schultz 1983). Two approaches can be taken to ameliorate this problem. The “health production function” approach estimates proximate determinants of health status using multi-stage estimation where the first-stage equation predicts the behavioral factors for each observation. However, this approach requires identifying instruments exogenous to child health status.³ The alternative is to estimate reduced-form “health demand functions,” linking health outcome to underlying variables affecting behavioral choices—for example, presence of health and family planning facilities in the community, or distance from that facility.

The demand function approach is a popular solution, particularly in analysis of mortality using survival models. However, by not including proximate determinants as explanatory variables, the approach is likely to understate the importance of factors such as health care. Moreover, it is these proximate determinants which are the most interesting for policymakers since they are most amenable to policy intervention. The remainder of this section reviews empirical evidence regarding which variables affect mortality outcomes, drawing on a meta-analysis carried out for this study (Chamarbagwala and others 2004).

Child-Specific Factors

Biological factors are important causes of death in infancy, particularly in the first month of life. Mother's age is usually found to have a convex (U-shaped) effect on infant mortality. Infants born to mothers under 20 and over 35 are at higher risk of death, because women under 20 are less likely to have fully developed reproductive systems, while the reproductive systems of women over 35 may be deteriorated. The relationship between mother's age and post-infant survival is due to other factors, for example because children of older women are more likely to be competing with other (younger) siblings; the relationship between mother's age on child mortality is therefore more likely to be linear, although children of very young mothers may suffer through the lack of maturity of the mother. This is more likely to be a factor when first births occur at a young age, as is the case in Bangladesh.

Child's gender has a strong relationship with survival chances. For biological reasons, males have a higher natural risk of death at all ages. However, it is well known that preference for boys over girls is strong throughout Asia, including Bangladesh (Croll 2001), with discrimination aggravated during crisis periods (see Bairagi 1986). Human interventions override biological factors at later stages in infancy, to the extent that, while neonatal mortality is typically higher among boys, under-five mortality rates are higher among girls. In the multivariate regression studies included in the meta-analysis, the coefficient on female child was insignificant or significantly negative for infant mortality, whereas in studies of child mortality the coefficient was insignificant or significantly positive (Chamarbagwala and others 2004). The estimated effect of sex on mortality is likely to be conservative due to sex-selective omissions in reported birth histories, which Bairagi, Islam, and Barua (1999) note will artificially lower mortality estimates among girls, although this effect is less in countries where the technology is not widely available.

The relationship between birth order and mortality risk is likely to be convex (U-shaped), reflecting mother's age, social preferences, and food availability for older children. On the one

hand, mortality is usually high among firstborns, particularly in infancy, partly because these births are more likely to occur before the mother has reached physical maturity. On the other hand, high birth order (i.e., being born later than other siblings) is expected to increase mortality risk for two reasons: physiologically, because women who have had many pregnancies are more likely to be physically depleted, and behaviorally, particularly where birth spacing is lower, due to constraints on household resources. However, women who have had more children are likely to be those experiencing higher mortality rates among their offspring. This will lead to spurious correlation between birth order and mortality risk due to unobserved correlations between siblings (frailty effects). It is important to control for this source of heterogeneity in mortality risk across families (see below).

Children are at higher risk of mortality when born after short intervals.^{4, 5} This reflects a number of factors (Koenig and others 1990):

- Biological factors among prenatals (fetal growth harmed by weakened nutritional reserves of the mother, who has been recently pregnant and lactating) and among infants where potential milk production is impaired.
- Socioeconomic and behavioral factors, i.e., competition among post-neonatal siblings for nutrition and mother's care, where parents' choices determine which child is favored. Either the younger child is neglected in favor of the older child, or vice versa, so that short subsequent birth interval becomes an important mortality risk factor—e.g., due to a new pregnancy, children are weaned earlier without receiving adequate nutritional substitutes for breast milk. Alternatively, both may be disadvantaged due to limited total family resources.
- Disease, since a higher number of young children in the household means that the likelihood of contracting contagions such as measles is higher.

Son preference may also have an indirect effect on girls' mortality rates when parents

choose to reduce birth interval because they are unhappy with the current gender composition of their offspring (Koenig and others 1990).

Miller and others (1992) find that mortality risk is higher among children born less than 15 months after the preceding child than those born more than 15 months after or those who are firstborns, reflecting the likelihood that, for birth intervals less than 15 months, conception occurred before the preceding child was 6 months old, therefore increasing chances that the child would have been weaned early.⁶ The meta-analysis found that short preceding birth interval increases infant mortality risk, while short subsequent birth interval increases child mortality risk (Chamarbagwala and others 2004).

The effects of birth interval, gender, and birth order are interlinked. In particular, in South Asia, mortality risks are likely to be high among girls who have one or more older sisters (Muhuri and Preston 1991; Muhuri and Menken 1997; Masset and White 2003) or among girls with no male siblings (Croll 2001).⁷ Some important interactions to consider are between birth order and sex, and between sex of index child and sex composition of older siblings (Masset and White 2003).⁸

Birth order and birth interval are proxies for fertility, so these relationships between birth order, birth interval, and mortality risk indicate that lower fertility means lower mortality. Although birth order and interval have the direct effects described above, they are also picking up the simultaneous relationship between declining fertility and mortality that occurs as the demographic transition proceeds and parents decide to invest in child quality rather than quantity. There will therefore be relationships between these variables and those such as use of health services for the child.

Community-Level Factors

Access to decent health care is an important determinant of infant and child survival. Babies delivered in modern health facilities by trained health professionals will have lower risks of death, since presence of a health professional reduces the mortality risk associated with complications during and after labor, while hygienic conditions reduce risk of infection of mother

and child. However, where home delivery is the norm, it can be the case that only complicated deliveries take place in health facilities, thus giving a perverse relationship. Immunization should prevent exposure to major killers such as measles, lowering risk of death among children immunized as well as having positive external effects for child survival among the non-immunized in a given community. The meta-analysis of Chamarbagwala and others (2004) indicates that these factors are indeed significantly correlated with mortality in multivariate studies.

Sanitary toilet facilities and safe drinking water improve survival chances by reducing exposure to diseases such as diarrhea. While it can be reasoned that water and sanitation are more likely to have greater effects as children get older and start using the services directly, Chamarbagwala and others (2004) show that, in fact, differentials in water and sanitation variables are more likely to be significant determinants of infant mortality in multivariate studies.

Electrification is usually found to have a positive impact on infant survival, but an insignificant impact on child survival (Chamarbagwala and others 2004). The channels through which access to electricity affects mortality are, however, contentious, though possible candidates are that electrification facilitates use of modern medicine in health facilities (e.g., a refrigerator allows hygienic storage of vaccines), improves transfer of health information (through television and radio), and improves domestic air quality. Electrification is, of course, highly correlated with locality (particularly rural versus urban) and it is also likely that electrification expansion favors less remote and better off communities (which are able to pay for the services), raising issues of reverse causality at community level. Controlling for socioeconomic status, broadly, should account for this.

Water, sanitation, electricity, and health services are labeled as community-level variables, because communities gain access to them at the same time (e.g., due to electrification programs, construction of a health center, etc.) and because it is recognized that there are strong positive externalities from improved services, e.g., in terms of disease control (the degree of coverage

in the community has an important effect on reducing morbidity and mortality among the community). However, where intra-community variation occurs due to variation in household income and preferences and where externalities do not exist, “community level” may be an erroneous classification. This is likely to be particularly true for aspects of health care such as antenatal care, where access is determined at the community level by presence of a nearby health facility, but whether parents seek professional health care is determined at family level by such factors as income and maternal education and preferences.

Most multivariate analyses control for physical locality (region and rural/urban sector), to represent unobserved determinants of mortality risk, mainly in terms of access to health care and clean water and sanitation, and possibly geographic socio-cultural factors. Chamarbagwala and others (2004) showed that urban location is frequently related to significantly lower mortality risk in multivariate models.

Household-Level Socioeconomic Factors⁹

Socioeconomic status determines nutrition, illness, and ability to access health care. Household socioeconomic status represents direct determinants of mortality such as nutrient intake and likelihood of affording health treatment. Multivariate studies generally find income or wealth to have a significantly positive impact on child survival, though no impact on infant survival prospects (Chamarbagwala and others 2004).

Education of household members is often used to proxy socioeconomic status as an indicator of potential earnings capacity of the labor force. In analyses of child health, however, there is a clear distinction between mother’s and father’s education. Whereas father’s education may affect child health indirectly through socioeconomic status, mother’s education has clear direct links to children for a number of reasons. Education can proxy for basic knowledge about nutrition and child care, including knowledge of good caring practices, health problems and treatment, and the importance of immunization and techniques such as use of oral rehydration salts

(ORS) to treat diarrhea. Educated mothers are more likely to seek treatment from health professionals, and see this as their right. Moreover, educated mothers may also have greater decision-making ability within their families and, therefore, greater say on child nutrition and health decisions (Caldwell 1979). Chamarbagwala and others (2004) find that parental, particularly maternal, education has a positive effect on infant and child survival in studies of mortality.

In a highly patriarchal society such as Bangladesh, marriage disruption caused by divorce, separation, or abandonment is potentially disastrous for women and therefore children. Where women have no support network, or are disowned by their families, they will be highly vulnerable and may have to remarry, or accept a new husband they know is already married. In such cases, the children may not fare better if they are not welcome to the new husband’s family. The effect of divorce is likely to be greater for infants because young children suffer more from reduced quantity and quality of mother’s care, e.g., because she is working to support the family (Bhuiya and Chowdhury 1997).

Unobserved Factors

Unobserved factors determining mortality among families include genetic or other family-specific practices altering childhood survival prospects and are referred to in the literature as “frailty” effects. As noted above, it is important to account for sources of unobserved heterogeneity in empirical estimation, since correlation between observations can lead to inconsistent and inefficient statistical estimates, as demonstrated for survival models by Heckman and Singer (1984). Statistical tools can be used to account for frailty, e.g., by assuming that observations at child level are correlated among mothers; however, when modeling frailty among siblings, it is also important to control for the death of the preceding child separately, since the association between immediate pairs of siblings may be stronger (Zenger 1993).

Data and Methodology

Data used for modeling mortality in Bangladesh come from the nationally representative Demo-

graphic and Health Survey. DHS collects data on complete fertility histories of all ever-married women aged 10-49, as well as data on health status and health care, contraception, women's knowledge, beliefs and caring practices, household composition, and other socioeconomic factors such as water and sanitation facilities, household asset ownership, and educational attainment. Data were collected over three rounds during the 1990s: 1993/94, 1996/97, and 1999/2000.

Most empirical studies analyze mortality determinants for infants (up to one year old) and children (between one and approximately five years old) separately. Biological and environmental factors should be more important for infants, while socioeconomic and behavioral factors become more important as children get older, though some behavioral factors, such as delivery assistance and antenatal care, are also important for neonatal survival prospects.¹⁰ It is also useful to divide infant mortality further into neonatal (first month after birth) and postnatal (second to twelfth months). Appendix C.1 summarizes results of other multivariate studies examining childhood mortality in Bangladesh.

The major econometric issue with mortality analysis is that survey data are right-censored: survival status of children born within the period of analysis is unknown past the survey date. This introduces a bias in conventional regression estimates, such as those obtained from probit/logit regression, unless observations are omitted for children born within x time periods before the survey, where x is the mortality rate being analyzed. Survival analysis, by accounting for right-censoring, enables all observations to be included and is therefore statistically more efficient.

A frequently used model of mortality is the Cox proportional hazards model.¹¹ The model assumes that the risk of death for any age can be calculated by adjusting the baseline risk for that age by an exponential set of factors. The model can be summarized as follows:

$$h_i(t) = h_0(t) \exp(\beta x_i)$$

where $h_i(t)$ represents the hazard (mortality) rate at time (t) for individual i and $h_0(t)$ is the

age-specific baseline hazard, which is the mortality risk at each age in the case where all explanatory variables are equal to zero (i.e., equivalent to the constant term in a standard linear regression); x_i is the matrix of covariates and the estimated parameters. Child mortality is modeled using the proportional hazards specification. For neonates and postnates, however, it is reasonable to drop observations on, respectively, children born in the month of the interview and children born in the year before the interview and estimate using a probit model. The probit model is summarized as:

$$P(y_i = 1) = \Phi(\beta x_i)$$

where $P(y_i = 1)$ is the probability of the event occurring for individual i , in the case of neonatal death in the first month after birth or for postnates between the second and twelfth months following birth, and Φ is the standard normal cumulative density.

A determinant can alter risk of death through two routes: where the effect of the factor changes over time (i.e., changes in the coefficient estimate); and where the distribution of the factor changes among the population (i.e., changes in the mean of the variable). A decomposition analysis is undertaken following model estimation in order to determine the latter.

As noted above, there are likely to be issues of unobserved heterogeneity of important proximate causes of health status relating to behavioral choices which will cause biased estimation of the "true" (technological) impact of the explanatory variables on health outcomes. Behavioral factors, such whether the mother seeks antenatal care during pregnancy and delivery assistance, are likely to be endogenous to health status—i.e., she seeks health care during pregnancy because she is experiencing complications or previously experienced complications during birth, which are likely to be correlated among her children due to frailty. On the other hand, immunization is likely to be less endogenous than antenatal care, since it is preventative rather than curative.¹² But even where reverse causality is not likely to be a serious issue, unobserved heterogeneity in preferences will lead to

erroneous estimates of the effect of decision variables, particularly those relating to caring practices. Some sources of heterogeneity can be controlled for with other explanatory variables—for example, maternal education and indicators of knowledge of modern health care practices. In addition, modeling frailty should account for some of the unobserved heterogeneity (though it is unable to account for the likelihood of behavior changing from birth to birth as it is assumed that frailty is constant across births for the same mother).

It is particularly likely that antenatal visits and delivery assistance are endogenous to child health, both because of reverse causality and unobserved heterogeneity with respect to mothers' behavioral choices. Under these conditions, consistency requires use of instrumental variables. Greene (1998; see also Greene 2000, p. 849) indicates that an endogenous system of limited dependent variable equations can be appropriately modeled as a multivariate probit model. Suppressing individual subscripts, the trivariate model can be expressed as:

$$P(y_1 = 1, y_2 = 1, y_3 = 1) = \Phi_3(\beta_1 x_1 + \gamma y_2 + \delta y_3, \beta_2 x_2, \beta_3 x_3, \rho_{21}, \rho_{31}, \rho_{32}).$$

Φ_3 indicates the trivariate standard normal density; y_1 , y_2 , and y_3 are, respectively, the probabilities of dying, receiving antenatal care, and delivery assistance; the equation estimating probability of dying includes antenatal care in the explanatory variables set, where γ and δ are the estimated parameters on antenatal visits and assisted delivery; $\beta_2 x_2$ and $\beta_3 x_3$ are the vectors of estimated coefficients and explanatory variables of the equations for antenatal visits and assisted delivery; ρ represents the estimated correlation between error terms of each equation—their statistical significance, therefore, provides evidence for endogeneity. Identifying instruments used in the equations for antenatal visits and delivery assistance are presence of health and information facilities in the community, distances from Upazilla and District headquarters; in addition, the equation for antenatal visits includes the non-self geographical area mean (by maternal education attainment and sex of child) of the dependent variable.

Estimation Results

The estimation strategy is to pool data for the three survey rounds and distinguish between two periods, the late-1980s to early-1990s and the mid-late 1990s. Due to the likelihood that the relationships between some covariates and mortality change during early childhood, separate regressions are estimated for neonatal mortality (less than 1 completed month), postnatal mortality (from 1 to 11 completed months), and child mortality (from 12 to 59 months).

Table C.1 presents bivariate estimates of mortality rates by various factors. Mortality rates are lower among wealthier households, though there is little difference between mortality rates among the bottom four quintiles. Mortality is higher in households with no electricity, poor sanitation, and non-piped (direct into the home) drinking water. The Divisions of Chittagong¹³ and Barisal have the highest mortality rates while Khulna has the lowest; girls fare worst in Chittagong. Mortality rates are lower among children whose mothers have formal education, especially when they have at least upper secondary education (grade 10 onwards), are not divorced or remarried, and are more mobile (can go out and/or can go to the health center). Child-specific characteristics associated with higher mortality risk include multiple birth, short preceding birth interval (less than 15 months), and death of previous child. Higher mortality rates are experienced by boys in the neonatal period, but by girls in the post-neonatal period. Children whose mothers received antenatal care, two or more tetanus toxoid (TT) vaccinations or who have been breastfed, have better survival chances.¹⁴ Mortality rates are lower among children living in communities where over half of the children have received at least one vaccination or received Vitamin A in the preceding six months. Babies delivered by a doctor have higher neonatal mortality rates than those delivered by nurses or traditional birth attendants (TBAs), suggesting that doctor-assisted births are more likely to occur for high-risk cases.

Neonatal Mortality

Table C.2 presents estimates of the probit analysis of neonatal mortality. The table reports coef-

TABLE C.1

Mortality Rates by Selected Characteristics

	Neonatal mortality rate (standard error)	Infant mortality rate (standard error)	Child mortality rate (standard error)
All children	0.047 (0.002)	0.077 (0.002)	0.033 (0.001)
Characteristics of household			
Wealth quintile 1	0.054 (0.004)	0.095 (0.005)	0.051 (0.004)
Wealth quintile 2	0.050 (0.004)	0.085 (0.005)	0.044 (0.004)
Wealth quintile 3	0.050 (0.004)	0.078 (0.005)	0.031 (0.003)
Wealth quintile 4	0.049 (0.004)	0.075 (0.004)	0.027 (0.003)
Wealth quintile 5	0.034 (0.003)	0.054 (0.004)	0.017 (0.002)
No electricity	0.052 (0.002)	0.084 (0.002)	0.037 (0.002)
Electricity	0.034 (0.003)	0.057 (0.004)	0.021 (0.002)
Unsanitary toilet	0.052 (0.002)	0.083 (0.003)	0.038 (0.002)
Sanitary toilet	0.038 (0.003)	0.063 (0.003)	0.022 (0.002)
Non-piped water	0.048 (0.002)	0.078 (0.002)	0.033 (0.001)
Piped water	0.030 (0.007)	0.049 (0.009)	0.016 (0.005)
Characteristics of mother			
Mother not attended school	0.052 (0.002)	0.088 (0.003)	0.042 (0.002)
Mother attended some primary	0.048 (0.004)	0.079 (0.005)	0.029 (0.003)
Mother finished primary	0.039 (0.005)	0.065 (0.006)	0.022 (0.004)
Mother attended lower secondary	0.042 (0.004)	0.054 (0.004)	0.010 (0.003)
Mother attended upper secondary or higher	0.014 (0.005)	0.019 (0.006)	0.003 (0.001)
Mother married	0.047 (0.002)	0.077 (0.002)	0.032 (0.001)
Mother divorced	0.061 (0.014)	0.113 (0.019)	0.047 (0.011)
Mother married once	0.047 (0.002)	0.076 (0.002)	0.032 (0.001)
Mother remarried	0.047 (0.006)	0.095 (0.009)	0.048 (0.007)
Mobility index=0	0.059 (0.004)	0.095 (0.005)	0.036 (0.005)
Mobility index=1	0.048 (0.002)	0.077 (0.003)	0.034 (0.002)
Mobility index=2	0.037 (0.003)	0.064 (0.004)	0.028 (0.003)
Characteristics of child			
Child male	0.053 (0.002)	0.081 (0.003)	0.029 (0.002)
Child female	0.042 (0.002)	0.073 (0.003)	0.037 (0.002)
Single birth	0.044 (0.002)	0.072 (0.002)	0.032 (0.001)
Multiple birth	0.261 (0.025)	0.404 (0.028)	0.105 (0.025)
Previous interval >15	0.045 (0.002)	0.074 (0.002)	0.031 (0.001)
Previous interval <15	0.099 (0.011)	0.171 (0.014)	0.064 (0.009)
Previous child survived	0.046 (0.002)	0.074 (0.002)	0.031 (0.001)
Previous child died	0.063 (0.006)	0.102 (0.007)	0.048 (0.005)
First-born child	0.069 (0.004)	0.099 (0.004)	0.023 (0.002)
Second or third birth order	0.038 (0.002)	0.063 (0.003)	0.033 (0.002)
Fourth or fifth birth order	0.037 (0.003)	0.067 (0.004)	0.038 (0.003)
Sixth or higher birth order	0.044 (0.004)	0.089 (0.006)	0.041 (0.004)
Rural*Born Dec–Jan	0.052 (0.004)	0.078 (0.005)	0.031 (0.003)
Rural*Born Feb–Mar	0.048 (0.004)	0.074 (0.005)	0.039 (0.004)
Rural*Born Apr–May	0.039 (0.004)	0.067 (0.006)	0.044 (0.005)
Rural*Born Jun–Jul	0.050 (0.005)	0.083 (0.006)	0.039 (0.005)
Rural*Born Aug–Sep	0.049 (0.004)	0.079 (0.006)	0.034 (0.004)
Rural*Born Oct–Nov	0.053 (0.004)	0.091 (0.005)	0.030 (0.003)
Delivered by nurse	0.034 (0.006)	0.068 (0.009)	0.017 (0.008)
Delivered by trained TBA	0.043 (0.006)	0.067 (0.007)	0.034 (0.011)

TABLE C.1 Mortality Rates by Selected Characteristics (*continued*)

	Neonatal mortality rate (standard error)	Infant mortality rate (standard error)	Child mortality rate (standard error)
Characteristics of child (<i>continued</i>)			
Delivered by untrained TBA	0.045 (0.002)	0.072 (0.003)	0.030 (0.003)
Delivered by other	0.053 (0.003)	0.089 (0.004)	0.034 (0.002)
Not breastfed	0.686 (0.014)	0.766 (0.014)	0.074 (0.022)
Breastfed	0.003 (0.000)	0.029 (0.001)	0.026 (0.002)
No antenatal visits	0.051 (0.002)	0.084 (0.003)	0.035 (0.002)
Some antenatal visits	0.036 (0.003)	0.058 (0.004)	0.019 (0.003)
Less than 2 TT vaccinations	0.056 (0.002)	0.092 (0.003)	0.035 (0.002)
2 or more TT vaccinations	0.039 (0.002)	0.064 (0.003)	0.023 (0.003)
Characteristics of community			
Located in Barisal	0.047 (0.005)	0.083 (0.007)	0.038 (0.005)
Located in Chittagong	0.049 (0.003)	0.082 (0.004)	0.042 (0.003)
Located in Dhaka	0.046 (0.003)	0.081 (0.004)	0.035 (0.003)
Located in Khulna	0.043 (0.004)	0.060 (0.005)	0.014 (0.003)
Located in Rajshahi	0.049 (0.003)	0.074 (0.004)	0.026 (0.003)
Female child*located in Barisal	0.031 (0.006)	0.069 (0.009)	0.034 (0.007)
Female child*located in Chittagong	0.045 (0.004)	0.079 (0.005)	0.050 (0.004)
Female child*located in Dhaka	0.041 (0.004)	0.076 (0.005)	0.038 (0.004)
Female child*located in Khulna	0.041 (0.006)	0.060 (0.007)	0.012 (0.003)
Female child*located in Rajshahi	0.044 (0.005)	0.070 (0.006)	0.032 (0.004)
Less than half of children in community received vitamin A supplement	0.049 (0.002)	0.081 (0.002)	0.035 (0.002)
More than half of children received vitamin A supplement	0.039 (0.033)	0.059 (0.004)	0.021 (0.003)
Less than half of children in community vaccinated	0.052 (0.004)	0.088 (0.005)	0.053 (0.006)
More than half of children in community vaccinated	0.046 (0.002)	0.075 (0.002)	0.030 (0.001)

ficient estimates, as well as marginal effects (dF/dx), which indicate the effect of a unit change in the explanatory variable on the probability of death. The regressions bear out the prior expectation that biological characteristics and inadequate access to public services are key factors impeding neonatal survival prospects.

Biological factors associated with higher neonatal mortality are gender (male), multiple birth, short preceding birth interval, death of previous sibling, and very low or very high birth order (the estimated turning point is between birth order 6 and 7).¹⁵ Mother's age has a convex effect on mortality risk, the coefficients suggesting that the mother's optimal age for childbirth is during her late 20s. While these factors are likely to reflect biological characteristics, the significantly negative coefficient on the interaction

between death of previous child and birth interval, suggests that newborns are less likely to die when there are fewer children competing for resources.

Maternal characteristics are important determinants of neonatal mortality. Newborns have better survival chances if their mothers are more mobile (able to go out to the health center). Mother's schooling has a negative effect on neonatal mortality, particularly upper secondary (grades 10 through 12) and higher education, as compared to the reference category of no schooling; mothers who have primary education also have fewer neonatal fatalities, though the estimate is imprecise.¹⁶ It is possible to examine some of the channels through which education lowers child mortality.¹⁷ Specification (2) includes a variable indicating number of modern

TABLE C.2

Neonatal Mortality Determinants: Probit Regression Estimates

	(1)			(2)			(3)			(4)		
	dF/dx	Coeff.	z	dF/dx	Coeff.	z	dF/dx	Coeff.	z	dF/dx	Coeff.	z
Wealth	-0.004	-0.044	-0.42	-0.002	-0.021	-0.20	-0.003	-0.041	-0.38	-0.002	-0.018	-0.17
Electricity	-0.009	-0.114**	-2.03	-0.009	-0.111*	-1.95	-0.009	-0.113**	-1.98	-0.009	-0.110*	-1.94
Piped water & sanitary toilet	-0.027	-0.485***	-2.62	-0.028	-0.502***	-2.72	-0.028	-0.505***	-2.72	-0.028	-0.505***	-2.73
Mother no formal ed +												
Mother primary	-0.005	-0.066	-1.47	-0.004	-0.047	-1.05	-0.005	-0.055	-1.17	-0.004	-0.047	-1.05
Mother lower secondary	-0.004	-0.043	-0.66	-0.001	-0.013	-0.20	-0.003	-0.031	-0.43	-0.001	-0.014	-0.22
Mother upper secondary or more	-0.023	-0.373**	-1.97	-0.020	-0.321*	-1.68	-0.022	-0.359*	-1.80	-0.020	-0.320*	-1.68
Contraceptive knowledge				-0.004	-0.053***	-4.29	-0.004	-0.054***	-4.32	-0.004	-0.053***	-4.30
Father no formal ed +												
father primary							0.003	0.037	0.82			
Father lower secondary							0.002	0.027	0.47			
Father upper secondary or more							0.006	0.066	0.67			
Multiple birth	0.244	1.204***	10.66	0.243	1.203***	10.63	0.242	1.203***	10.62	0.242	1.203***	10.62
Age at birth	-0.004	-0.047**	-2.15	-0.004	-0.043**	-2.01	-0.004	-0.044**	-2.04	-0.004	-0.043**	-2.01
Age at birth sq	0.000	0.001**	1.99	0.000	0.001*	1.81	0.000	0.001*	1.84	0.000	0.001*	1.81
Prev child died	0.019	0.200***	3.30	0.018	0.191***	3.16	0.018	0.191***	3.15	0.018	0.191***	3.15
Prec interval <15	0.085	0.619***	6.46	0.083	0.615***	6.42	0.083	0.615***	6.43	0.083	0.613***	6.40
Prev child died x prec Interval	-0.027	-0.473***	-3.07	-0.027	-0.484***	-3.14	-0.027	-0.485***	-3.16	-0.027	-0.481***	-3.12
Birth order	-0.017	-0.197***	-5.68	-0.016	-0.186***	-5.39	-0.016	-0.186***	-5.37	-0.016	-0.186***	-5.38
Birth order sq	0.001	0.016***	5.39	0.001	0.015***	5.12	0.001	0.015***	5.11	0.001	0.015***	5.11
Female child	-0.010	-0.123***	-3.59	-0.010	-0.123***	-3.58	-0.010	-0.123***	-3.59	-0.026	-0.315***	-2.78
Female head of hh	-0.004	-0.046	-0.57	-0.004	-0.051	-0.62	-0.004	-0.053	-0.64	-0.004	-0.051	-0.62
Mother's mobility	-0.006	-0.069**	-2.46	-0.005	-0.057**	-2.02	-0.005	-0.056**	-1.99	-0.005	-0.057**	-2.02
Mother divorced	-0.001	-0.013	-0.10	-0.001	-0.010	-0.08	-0.001	-0.009	-0.07	-0.001	-0.012	-0.10
Mother remarried	-0.002	-0.026	-0.35	-0.002	-0.030	-0.40	-0.003	-0.030	-0.41	-0.002	-0.028	-0.37
Antenatal visits	-0.009	-0.110**	-2.42	-0.008	-0.095**	-2.10	-0.008	-0.097**	-2.13	-0.008	-0.095**	-2.10
TBA delivery	-0.007	-0.081**	-2.22	-0.006	-0.075**	-2.04	-0.006	-0.074**	-2.02	-0.006	-0.075**	-2.05
TTBA delivery	0.009	0.096	0.82	0.010	0.111	0.94	0.010	0.112	0.94	0.010	0.109	0.92
TTBA delivery x period 1995-99	-0.017	-0.239	-1.55	-0.017	-0.245	-1.59	-0.017	-0.249	-1.62	-0.017	-0.243	-1.58
Barisal + Chittagong	0.002	0.020	0.30	-0.002	-0.030	-0.45	-0.002	-0.027	-0.40	-0.009	-0.118	-1.38
Dhaka	-0.001	-0.012	-0.18	-0.003	-0.032	-0.48	-0.002	-0.027	-0.41	-0.008	-0.105	-1.23
Khulna	-0.005	-0.067	-0.90	-0.006	-0.075	-0.99	-0.006	-0.070	-0.93	-0.013	-0.183*	-1.83
Rajshahi	-0.002	-0.026	-0.38	-0.003	-0.034	-0.50	-0.003	-0.030	-0.45	-0.009	-0.114	-1.29
Female x Chittagong										0.020	0.214*	1.66
Female x Dhaka										0.017	0.182	1.39
Female x Khulna										0.026	0.256*	1.69
Female x Rajshahi										0.019	0.198	1.48
Rural	-0.001	-0.016	-0.24	-0.001	-0.012	-0.17	-0.001	-0.012	-0.18	-0.001	-0.010	-0.15
Born In Jan-Feb x rural + born In Feb-Mar x rural	-0.006	-0.068	-1.07	-0.006	-0.070	-1.09	-0.006	-0.070	-1.09	-0.006	-0.071	-1.11
Born in Apr-May x rural	-0.012	-0.158**	-2.27	-0.012	-0.162**	-2.32	-0.012	-0.161**	-2.32	-0.012	-0.163**	-2.35
Born in Jun-Jul x rural	-0.003	-0.038	-0.57	-0.003	-0.041	-0.62	-0.003	-0.041	-0.62	-0.003	-0.041	-0.63
Born in Aug-Sep x rural	-0.007	-0.093	-1.52	-0.008	-0.098	-1.60	-0.008	-0.098	-1.61	-0.008	-0.098	-1.61
Born in Oct-Nov x rural	-0.003	-0.035	-0.62	-0.003	-0.038	-0.68	-0.003	-0.038	-0.68	-0.003	-0.039	-0.69
Period 1995-99	-0.005	-0.063*	-1.72	-0.006	-0.075**	-2.04	-0.006	-0.073**	-1.99	-0.006	-0.075**	-2.05
Constant		-0.346	-1.21		-0.204	-0.71		-0.201	-0.70		-0.129	-0.45
# Obs	18,646			18,646			18,646			18,646		
# Deaths	876			876			876			876		
Wald chi-squared	318.5***			329.5***			330.7***			334.2***		
Log-likelihood	-3,272.9			-3,262.2			-3,261.7			-3,261.0		
Pseudo R-squared	0.060			0.063			0.063			0.063		

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively; + reference category; x interactive term; dF/dx marginal effect.

contraceptive methods known by the mother—a proxy for general knowledge of modern health and child care practices—which is highly significantly negative; schooling remains significant (though its estimated impact is marginally reduced). This suggests that the impact of maternal formal schooling on neonate health operates outside knowledge, though it is possible that maternal education is approximating household socioeconomic status, if the wealth index measures this inaccurately. Indicators of paternal educational attainment were added, in order to isolate the impact of socioeconomic status from maternal education with more certainty (specification 3). Inclusion of paternal schooling enhances significance of maternal schooling, possibly by indicating educated women's enhanced decision making in the household.

Socioeconomic status is a less important determinant of neonatal mortality risk: there is no estimated impact of wealth (nor of paternal education).¹⁸ Similarly, location (division of residence, rural/urban) does not exert an important effect on neonatal survival chances, once controlling for other factors.¹⁹ Interactions between division of residence and gender of child were included in order to test for regional differences in treatment of girls and boys (table C.2, specification 4). The results provide some evidence of gender discrimination, since newborns are significantly less likely to die in Khulna, the wealthiest division by the late 1990s, though only if they are boys, while newborn girls in Chittagong are significantly more likely to die than boys. Rural seasonality is also a significant factor behind neonatal survival, which is most likely for a child born in April-May (which follows the *boro* rice harvest and is a period of high food availability) or in August-September (when maternal calorie deficiency is least severe); in contrast neonatal mortality risk is highest during December-January (when maternal calorie deficiency is also highest) (see Bloem and others 2003).

Services such as clean water and sanitation, electrification, and antenatal and delivery care are key determinants of neonatal survival. Newborns have significantly better survival chances when they live in households with both

pipled water into the dwelling and sanitary toilet facilities (septic tank). In addition, newborns living in electrified dwellings have significantly better survival chances than others; this result is significant in specifications including a wide variety of socioeconomic controls such as wealth and parental education. However, when rural and urban regressions are estimated separately, electricity is significant in urban only (results not reported), which suggests electricity may be proxying for factors such as location in a slum.

Newborns whose mothers had one or more antenatal visit are estimated to have significantly lower mortality rates. Children delivered by TBAs had higher survival chances, while a positive impact on survival of trained TBAs was found for the later period (late 1990s) only, suggesting that quality of care provided by trained-TBAs improved over time. However, as noted above, these coefficients may overstate the impact of health care on survival. In the case of delivery assistance, home delivery assisted by a TBA is the norm—over 60 percent of children are delivered in this way—but the decision to use a TBA may be affected by whether she is perceived to deliver better quality services due to modern training. The results of re-estimation using a trivariate probit model to account for the endogeneity are presented in table C.3.

The probit for one or more antenatal visits is relatively well fit (pseudo-R-squared=0.17) and identifying instruments are highly significant: presence of a satellite clinic offering antenatal care in the community significantly increases access to antenatal care, as does access to a community television in rural areas, which is a proxy for information transfer; distances from Upazilla and district headquarters reduce the likelihood that women receive antenatal care; in addition, the non-self mean of the dependent variable is highly significant. Turning to the remaining explanatory variables, pregnant women are more likely to demand antenatal care if they are educated, if they are older, if they have good knowledge of modern health practices (proxied by knowledge of contraception), when they are identified as the head of the household, and when they are more freely mobile. Short preceding birth interval has a negative impacts on

TABLE C.3

Determinants of Neonatal Death, Antenatal Visits, and Delivery by Trained TBA: Trivariate Probit Estimates

	(1)						(2)					
	Neonatal Death		Antenatal Visit		TTBA Delivery		Neonatal Death		Antenatal Doctor/Nurse		TTBA Delivery	
	Coeff.	z	Coeff.	z	Coeff.	z	Coeff.	z	Coeff.	z	Coeff.	z
Wealth	-0.028	-0.27	0.574***	8.17	0.223**	2.32	0.016	0.15	0.642***	9.52	0.223**	2.32
Electricity	-0.115**	-2.02	0.252***	6.82	0.086*	1.67	-0.098*	-1.71	0.266***	7.58	0.086*	1.67
Piped water & sanitary toilet	-0.504***	-2.73					-0.483***	-2.61				
Mother primary	-0.049	-1.09	0.137***	4.23	0.070	1.60	-0.039	-0.87	0.168***	5.46	0.070	1.59
Mother lower secondary	-0.023	-0.34	0.382***	8.44	0.163***	2.84	0.014	0.21	0.432***	10.14	0.162***	2.82
Mother upper secondary or more	-0.346*	-1.80	0.682***	6.58	-0.312**	-2.37	-0.274	-1.42	0.728***	7.85	-0.311**	-2.37
Contraceptive knowledge	-0.054***	-4.34	0.093***	10.02	0.053***	4.37	-0.049***	-3.94	0.100***	11.41	0.053***	4.33
Multiple birth	1.208***	10.65	0.011	0.09	0.298**	2.12	1.200***	10.58	-0.231*	-1.78	0.303**	2.16
Age at birth	-0.044**	-2.03	0.021***	5.87	0.011**	2.30	-0.042*	-1.92	0.022***	6.49	0.011**	2.30
Age at birth sq	0.001*	1.83					0.001*	1.78				
Prev child died	0.190***	3.15	0.054	1.30	-0.006	-0.11	0.191***	3.15	0.017	0.39	-0.005	-0.09
Prec interval <15	0.617***	6.46	-0.165*	-1.79			0.617***	6.40	-0.104	-1.10		
Prev child died x prec interval	-0.490***	-3.18	0.284**	2.16			-0.480***	-3.11	0.276**	2.04		
Birth order	-0.184***	-5.33	-0.146***	-6.44	-0.048***	-3.08	-0.193***	-5.56	-0.130***	-5.76	-0.048***	-3.07
Birth order sq	0.015***	5.10	0.003	1.52			0.016***	5.18	0.002	1.05		
Female child	-0.123***	-3.58	0.021	0.93	-0.001	-0.05	-0.123***	-3.59	0.015	0.64	-0.001	-0.04
Female head of hh	-0.049	-0.60	0.113**	2.02	0.133*	1.85	-0.041	-0.49	0.133**	2.47	0.134*	1.86
Mother's mobility	-0.059**	-2.08	0.102***	4.82			-0.056**	-1.98	0.070***	3.36		
Mother divorced	-0.007	-0.05	-0.121	-1.06			-0.012	-0.09	-0.131	-1.15		
Mother remarried	-0.028	-0.37	-0.086	-1.54			-0.030	-0.40	-0.064	-1.12		
Antenatal visits	-0.031	-0.47										
Antenatal visit by doctor/nurse							-0.278***	-4.01				
TBA delivery	-0.074**	-2.02					-0.074**	-2.01				
TTBA delivery	-0.015	-0.11					0.007	0.05				
TTBA delivery x period 1995-99	-0.234	-1.53					-0.245	-1.59				
Chittagong	-0.031	-0.46	0.020	0.42	-0.025	-0.39	-0.029	-0.43	0.030	0.66	-0.027	-0.41
Dhaka	-0.032	-0.48	-0.009	-0.19	0.009	0.14	-0.032	-0.48	0.026	0.60	0.008	0.12
Khulna	-0.080	-1.07	0.086	1.64	-0.268***	-3.54	-0.079	-1.05	0.076	1.54	-0.271***	-3.59
Rajshahi	-0.037	-0.54	0.084*	1.75	-0.093	-1.41	-0.037	-0.54	0.045	0.98	-0.094	-1.43
Rural	-0.003	-0.04	-0.303***	-7.41	0.157***	2.71	-0.029	-0.43	-0.271***	-6.99	0.158***	2.72
Born in Feb-Mar x Rural	-0.070	-1.10					-0.067	-1.05				
Born in Apr-May x Rural	-0.161**	-2.32					-0.159**	-2.28				
Born in Jun-Jul x Rural	-0.042	-0.63					-0.041	-0.62				
Born in Aug-Sep x Rural	-0.098	-1.61					-0.093	-1.53				
Born in Oct-Nov x Rural	-0.039	-0.69					-0.034	-0.60				
Period 1995-99	-0.075**	-2.03	0.240***	9.15	0.239***	6.72	-0.072*	-1.94	0.082***	3.25	0.238***	6.71
Distance to Thana hq			-0.013***	-3.69	-0.021***	-4.43			-0.011***	-3.21	-0.022***	-4.46
Distance to District hq			-0.005***	-3.80	-0.003*	-1.81			-0.003**	-1.99	-0.003*	-1.83
Antenatal clinic			0.107***	3.98	0.103***	2.92			0.088***	3.41	0.102***	2.88
Community tv x rural			0.073**	2.34					0.056*	1.83		
Community tv					-0.273***	-6.80					-0.273***	-6.80
Mean-antenatal visit			1.379***	7.58					0.776***	5.19		
Constant	-0.210	-0.74	-1.494***	-12.20	-2.134***	-13.98	-0.218	-0.76	-1.731***	-14.59	-2.132***	-13.91
Rho (2,1)	-0.035	-1.29					0.001	0.02				
Rho (3,1)	0.056	1.49					0.058	1.52				
Rho (3,2)	0.172***	8.17					0.151***	7.37				
# Obs	18,646						18,646					
# Deaths	876						876					
Wald chi-squared	3,172.2***						3,188.0***					
Log likelihood	-5361.0						-5169.0					

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively; X interactive term.

demand for antenatal care, though the positive coefficient on the interaction between previous interval and death of previous sibling indicates that women with fewer young children are better able to attend the clinics. Finally, women demand less antenatal care the higher the child's birth order, which is to be expected as they have experience from previous births. Household variables—electrification and, particularly, wealth—exert a strong positive influence on demand for antenatal care. Finally, there is evidence that pregnant women are more likely to access services in the Divisions of Khulna and Rajshahi and during the mid-late 1990s and generally less likely to access them in rural areas.²⁰

Identifying variables are also highly significant in the probit for trained-TBA-assisted delivery (pseudo-R-squared=0.05): presence of a community antenatal clinic increases the likelihood of receiving the delivery assistance of a trained TBA, while distances from Upazilla and District headquarters and access to community television (by promoting use of professional attendants, i.e., doctors, nurses, midwives, family welfare officers) reduce it. The remaining explanatory variables are of the expected sign: household-level variables such as wealth, electricity, rural location, female headship, mother's age and contraceptive knowledge are positively correlated with trained-TBA delivery incidence. Multiple births are also more likely to be delivered by trained TBAs. Children in Khulna Division are less likely to be delivered by trained TBAs, but the incidence of delivery by health professionals is highest there. For maternal education, while mother's education to lower secondary level is associated with greater probability of receiving delivery assistance from a trained TBA, women who are educated beyond this level are less likely to.

The trivariate probit specification does not significantly alter the estimated effect of trained TBA on neonatal mortality, which remains marginally insignificant in the latter period. However, the coefficient on antenatal care in the neonatal mortality regression becomes insignificant when using the multivariate probit specification (table C.3 specification 1). The estimates of \hat{U} (RHO) indicate that errors are negatively

correlated between antenatal care and neonatal mortality equations (RHO 2,1) and positively correlated between trained-TBA and neonatal mortality equations (RHO 3,1), though the coefficients are marginally insignificant.²¹ The insignificance of antenatal care may be because quality of service is not being accounted for—it is highly unlikely that antenatal care has no impact on neonatal survival. Table C.3 specification (2) accounts for quality of antenatal care by re-estimating the trivariate probit where the second equation models antenatal visits to a qualified doctor or nurse. The fit of the probit for antenatal visit attended by doctor or nurse is again high (pseudo-R-squared=0.16) and identifying instruments are highly significant, but, in contrast to above, accounting for quality of antenatal care and potential endogeneity with respect to child health, antenatal visits (with qualified doctor/nurse) significantly reduce risk of neonatal mortality.²²

Postnatal Mortality

For the postnatal infant period, services are less prominent determinants of mortality risk (table C.4). Access to electricity and clean water and sanitation are not significantly correlated with postnatal mortality. Whether the child received vitamin A supplementation in the past six months, proxied by the (non-self) share of children in the community receiving vitamin A, has the expected negative effect on postnatal mortality, though is marginally insignificant. Child-level factors such as multiple birth, short preceding birth interval, and mother's age of delivery have a negative impact on postnatal mortality. However, it is also clear that socioeconomic level is an important determinant of postnatal mortality, demonstrated most obviously by the significance of wealth, as well as the estimated linear negative effect of maternal delivery age and the observation that the effect of fate of the preceding child is entirely through the interaction with the preceding interval, both of which suggest competition for resources among siblings increases postnatal mortality.

Of mother-level variables, coefficients on maternal mobility index and marital status are insignificant. Maternal educational attainment to

TABLE C.4

Postnatal Mortality Determinants: Probit Estimates

	(1)			(2)			(3)			(4)		
	dF/dx	Coeff.	z	dF/dx	Coeff.	z	dF/dx	Coeff.	z	dF/dx	Coeff.	z
Wealth	-0.019	-0.320**	-2.04	-0.018	-0.301*	-1.92	-0.015	-0.248	-1.55	-0.010	-0.167	-1.01
Electricity	0.002	0.027	0.38	0.002	0.033	0.45	0.002	0.035	0.49	0.003	0.042	0.58
Piped water & sanitary toilet	-0.003	-0.053	-0.25	-0.004	-0.072	-0.35	-0.005	-0.083	-0.40	-0.003	-0.059	-0.28
Mother no formal ed + Mother primary	0.003	-0.048	-0.84	-0.002	-0.032	-0.55	-0.002	-0.032	-0.55	0.000	0.002	0.04
Mother lower secondary	-0.015	-0.314***	-3.15	-0.014	-0.286***	-2.85	-0.014	-0.282***	-2.81	-0.009	-0.162	-1.63
Mother upper secondary or more	-0.019	-0.476	-1.59	-0.017	-0.428	-1.43	-0.017	-0.428	-1.44	-0.007	-0.129	-0.43
Contraceptive knowledge				-0.003	-0.045***	-2.72	-0.003	-0.045***	-2.72	-0.003	-0.044***	-2.64
Father no formal ed + father primary										-0.005	-0.086	-1.52
Father lower secondary										-0.007	-0.131*	-1.83
Father upper secondary or more										-0.018	-0.427***	-2.79
Multiple birth	0.149	1.020***	8.30	0.150	1.028***	8.36	0.149	1.025***	8.33	0.149	1.031***	8.36
Age at birth	-0.001	-0.013*	-1.91	-0.001	-0.013**	-1.98	-0.001	-0.014**	-2.00	-0.001	-0.012*	-1.82
Prev child died	0.002	0.031	0.40	0.002	0.026	0.34	0.001	0.021	0.27	0.001	0.017	0.22
Prec interval <15	0.067	0.636***	5.59	0.067	0.639***	5.60	0.065	0.631***	5.56	0.064	0.628***	5.53
Prev child died x prec interval	-0.015	-0.335*	-1.77	-0.015	-0.340*	-1.80	-0.015	-0.339*	-1.80	-0.014	-0.329*	-1.75
Birth order	-0.005	-0.075**	-2.00	-0.004	-0.065*	-1.75	-0.005	-0.083**	-2.15	-0.005	-0.084**	-2.16
Birth order sq	0.001	0.013***	4.26	0.001	0.012***	3.97	0.001	0.012***	3.98	0.001	0.012***	3.95
Female child	0.004	0.062	1.38	0.004	0.060	1.32	-0.004	-0.060	-0.76	-0.004	-0.060	-0.75
Female x birth order							0.002	0.035*	1.88	0.002	0.034*	1.85
Female x older sister							0.011	0.159	1.43	0.011	0.160	1.44
Female x older sister x wealth							-0.039	-0.656*	-1.70	-0.038	-0.656*	-1.68
Female head of hh	0.007	0.117	1.26	0.007	0.116	1.24	0.007	0.110	1.18	0.007	0.120	1.28
Mother's mobility	0.000	-0.002	-0.05	0.001	0.010	0.26	0.001	0.009	0.25	0.001	0.009	0.24
Mother divorced	0.015	0.202	1.37	0.015	0.205	1.38	0.015	0.214	1.44	0.015	0.210	1.41
Mother remarried	0.006	0.098	1.13	0.006	0.098	1.13	0.006	0.100	1.14	0.006	0.096	1.10
Barisal + Chittagong	-0.006	-0.104	-1.30	-0.008	-0.143*	-1.74	-0.008	-0.145*	-1.76	-0.008	-0.153*	-1.86
Dhaka	-0.003	-0.051	-0.63	-0.004	-0.065	-0.80	-0.004	-0.068	-0.84	-0.005	-0.083	-1.02
Khulna	-0.013	-0.276***	-2.69	-0.013	-0.278***	-2.71	-0.013	-0.280***	-2.72	-0.014	-0.296***	-2.89
Rajshahi	-0.009	-0.158*	-1.88	-0.009	-0.164*	-1.95	-0.009	-0.166**	-1.98	-0.009	-0.176**	-2.09
Rural	-0.013	-0.189**	-2.16	-0.013	-0.189**	-2.16	-0.013	-0.191**	-2.17	-0.013	-0.190**	-2.15
Born in Jan-Feb x rural + Born in Feb-Mar x rural	-0.002	-0.029	-0.34	-0.002	-0.029	-0.35	-0.002	-0.035	-0.42	-0.002	-0.038	-0.45
Born in Apr-May x rural	0.009	0.129	1.46	0.008	0.128	1.45	0.008	0.127	1.43	0.008	0.122	1.38
Born in Jun-Jul x rural	0.005	0.082	0.92	0.005	0.085	0.95	0.005	0.085	0.96	0.005	0.081	0.91
Born in Aug-Sep x rural	0.001	0.024	0.29	0.001	0.024	0.29	0.001	0.024	0.29	0.001	0.023	0.28
Born in Oct-Nov x rural	0.010	0.150**	2.03	0.010	0.151**	2.04	0.010	0.150**	2.02	0.010	0.148**	2.00
Mean-vitamin A	-0.008	-0.139	-1.45	-0.008	-0.132	-1.36	-0.008	-0.134	-1.38	-0.008	-0.135	-1.40
Period 1995-99	-0.003	-0.054	-1.17	-0.004	-0.066	-1.43	-0.004	-0.065	-1.42	-0.004	-0.075	-1.61
Constant		-1.434***	-7.04		-1.277***	-6.07		-1.218***	-5.67		-1.222***	-5.67
# Obs	14,285			14,285			14,285			14,285		
# Deaths	451			451			451			451		
Wald chi-squared	196.1***			202.6***			209.4***			214.7***		
Log-likelihood	-1,865.6			-1,861.1			-1,857.6			-1,853.0		
Pseudo R-squared	0.058			0.060			0.062			0.064		

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively; + reference category; x interactive term; dF/dx marginal effect.

lower secondary level leads to lower mortality risk in postnatal infancy, as does mother's health knowledge, as proxied by knowledge of contraceptives (specification 2). However, the magnitude and significance of maternal education is reduced when paternal education is included (specification 4).

Girls of high birth order are more likely to be discriminated against and the interaction between gender of child and sex composition of elder children provides some evidence that girls have higher mortality risk if they have living elder sisters at birth, though only among poorer households (specification 3).

There is some locational variation in postnatal mortality in multivariate analysis, as children in rural areas and in Khulna and Rajshahi have lower estimated mortality risk when controlling for other factors.²³ The negative estimated impact of urban areas on postnatal survival may reflect that, accounting for socioeconomic factors and services, urban areas are more risky environments for infants, e.g., due to increased exposure to disease in areas of high population density, greater likelihood of poor communities being directly adversely affected by annual floods (e.g., because they are forced to live on river-banks), and due to greater environmental pollution. Seasonality is again a significant factor in rural areas: postnatal infants are more likely to die if born in the months of October and November, which is the counterpart to the higher neonatal mortality observed in December-January.

It is possible to account for (self-reported) annual food availability at household level and mother's recollection of birth weight in the 1999/2000 survey round, as well as mother's agency, using an index constructed from mother's reported decisionmaking in the household.²⁴ Table C.5 presents results for 1999/2000 survey data for infant mortality, which pools observations for neonates and postnates in order to maximize number of observed deaths. Specification (1) indicates that low birth weight (moderately small and very small) children are significantly more likely to die in the infant period, but whether the household experiences food deficit during some or all of the year has no significant effect on infant mortality. Specification

(2) reports results for rural areas only and includes a variable indicating whether the household was situated in a Bangladesh Integrated Nutrition Project (BINP) area. While being in a BINP area does not significantly reduce infant mortality, the interaction with mother's schooling suggests that better educated mothers are more able to benefit from the BINP project, e.g., because they are more likely to understand child growth charts and to follow educational messages such as beneficial food practices (see Annex G).²⁵ In neither specification is mother's agency significantly correlated with infant mortality.

Child Mortality

Table C.6 presents multivariate analysis of child mortality using Cox proportional hazards model.²⁶ The model is stratified by poverty status (where poverty is defined using a threshold of the wealth index), allowing the baseline hazard to differ between poor and non-poor.²⁷ The most significant factors explaining differentials in child mortality are household wealth, vaccinations, and maternal education.

Coefficients on (non-self) community shares of children receiving vitamin A supplementation and having no vaccinations have the expected signs, indicating that access to health services has a direct positive effect on child survival prospects. Formal education of mothers reduces child mortality risk, but in contrast with results for infants, the effect of mother's knowledge of modern contraception is negative but insignificant in the child mortality regressions (specification 2); this may indicate that contraceptive knowledge is a better representation of knowledge of good care practices for infants regarding health care and feeding. Access to piped drinking water and sanitary toilet facilities negatively correlates with child mortality, though, somewhat surprisingly, this effect is insignificant. However, the interaction between schooling and water and sanitation is significantly negative, indicating complementarities in the improvement of child health status: intuitively, access to piped water and clean sanitation is unlikely to improve survival chances alone, mothers must know how to use facilities properly and children must also learn to use them, which is more likely

TABLE C.5

Infant Mortality Determinants in 1999: Probit Estimates

	(1)			(2 Rural)		
	dF/dx	Coeff.	z	dF/dx	Coeff.	z
Wealth	-0.017	-0.156	-0.88	-0.009	-0.083	-0.40
Electricity	-0.007	-0.066	-0.78	-0.009	-0.083	-0.82
Piped water & sanitary toilet	-0.017	-0.178	-0.66			
Mother no formal ed +						
Mother primary	-0.016	-0.156**	-2.14	-0.018	-0.171**	-2.02
Mother lower secondary	-0.012	-0.111	-1.08	-0.001	-0.013	-0.12
Mother upper secondary or more	-0.043	-0.621**	-2.36	-0.044	-0.691	-1.59
Contraceptive knowledge	-0.009	-0.081***	-3.97	-0.011	-0.097***	-4.14
Multiple birth	0.341	1.352***	7.77	0.362	1.418***	7.34
Age at birth	-0.004	-0.038	-1.09	-0.002	-0.018	-0.45
Age at birth sq	0.000	0.001	1.04	0.000	0.000	0.44
Prev child died	0.000	0.002	0.02	-0.006	-0.054	-0.42
Prec interval <15	0.111	0.642***	4.05	0.102	0.614***	3.25
Prev child died x prec interval	-0.027	-0.302	-1.10	-0.014	-0.143	-0.47
Birth order	-0.022	-0.200***	-3.45	-0.024	-0.218***	-3.27
Birth order sq	0.002	0.017***	3.26	0.002	0.018	3.10
Female child	-0.009	-0.082	-1.46	-0.005	-0.049	-0.77
Female head of hh	0.006	0.055	0.44	0.001	0.008	0.06
Mother's mobility	-0.001	-0.005	-0.09	0.004	0.034	0.49
Mother's agency	0.000	0.004	0.37	0.001	0.006	0.52
Barisal + Chittagong	-0.010	-0.091	-0.84	-0.011	-0.100	-0.83
Dhaka	-0.005	-0.046	-0.41	-0.009	-0.085	-0.68
Khulna	-0.024	-0.258**	-2.05	-0.025	-0.269*	-1.88
Rajshahi	-0.009	-0.080	-0.70	-0.008	-0.077	-0.61
Rural	-0.013	-0.113	-1.16			
Born in Jan-Feb x rural +						
Born in Feb-Mar x rural	-0.027	-0.287**	-2.45	-0.027	-0.291**	-2.47
Born in Apr-May x rural	0.001	0.006	0.06	0.000	0.003	0.03
Born in Jun-Jul x rural	0.000	-0.002	-0.01	0.001	0.006	0.05
Born in Aug-Sep x rural	-0.003	-0.025	-0.25	-0.001	-0.009	-0.08
Born in Oct-Nov x rural	-0.003	-0.027	-0.28	-0.002	-0.021	-0.23
Mean-vitamin A	-0.018	-0.164	-1.49	-0.012	-0.106	-0.83
Food deficit	0.005	0.041	0.58	0.005	0.047	0.60
Low birth weight	0.023	0.192***	2.87	0.025	0.209***	2.77
GINP		0.022	0.180	1.01		
GINP x mother's schooling		-0.055	-0.509***	-2.87		
Constant	-0.141	-0.300	-0.491	-0.94		
# Obs	5,626		4,172			
# Deaths	382		289			
Wald chi-squared	175.2***		154.3***			
Log-likelihood	-1,272.5		-939.7			
Pseudo R-squared	0.080		0.087			

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively; + reference category; x interactive term.

TABLE C.6

Child Mortality Determinants: Cox Regression Estimates

	(1)			(2)			(3)			(4)			(5)		
	Haz. ratio	Coeff.	z	Haz. ratio	Coeff.	z	Haz. ratio	Coeff.	z	Haz. ratio	Coeff.	z	Haz. ratio	Coeff.	z
Wealth	0.269	-1.313***	-3.03	0.278	-1.280***	-2.98	0.279	-1.278***	-2.98	0.313	-1.161***	-2.71	0.272	-1.301***	-3.02
Electricity	1.044	0.043	0.31	1.041	0.040	0.29	1.041	0.040	0.28	1.053	0.052	0.37	1.021	0.021	0.15
Piped water & sanitary toilet	0.683	-0.381	-0.64	1.797	0.586	0.98	1.814	0.596	0.99	1.840	0.610	1.01	1.769	0.570	0.93
Mother no schooling +															
Mother primary	0.836	-0.179	-1.58	0.853	-0.159	-1.39	0.860	-0.151	-1.32	0.914	-0.090	-0.78	0.834	-0.181	-1.57
Mother lower secondary	0.747	-0.291	-1.52	0.789	-0.237	-1.25	0.811	-0.210	-1.10	0.957	-0.044	-0.21	0.731	-0.313*	-1.66
Mother higher secondary or more	0.111	-2.198**	-2.14	0.147	-1.914*	-1.89	0.157	-1.849*	-1.83	0.216	-1.533	-1.49	0.121	-2.112**	-2.09
Water & sanitary x mother ed				0.281	-1.270**	-1.98	0.277	-1.283**	-2.00	0.284	-1.260*	-1.95	0.276	-1.287**	-1.98
Contraceptive knowledge				0.966	-0.035	-1.08	0.961	-0.039	-1.22	0.964	-0.037	-1.15	0.964	-0.036	-1.13
Age at birth	0.971	-0.029**	-2.29	0.971	-0.029**	-2.32	0.967	-0.034***	-2.58						
Birth order	1.106	0.100***	2.94	1.105	0.100***	2.94	1.090	0.087**	2.10	0.837	-0.177	-1.59			
Female	1.284	0.250***	2.78	1.287	0.252***	2.81	0.509	-0.675**	-1.99	0.788	-0.238	-1.56			
Female x birth order							1.279	0.246**	2.56	0.603	-0.507*	-1.76			
Female x birth order sq							0.978	-0.022**	-2.32	0.969	-0.032**	-2.49			
Previous child died	1.317	0.275**	2.35	1.312	0.272**	2.31	1.283	0.249**	2.11	1.086	0.083**	2.02			
Maternal mobility	0.984	-0.016	-0.22	0.994	-0.006	-0.08	0.988	-0.012	-0.16	0.504	-0.685**	-2.02	0.836	-0.179	-0.64
Mother divorced	1.450	0.372	1.55	1.436	0.362	1.51	1.460	0.378	1.58	1.284	0.250***	2.59			
Mother Remarried	1.369	0.314**	1.99	1.372	0.317**	2.00	1.393	0.331**	2.09	0.978	-0.022**	-2.33			
Barisal + Chittagong	0.980	-0.021	-0.13	0.954	-0.047	-0.30	0.718	-0.332	-1.59	0.985	-0.015	-0.20	0.992	-0.008	-0.10
Dhaka	0.796	-0.228	-1.39	0.789	-0.237	-1.44	0.651	-0.429**	-1.96	1.456	0.376	1.56	1.404	0.339	1.42
Khulna	0.438	-0.826***	-3.51	0.438	-0.827***	-3.52	0.444	-0.812***	-2.58	1.385	0.326**	2.06	1.300	0.262*	1.66
Rajshahi	0.660	-0.416***	-2.44	0.659	-0.417**	-2.45	0.503	-0.688***	-2.89						
Female x Chittagong							1.735	0.551*	1.75	0.715	-0.335	-1.60	0.717	-0.332	-1.59
Female x Dhaka							1.496	0.403	1.22	0.638	-0.449**	-2.04	0.642	-0.444**	-2.02
Female x Khulna							1.008	0.008	0.02	0.436	-0.829***	-2.63	0.443	-0.814***	-2.59
Female x Rajshahi							1.702	0.532	1.52	0.498	-0.697***	-2.92	0.500	-0.692***	-2.91
Rural	0.929	-0.073	-0.48	0.929	-0.074	-0.49	0.926	-0.077	-0.51	1.737	0.552*	1.76	1.748	0.559*	1.78
Mean-no vaccination	1.579	0.457***	2.57	1.551	0.439**	2.45	1.550	0.439**	2.46	1.509	0.412	1.25	1.476	0.389	1.18
Mean-vitamin A	0.638	-0.449**	-2.19	0.639	-0.448**	-2.17	0.653	-0.427**	-2.07	1.009	0.009	0.02	0.961	-0.040	-0.09
Period 1995–99	0.649	-0.432***	-3.27	0.642	-0.443***	-3.35	0.645	-0.439***	-3.32	1.714	0.539	1.54	1.645	0.498	1.42
# Obs	22,527			22,527			22,527			0.932	-0.071	-0.47	0.932	-0.070	-0.46
# Deaths	558			558			558			1.543	0.434**	2.43	1.568	0.450**	2.52
Wald chi-squared	129.1***			130.4***			144.4***			0.657	-0.420**	-2.04	0.625	-0.470**	-2.26
Log-likelihood	-5206.0			-5203.6			-5197.5			0.638	-0.450***	-3.40	0.627	-0.467***	-3.54

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively; + reference category; x interactive term.

where their mothers are educated. Maternal education becomes insignificant when controlling for paternal education (specification 4). Of the remaining mother-level variables, mobility is insignificant, but children of mothers who are remarried have significantly higher mortality risk, which may indicate children are at risk of being rejected by the new family.

The regressions indicate evidence for discrimination against girl children, who have signif-

icantly higher chances of dying than boys between age 1 and 5 years. Breaking down this effect, it is girls of higher birth order who fare worse (though, as estimated by the quadratic term, the birth order effect levels off after the sixth child) (specification 3). Children living in Khulna and Rajshahi Divisions have significantly better survival chances than elsewhere, while interactions with gender find that girls living in Chittagong Division face the highest levels of discrimination.

Maternal age at birth, birth order, and fate of previous child are all significantly correlated with child mortality risk; maternal age has a linear positive relationship and birth order a linear negative relationship with child mortality. These factors may reflect fertility and therefore be endogenous to mortality. Thus, specification (5) omits these variables, the main effect being to improve significance of education variables.

Finally, we re-estimate the child mortality model for 1999/2000 survey round, in order to include measures of food deficiency, low birth weight, mother's agency, and effects of the BINP. Table C.7 indicates mother's agency has a nega-

tive impact on child mortality, though is marginally insignificant, while low birth weight children are also more likely to die in the post-infant period. There is some evidence that children of food deficit households are more likely to die, especially if they are female—this result is statistically significant in rural areas (specification 2). However, the equation including both wealth and incidence of food deficit is over-specified: the estimated coefficient on food deficit, without controlling for wealth, is positive and significant at less than 5 percent for both girls and boys (results not reported). Finally, the regression model for rural areas suggests that the ef-

TABLE C.7 Child Mortality Determinants in 1999: Cox Regression Estimates

	(1)			(2 Rural)		
	Haz. ratio	Coeff.	z	Haz. ratio	Coeff.	z
Wealth	0.152	-1.886***	-3.06	0.105	-2.259***	-3.14
Electricity	0.983	-0.017	-0.09	1.112	0.106	0.51
Piped water & sanitary toilet	0.372	-0.988	-0.96			
Mother no schooling +						
Mother primary	0.932	-0.071	-0.45	0.978	-0.023	-0.13
Mother secondary or higher	0.835	-0.181	-0.71	0.963	-0.038	-0.13
Contraceptive knowledge	0.947	-0.054	-1.13	0.952	-0.049	-0.94
Age at birth	0.961	-0.039**	-2.26	0.962	-0.038**	-2.00
Birth order	1.145	0.136**	2.41	1.163	0.151**	2.47
Female	0.763	-0.271	-0.76	0.681	-0.383	-0.94
Female x birth order	1.289	0.254*	1.83	1.320	0.277*	1.79
Female x birth order sq	0.973	-0.027*	-1.93	0.970	-0.031**	-1.96
Previous child died	0.872	-0.137	-0.74	0.843	-0.171	-0.84
Maternal mobility	0.979	-0.022	-0.18	0.975	-0.025	-0.19
Mother agency	0.964	-0.037	-1.60	0.962	-0.039	-1.51
Mother divorced	1.664	0.509	1.47	1.673	0.514	1.38
Mother remarried	1.440	0.365*	1.66	1.495	0.402*	1.72
Barisal + Chittagong	0.975	-0.026	-0.11	0.871	-0.138	-0.55
Dhaka	0.837	-0.177	-0.72	0.802	-0.221	-0.84
Khulna	0.517	-0.660**	-2.06	0.531	-0.634*	-1.86
Rajshahi	0.727	-0.318	-1.25	0.703	-0.352	-1.32
Rural	0.837	-0.178	-0.97			
Mean-no vaccination	2.050*	0.718*	1.79	2.081	0.733*	1.64
Mean-vitamin A	0.950	-0.052	-0.16	0.890	-0.117	-0.30
Food deficit	1.300	0.263	1.32	1.501	0.406*	1.74
Food deficit x male	0.778	-0.250	-0.88	0.651	-0.429	-1.30
Low birth weight	1.550	0.438***	2.60	1.597	0.468**	2.47
BINP				1.282	0.249	0.74
BINP x mother education				0.529	-0.637**	-2.05
# Obs	11,493			8,522		
# Deaths	249			196		
Wald chi-squared	108.3***			105.3***		
Log-likelihood	-2,417.2			-1,808.6		

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively; x interactive term.

fect of BINP on child mortality was dependent on the level of maternal education: BINP had no effect on mortality for children, unless, as indicated by the significantly negative interaction between BINP and mothers' schooling, their mothers were educated.

Decompositions

Following Fairlie (2003), for the probit model the total decomposition between two periods is estimated as:

$$\bar{Y}_2 - \bar{Y}_1 = \sum_{i=1}^{N_2} \frac{\Phi(\beta X_{i2})}{N_2} - \sum_{i=1}^{N_1} \frac{\Phi(\beta X_{i1})}{N_1}.$$

That is, the change in mean probability of dying over time is calculated as the difference in means of the predicted probabilities over the two sub-samples, corresponding to periods 1 and 2. However, in order to calculate the specific contributions to the change for an individual explanatory variable requires:

$$\frac{1}{N_1} \sum_{i=1}^{N_1} \Phi(\beta W_{i1} + \delta Z_{i2}) - \Phi(\beta W_{i1} + \delta Z_{i1})$$

where vector W and variable Z comprise the total explanatory variables set, X . The contribution of variable Z to the total change over time is calculated as the change in the average predicted probability from replacing the period 1 distribution with the period 2 distribution of variable Z , holding constant the distributions of the other variables (Fairlie 2003: 4). Calculation of the decomposition requires matching observations from the two sub-samples; observations from both periods are matched according to the rank of predicted probabilities calculated from the regression estimates.²⁸

For the proportional hazards model, Masset and White (2003) have shown that the percentage contribution of each variable to the change in predicted hazard can be calculated at the means of the data:

$$\frac{\bar{Y}_2}{\bar{Y}_1} - 1 = \exp[\beta(\bar{Z}_2 - \bar{Z}_1)] - 1.$$

Tables C.8 and C.9 present results of the decompositions of coefficient estimates into their shares of the change in predicted mortality over

time. The decompositions for neonatal and postnatal mortality (table C.8) indicate that, in addition to the large (adverse) impacts of mother's age at birth and birth order on mortality, expansion of electrification, antenatal care by trained doctor or nurse, and share of mothers completing secondary school were the biggest contributors to reducing neonatal mortality, while the improvement in wealth had the biggest impact on reducing postnatal mortality. The decomposition for child mortality (table C.9) indicates that improvements in wealth, maternal secondary education, and immunization have contributed the largest impact on reducing child mortality during the decade.²⁹

Cost Effectiveness Analysis

Analysis of cost effectiveness requires an estimate of the actual number of lives saved by an intervention. Table C.10 shows the actual numbers that are used as the base case for the simulations. Data on population, crude birth rate, and infant and child mortality are taken from Khan and Yoder (1998) and the World Development Indicators database. The crude birth rate is used to calculate the number of live births each year. The infant mortality rate can be used to calculate how many of those born in each year will die as infants, giving total infant deaths. Where necessary for the calculations it is assumed that half of infant deaths are neonatal (figures not shown). It is thus possible to calculate how many of the children born that year survive to age one. Using the CMR it can be calculated how many of those children will die before their fifth birthday. These deaths are spread over the four years in the proportions, 0.5, 0.3, 0.15, and 0.05. Hence, the number of child deaths each year can be calculated. These can be added to infant deaths to get under-five deaths (not shown).

The first simulation uses the result of the cross-country analysis that immunization accounts for 30 percent of the decline in mortality since the 1980s. Immunization coverage increased markedly from 1987. The simulation assumes no reduction in mortality rates from 1988 (table C.11); the number of deaths are calculated as for the base case. The difference in the number of under-five deaths between the two cases

TABLE C.8**Decomposition (%)
for Neonatal and
Postnatal Probit Models**

	Neonatal	Postnatal
Wealth	0.30	-2.28
Electricity	-2.44	0.74
Piped water & sanitary toilet	-0.47	-0.01
Mother primary	-0.02	0.02
Mother lower secondary	0.78	-0.93
Mother upper secondary or higher	-1.51	-0.31
Contraceptive knowledge	0.12	3.01
Multiple birth	0.85	4.13
Age at birth	14.62	3.79
Age at birth sq	14.89	
Prev child died	-0.88	-0.14
Prec interval <15	0.76	4.50
Prev child died x prec interval	2.00	2.01
Birth order	39.12	21.88
Birth order sq	2.17	13.52
Female child	0.49	0.63
Female x birth order		0.18
Female x older sister		0.94
Female x older sister x wealth		0.28
Female head of hh	0.00	0.08
Mother's mobility	-0.78	0.20
Mother divorced	0.03	-0.38
Mother remarried	0.07	-0.05
Antenatal visits by doctor/nurse	-4.27	
Chittagong	-0.26	-0.40
Dhaka	0.27	1.34
Khulna	-0.35	-0.76
Rajshahi	0.28	2.19
Rural	0.79	5.72
Born in Feb-Mar x rural	0.56	0.20
Born in Apr-May x rural	0.83	0.03
Born in Jun-Jul x rural	0.17	0.01
Born in Aug-sep x rural	0.45	-0.04
Born in oct-Nov x rural	0.18	0.61
Mean-vitamin A		-0.65

Note: x interactive term.

is the number of deaths averted by the reduction in mortality; 30 percent of these are attributed to immunization.

The second simulation is based on the DHS analysis, which finds that immunization reduces the probability of death by 50 percent. The hazard ratio for the Cox regression using 1999 data was 2.05 (table C.7), but using pooled data, and a without-self mean of immunization to account for potential endogeneity, the ratio falls to 1.55,

which is the figure used here. To use this information it is necessary to know the mortality rate for unimmunized children. This can be done using the fact that the observed mortality rate is a weighted average of that for immunized and unimmunized children:

$$MR = i MR^{IMM} + (1-i) MR^{UNIMM}$$

where i is the rate of immunization coverage. The odds ratio tells us that

$$\frac{MR^{UNIMM}}{MR^{IMM}} = \frac{3}{2}$$

so given i (for which DPT coverage is used) and MR , it is possible to calculate MR^{UNIMM} . This mortality rate is used to calculate the number of deaths in the absence of immunization for infants and children separately (children not shown). It is assumed that immunization only affects postnatal deaths among infants.

The number of deaths averted provides the denominator for the cost-effectiveness calculation. Budget data for the numerator are taken from Khan and Yoder (1998). EPI expenditure data are provided from the start of the program until 1998. A more detailed calculation for 1997/98 captures all costs associated with immunization services, including those outside of EPI, which are about 30 percent greater than the direct EPI costs. Two calculations are made: (1) total immunization costs in 1998/deaths averted in 1998; and (2) program costs (from Khan and Yoder inflated for additional years and 30 percent to capture immunization costs not paid through EPI)/deaths averted over the whole period.

The results of these calculations are summarized in table C.12. The range is quite large, from just under US\$100 to nearly US\$300 per life saved. These results compare with the US\$137 reported by Khan and Yoder (1998). These calculations suggest that that result may be something of an underestimate, but that a value of US\$200 per life saved is not unreasonable.

Conclusions

The exercise to examine the determinants of mortality in the neonatal, postnatal, and child periods has yielded some interesting results:

- Household incomes, as represented in this paper by wealth, exert a significant effect on post-neonatal survival, particularly in post-infancy when income contributed the most of any one variable to reducing mortality over time.
- Mothers' education is an important factor behind variation in mortality risk across children and over time. Mothers' education, particularly at secondary or higher level, has a significantly positive impact on child survival; this is observed over the impact of household wealth, as well as mothers' knowledge of modern contraception, an indicator of general health knowledge, which itself is a significant determinant of infant mortality differentials. The results suggest that mother's health knowledge has an important effect on survival chances, but also that children of better educated mothers are more likely to survive. The latter may be due to more highly educated mothers having greater say in household decisionmaking. Evidence also indicates that neonates have better survival chances when their mother is free to leave the dwelling and visit the health center, while divorce and remarriage can be detrimental to survival in childhood.
- Health services contribute significantly to improving childhood survival chances. Mortality is lower among newborns whose mothers attended antenatal visits with a trained doctor or nurse and immunization had a strong impact on reducing mortality in childhood in Bangladesh in the 1990s. These effects are robust to use of methods that attempt to account for endogeneity and unobserved heterogeneity in mothers' preferences.
- Access to piped drinking water and sanitary toilet facilities were found to have a significant effect on reducing mortality among neonates, and among children in households where the mother has formal schooling.
- The multivariate analysis indicates that gender discrimination against girls remains strong. While the effect is quantitatively greatest among children, as would be expected, the evidence suggests that higher birth order

TABLE C.9		Decomposition for Child Mortality: Proportional Hazards Model		
	Coeff.	Late 80s –early 90s	Mid- late 90s	Decomposition (%)
Predicted hazard at means	0.053	0.041		
Wealth	–1.308***	0.240	0.332	–11.4
Electricity	0.023	0.178	0.288	0.3
Piped water & sanitary toilet	0.495	0.025	0.023	–0.1
Water & sanitary	0.495	0.025	0.023	–0.1
x mother ed	–0.787***	0.059	0.058	0.1
Mother some primary	–0.180	0.258	0.274	–0.3
Mother compl primary	–0.290	0.131	0.174	–1.3
Mother secondary or more	–2.20**	0.019	0.029	–2.2
Contraceptive knowledge	–0.035	3.663	3.767	–0.4
Female	–0.181	0.495	0.492	0.1
Maternal mobility	–0.01	1.117	1.202	–0.1
Mother divorced	0.335	0.019	0.024	0.1
Mother remarried	0.253	1.056	1.061	0.1
Chittagong	–0.349*	0.313	0.287	0.1
Dhaka	–0.46**	0.295	0.305	–0.4
Khulna	–0.834***	0.109	0.103	0.5
Rajshahi	–0.715***	0.219	0.242	–1.6
Female x Chittagong	0.561*	0.150	0.138	–0.7
Female x Dhaka	0.389	0.146	0.151	0.2
Female x Khulna	–0.036	0.058	0.051	0.0
Female x Rajshahi	0.502	0.109	0.118	0.5
Rural	–0.054	0.899	0.832	0.4
Mean-no vaccination	0.447**	0.340	0.180	–6.9
Mean-vitamin A	–0.466**	0.255	0.271	–0.7

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively; x interactive term.

girls are more likely to die in both the post-natal and post-infant periods. Analysis of DHS 1999/2000 data indicated that girl children in food deficit households are more likely to be discriminated against.

- The analysis of DHS 1999/2000 survey round data showed that infants and children living in BINP areas tended to have better survival chances where their mothers were formally educated, suggesting educated mothers were more able to understand and act upon the messages of the project and/or were more likely to take complementary actions such as use modern medical care.

TABLE C.10 Under-Five Death Calculation Based on Actual Data (base case)

Year	Population	IMR	CMR	Crude birth rate	No. of children born	Infant deaths	Children alive at age one	Child deaths of this cohort	Year 1 (.5)	Year 2 (.3)	Year 3 (.15)	Year 4 (.05)	Child deaths per year
1981	89912	126	83.9	43.0	3866	485	3381	284	142	85	43	14	
1982	92061	122	80.6	42.0	3867	472	3395	274	137	82	41	14	
1983	94169	119	77.2	41.0	3861	458	3403	263	131	79	39	13	
1984	96335	115	73.8	40.0	3853	444	3409	252	126	75	38	13	260
1985	98531	112	70.4	39.0	3843	430	3412	240	120	72	36	12	249
1986	100758	113	63.0	38.0	3829	431	3398	214	107	64	32	11	230
1987	103025	113	55.6	37.0	3812	431	3381	188	94	56	28	9	207
1988	105282	107	54.8	35.6	3744	402	3342	183	91	55	27	9	192
1989	107587	102	53.9	34.1	3671	373	3298	178	89	53	27	9	183
1990	109922	96	53.1	32.7	3592	345	3247	172	86	52	26	9	176
1991	111455	92	47.0	31.2	3482	320	3162	149	74	45	22	7	162
1992	113729	88	40.9	29.8	3389	298	3091	127	63	38	19	6	143
1993	115981	84	41.7	29.5	3417	287	3130	131	65	39	20	7	134
1994	118161	80	43.0	29.1	3441	274	3167	136	68	41	20	7	134
1995	120382	75	44.3	28.8	3465	260	3205	142	71	43	21	7	138
1996	122561	71	41.3	28.4	3486	247	3239	134	67	40	20	7	137
1997	124706	67	38.4	28.1	3504	233	3271	125	63	38	19	6	131
1998	126579	62	35.4	28.1	3559	222	3337	118	59	35	18	6	124
1999	128797	58	32.5	28.1	3624	211	3413	111	55	33	17	6	116
2000	131050	54	29.6	28.2	3690	199	3491	103	52	31	15	5	109
2001	133345	51	27.9	28.2	3758	192	3566	100	50	30	15	5	103
2002	135684	48	26.3	28.2	3826	184	3643	96	48	29	14	5	99

TABLE C.11 Mortality Simulations

Year	Simulation 1: no mortality reduction since 1985					Simulation 2: immunization reduces mortality by 50 percent						
	IMR	CMR	Infant deaths	Child deaths	Difference vs base case	IMR	IMR for immunized	IMR for unimmunized	CMR	Infant deaths	Child deaths	Difference vs base case
1981	126	84	485		126	84	126	84	487	305	0	
1982	122	81	472		122	82	122	81	473	294	0	
1983	119	77	458		119	79	119	77	460	281	0	
1984	115	74	444	260	0	115	77	116	74	446	269	204
1985	112	70	430	249	0	112	75	113	70	433	257	198
1986	112	70	429	243	11	113	76	114	63	437	230	199
1987	112	70	427	240	29	113	78	116	56	444	206	198
1988	112	70	419	236	62	107	76	113	55	424	204	182
1989	112	70	411	233	88	102	81	122	54	446	222	155
1990	112	70	402	228	109	96	83	125	53	448	232	125
1991	112	70	390	222	130	92	81	122	47	425	204	107
1992	112	70	380	216	155	88	75	113	41	382	168	101
1993	112	70	383	214	176	84	74	112	42	381	179	96
1994	112	70	385	214	192	80	74	110	43	380	194	84
1995	112	70	388	215	206	75	65	97	44	337	190	79
1996	112	70	390	217	224	71	63	95	41	332	185	73
1997	112	70	392	218	246	67	60	90	38	315	173	67
1998	112	70	399	220	273	62	57	86	35	306	166	63
1999	112	70	406	224	302	58	53	80	33	289	155	59
2000	112	70	413	228	333	54	50	75	30	276	145	56
2001	112	70	421	232	358	51	47	71	28	265	140	54
2002	112	70	429	236	382	48	45	67	26	256	136	51

TABLE C.12**Decomposition for Child Mortality: Proportional Hazards Model**

	Simulation 1	Simulation 2
1998 only	224	292
Whole program	192	90

APPENDIX C.1: Mortality Regression Results for Bangladesh Data

Study	Dataset	Dependent variable	Explanatory variables	Estimation method
Bairagi, Sutradhar, and Alam (1999)	DSS 1966-94	Infant mortality	Mother literate(--), boy(++), muslim(--), mother's age (--), birth order(--)+ square(++).	Hazard
		Child mortality	Mother literate(--), boy(--), muslim(++), mother's age (--), birth order(++)+ square(--).	Hazard
Bhuiya and Chowdhury (1997)	DSS 1975-89	Infant mortality	Mother's education(--), mother's age(--), divorced(++), girl(--), MCH project area.	Logit (infant)
		Child mortality	Mother's education(--), mother's age(++), divorced(++), girl(++), MCH project area(--).	Hazard (child)
Bhuiya and Streatfield (1991)	DSS 1982-84	Under-three mortality (children surviving to 6 mos.)	Mother's education (--), wealth(--), mother's age, girl child(++), age of child(++), boys with more highly educated mothers (-), girls with more highly educated mothers(+), MCH project area (Matlab) (--).	Hazard
Howlader and Bhuiyan (1999)	DHS 1996	Neonatal mortality	Male(++), mother aged over 20 (--), birth order(++), birth interval over 18 mos (--), previous child died(++), mother's education(--), urban, flush toilet, piped drinking water, electricity(--), landholding, antenatal care(--), assisted delivery, hospital delivery, immunization(--), tetanus vaccination(++), breastfed, visited by family planning worker.	Logit
	DHS 1996	Infant mortality	Male(++), mother aged over 20 (--), birth order(++), birth interval over 18 mos (--), previous child died(++), mother's education(--), urban, flush toilet(--), piped drinking water, electricity(--), landholding, antenatal care(--), assisted delivery, hospital delivery, immunization(--), tetanus vaccination(++), breastfed(+), visited by family planning worker(--).	Logit
	DHS 1996	Child mortality	Male(--), mother aged over 20 (--), birth order(--), birth interval over 18 mos, previous child died(++), mother's education(--), urban(--), flush toilet, piped drinking water, electricity, landholding(--), antenatal care(--), assisted delivery, hospital delivery, immunization(--), tetanus vaccination(-), breastfed(++), visited by family planning worker(--).	Logit
Kabir and Amin (2003)	Bangladesh Fertility Survey (BFS) 1989	Prop. of dead children born to each mother relative to expected mortality rate (standardised by age of deaths)	Mother's education (--), father's education (--), father's occupation (sig), religion, rural(++), region (sig), water supply, toilet (--).	OLS
Miller and others (1992)	DSS and Determinants of Natural Fertility Survey (DNFS), Matlab 1983/84	Mortality among children <24 mos.	Short conception interval interacted with higher birth order(++), proportion of previous children dead(++), mother's age(--), mother's education(-), breastfeeding(--).	Hazard
Muhuri and Menken (1997)	DSS 1981/82	Child mortality	Mother's education, wealth(--), boy with two or more older brothers and any sisters (++), girls with 1 or more older sister and any brothers(++), second child with older brother(++), birth-to-conception interval(--), birth-to-conception interval interacted with girl sex by age of child(++ for older girls), living in MCH project area (Matlab)(--), family with more schooling lives in control area(--).	Logit
Muhuri and Preston (1991)	DSS 1981/82	Under-five mortality (children surviving past 6 mos.)	Mother's education(--), wealth(--), girl with one older brother, girls with two or more older brothers(--), girl with one or more older sister(++), birth order dummies, girl in poor household(-), born in MCH project area (Matlab)(--), girl born in Matlab.	Logit
Razzaque and others (1990)	DSS	Neonatal mortality	Mother aged under 20-34(--), boy(++), wealth, famine-born, famine-conceived, famine-born interacted with sex, wealth and mother's age.	Logit
	DSS	Post-neonatal mortality	Mother aged under 20-34, boy, wealth(--), famine-born(++), famine-conceived, famine-born interacted with sex, wealth and mother's age.	Logit
	DSS	Child mortality	Mother aged under 20-34, boy(--), wealth(--), famine-born, famine-conceived, boys born in famine(++).	Logit
Zenger (1993)	DSS 1968-82	Neonatal mortality	Mother's education, mother under 20 years old(++), mother over 29 years old(+), girl child(--), Hindu(++), birth order(++), previous child died(++), post-famine birth(--), child died(++), post-famine birth(--).	Logit

Notes: +++, ++, + indicates positive correlation at significance level <1%, <5%, <10%; ---, --, - indicates negative correlation at significance level <1%, <5%, <10%.

APPENDIX C.2: Measuring Socioeconomic Status: Wealth, Assets, and Welfare Outcomes in Bangladesh

This paper reviews measures of socioeconomic status (SES), with a particular focus on wealth. The objective is to provide guidance in constructing a suitable index, which can be used to measure SES using Demographic and Health Surveys (DHS). DHS collect data on asset ownership at the household level, which can be converted into indicators of household wealth status. There are various ways of doing this, and indices have been constructed from different types of assets using different estimation methodologies. However, attention to methodology is important because alternative indices have been shown not only to identify different households as “poor,” but also to generate significantly different measures of health inequality between “non-poor” and “poor.” This paper reviews the types of indicators that are usually used for socioeconomic status, focusing specifically on assets to measure SES and summarizes the issues involved in aggregating assets into a composite index of wealth. Finally, the paper presents results of construction of an asset index for Bangladesh DHS data.

Indicators of Socioeconomic Status

Socioeconomic status is a well-established term, originating in sociological literature (see Bollen and others 1999).¹ Socioeconomic status (SES) can be broadly defined as indicating an individual or household’s relative position according to some social and/or economic stratification scale, where the latter may be determined with reference to a broad or narrow set of characteristics. The term “status” implies a relative approach to measuring well-being, though some measures, such as income/consumption, can be interpreted absolutely, given adjustments for differences across space and time. Given this conceptual ambiguity, socioeconomic status is measured in a variety of ways, the most common being with economic indicators (such as income, expenditure, and asset ownership). This

is because economic indicators are relatively easy to measure—unlike concepts of social standing, which are measured with much greater difficulty if at all—and because social and economic indicators are assumed to be strongly correlated. However, most studies also account for factors that arguably reflect social standing to a greater extent than income/wealth alone, including human capital and/or occupation.

Household income and consumption expenditure are the most commonly used measures of living standards. Consumption and income are estimated from surveys, such as the World Bank-sponsored Living Standards Measurement Surveys (LSMS), which collect detailed household-level information on income and consumption items, including imputed estimates of self-employed income and consumption of own produce. White and Masset (2003) provide a description of the steps that must be taken to ensure comparability of income and expenditure across households and time.

Indicators of physical asset ownership such as durable goods, housing quality, and access to basic services such as water and sanitation, are often used to measure socioeconomic status where income or expenditure data are unavailable, as in the case of research based on Demographic and Health Survey (DHS) data. DHS collect nationally representative information on health, nutrition, and fertility. The DHS asset module can be used to construct an index of wealth, though the main problem is in aggregation, since data on prices, which would facilitate comparisons of different assets, are not available. There are also issues regarding the choice of assets to include in the index and adjustment for household size, composition, and economies of scale. These topics are taken up below.

However, it should be clear that income/consumption and assets are not equivalent indicators. There is a theoretical justification for using assets to measure SES. Economic theory

distinguishes current income from permanent income (wealth), the latter providing a better representation of living standards over time. If credit and insurance markets operated perfectly (and foresight were perfect), households would be able to plan expenditures and borrow/save accordingly to minimize consumption fluctuations across seasons and over economic and life cycles, in order to stabilize life-time consumption. Under these conditions, current expenditure provides the best indication of permanent income (wealth). In practice, though, particularly in poor countries, household expenditure tends to fluctuate with current income because problems in saving and borrowing (and accessing insurance) prevent consumption smoothing over time (Baulch and Hoddinott 2000).² Under such conditions, assets are preferable to consumption for measuring wealth, and there is some empirical evidence supporting this; for example, using survey data from Indonesia, Nepal and Pakistan, Filmer and Pritchett (1998) argue assets to be preferable to consumption for measuring wealth because they rank households more consistently over time.

On the other hand, income represents human capital attainment and remittance flows—important determinants of current living standards that physical assets are less likely to account for. Note also that, for these reasons, the stock of physical wealth would tend to be more unequally distributed than income flows. Human capital assets have been included in asset indices by some (Sahn and Stifel 2000) to reduce the bias in measuring wealth when relying on physical assets alone; the problem with doing so is that it precludes estimation of a separate education effect.

Generating an Index

This section describes the main issues in generating an index from asset data collected by household surveys.

Before moving on, it is important to note that there is no reason to generate an index unless it is needed for some analytical purpose, e.g., in order to analyze the relationship between “wealth” and another variable. If not, and household wealth merely needs to be controlled for, as-

sets can enter separately in regressions. The main arguments against this are: (1) reduction in degrees of freedom, but this is not usually an issue with analysis of household survey data sets, given that the number of observations is often in the thousands; and (2) a high correlation between the different assets will result in multicollinearity, undermining their statistical significance.

Where households need to be ranked by wealth, in order to determine the correlation between wealth and another variable, an index can be generated by aggregating assets:

$$A_i = \psi_1 a_{i1} + \dots + \psi_n a_{in} \quad (1)$$

where A_i is the total asset score for each household i , a_n are individual asset types and ψ_n are the weights. Variables indicating ownership/access to assets are measured categorically (1 = yes; 0 = no). In all other cases, scale equivalence must be ensured, where necessary by adjusting variables to range between 0 and 1, otherwise the index will contain “hidden weights” since variables ranging over bigger scales will implicitly be weighted more heavily.³ Young Lives (2002) cites as an example of the latter the education component of UNDP’s Human Development Index (which is measured as one-third mean years of schooling to two-thirds literacy), which in the first year of implementation failed to adjust for scale equivalence of literacy (measured in percentage terms) and mean years of schooling (ranging between about 2 and 10); in the resulting education index, the majority of variation was due to literacy because of its much greater variance. One problem with rescaling is that it is sensitive to outliers, which may therefore have to be removed from the data beforehand.

The following hypothetical example (table C2.1) demonstrates the calculation of a wealth score for two households, assuming a simple index of seven household durable goods (ownership of a wardrobe, chair, and table, which all have unitary weights; ownership of telephone, TV, and bicycle, which each carry weights of 2; and ownership of a motor vehicle, with weight 3). Household 1 owning all of the items would therefore have asset score 12; household 2 own-

ing chair, table, and bicycle would be given an asset score of 4.

In the preceding example, weightings were chosen arbitrarily, though were intended to be sensible. An alternative is to allow weightings to be determined by the variation in the data using statistical methods, for example principal components analysis (PCA), as used by Filmer and Pritchett (1998). Table C2.10 presents a calculation of asset scores for two hypothetical households using weightings estimated for Bangladesh by Gwatkin and others (2000).⁴ Household A—which has electricity, radio, TV, bicycle, piped drinking water outside residence, septic tank, cement flooring, brick walls, and finished (cement/concrete) roof—has a high asset score of 3.4. Household B—which is “poorer”—works its own land, owns a radio and bicycle, has a surface well for drinking water, uses a pit latrine, and has wooden flooring and rudimentary (wooden) walls and (tin) roof, and is assigned an asset score of 0.7. If the same dwelling had only natural (earth) flooring, (bamboo) walls and (thatched) roof, it would have an asset score of –0.2.

There are three conceptual issues associated with generating an index: choosing the individual components that make up the index, determining the weighting scheme and controlling for the size and structure of the household (Gwatkin and others 2000). These issues are explored in the following sub-sections.

Choosing the Components of the Index

Asset information commonly collected in household surveys can be categorized into two main groups, or five sub-groups. Table C2.2 presents a list of assets for which data are collected in three such surveys, the Bangladesh Household Income and Expenditure Survey (HIES) 2000/01, the Bangladesh DHS 1999/2000, and the Young Lives project.⁵ The variables include:

- (1) Household assets, such as:
 - consumer durables, such as a bicycle, motor vehicle, refrigerator, radio, table, TV, and telephone;
 - housing quality, such as the types of material used to construct floor, walls and

TABLE C2.1		Simple Example of Asset Index	
	Weighting	Household 1	Household 2
Wardrobe	1	Y	N
Chair	1	Y	Y
Table	1	Y	Y
Telephone	2	Y	N
TV	2	Y	N
Bicycle	2	Y	Y
Motor vehicle	3	Y	N
Total value		12	4

Note: Y = household owns item; N = household does not own item.

ceiling and number of rooms in the dwelling;

- productive assets, such as landholdings, livestock, machinery, human capital; financial assets, such as savings.
- (2) Basic amenities, such as drinking water source, toilet facility, electricity, and fuel used for cooking, which are likely to be strongly determined by location.

The distinction between “household variables” and basic amenities is important, due to the fact that many amenities are either publicly provided or dependent on the availability of infrastructure, which is in turn strongly correlated with locational factors such as rural or urban residence, region, etc. In other words, lack of a particular facility does not necessarily indicate “poverty” at the level of the household. Of course, the classification presented above is primarily illustrative and simplified. Basic amenities coverage will vary by country/region, and will therefore be less indicative of location in countries/regions where coverage is greater. Other factors classified as household assets may also be locational in nature, e.g., productive assets and (landline) telephone, and households are unlikely to own durables such as a television or refrigerator where electricity is unavailable. On the other hand, toilet facility may be household specific.⁶

One of the key problems in using assets, broadly defined, is that some assets, while useful to proxy SES, are also important indicators of

TABLE C2.2**Assets Included in Three Household Questionnaires**

Asset group	Bangladesh HIES (2000/01)	Bangladesh DHS (1999/2000)	Young Lives (2002)
Productive assets	Land ownership Agricultural assets including tractor, farm equipment, livestock, trees	Land ownership	Land ownership Agricultural assets including tractor, farm equipment, livestock
Financial assets	Savings, investments		Savings
Household durables	TV, radio, cassette player, VCR, dish antenna/decoder, camera/camcorder, bicycle, motorcycle/scooter, motor vehicle, refrigerator/freezer, washing machine, fans, heaters, sewing machine, pressure lamps/petromax, carpet, furniture, kitchen items, watch, clock, telephone/mobile phone, other	TV, radio, almirah (wardrobe), table/chair, bench, watch/clock, cot/bed, bicycle, motorcycle, sewing machine, telephone	(Specified to be in working condition): TV, radio, refrigerator, bicycle, motorbike/scooter, motor vehicle, landline telephone, mobile phone, sewing machine (plus two other country-specific assets)
Housing quality	Floor, roof material Number of rooms & area of dwelling Home ownership	Floor, roof, wall material Home ownership	Floor, roof material Number of rooms Home ownership
Amenities	Toilet facility Drinking water source Electricity	Toilet facility Drinking water source Electricity	Toilet facility Drinking water source Electricity Fuel used for cooking

other aspects of quality of life of interest to the researcher. That is to say, including different types of assets with separate relationships with a variable of interest, such as health status, may obscure effects operating directly from indirect effects operating via economic status (Montgomery and others 2000). For example, aspects of housing quality such as flooring and amenities such as water, sanitation, and energy source⁷ are likely to have separate direct effects on health outcomes, e.g., by affecting exposure to infections, in addition to any indirect effect which operates through their use as an indicator of income or wealth. In comparison, consumer durables are likely to influence health status only indirectly through their correlation with income and wealth, though possibly ownership of a radio, TV, and telephone gives the household better access to information regarding good health practices and emergencies, especially in remote areas. Human capital of the main care-

taker, embodied in existing knowledge and ability to process new information (as strongly affected by education outcomes), is likely to have important direct effects on child health outcomes. Other productive assets are more likely to have indirect implications for health via wealth (e.g., where they act as methods of saving).

Therefore, an alternative classification may be more relevant to a study examining causal relationships, by classifying variables according to their “direct” and “indirect” impacts on a variable of interest (Houweling and others 2003). Note, however, that the definition of “direct” and “indirect” depends on the specific health outcome in question. Table C2.3 presents a matrix showing the relationship between these alternative classification schemes for health/survival status.

Given the variety of assets that are often available, and the fact that they may indicate or determine (directly or indirectly) separate aspects of well-being, there is usually a large menu of po-

TABLE C2.3

Asset Classification for Direct and Indirect Impacts on Health Status

Asset type	Impact on health/survival	
	Direct	Indirect
Consumer durables		Table, bed, radio, TV, bicycle, motor vehicle, telephone, refrigerator
Housing quality	Type of material used on construct floor, number of people per room	Type of material used to construct walls and ceiling
Productive assets	Human capital (education, knowledge)	Landholdings, livestock, farm implements, machinery
Amenities	Drinking water source, toilet facility, fuel source	

tential indices to choose from. However, recent research has demonstrated empirically that different asset indices are not equivalent, as they are often imperfectly correlated across households due to the different types of assets included. For example, Houweling and others (2003) demonstrate that only about half of households in Indonesia and Uganda are identified in the same wealth quintile of the population by asset indices created from household durables ownership alone and an index that also includes housing quality and access to amenities (water, sanitation, and electricity); most (about three-quarters) of the remaining households move one wealth group up or down.

In sum, there are good reasons to exclude certain types of assets from an index of household socioeconomic status:

- Different types of assets represent different aspects of “wealth.” Productive assets, while representing household wealth status within livelihood groups, are unsuitable for measuring wealth between livelihood groups—for example, households that do not own cattle may not be poor, but simply do not rely on herding as a livelihood. Productive assets should therefore be excluded from the asset index. However, where productive assets represent an important aspect of wealth, such as land or livestock ownership in rural areas (or, for example, where there are limited numbers of non-productive assets, or where their distribution is

highly skewed), it may be necessary to include some productive assets in order to distinguish between households. Inclusion of agricultural land may be desirable given a large rural population, despite the fact that it would also tend to mis-categorize households in urban areas as poorer than they may actually be.

- Access to certain assets will be determined by location as opposed to purchasing power of the household. Most obviously, households in rural areas are less likely to have access to amenities such as electricity, water, and sanitation, though some would not be considered poor as more conventionally defined. Therefore, amenities which are highly locational should not be used to measure household wealth. The effect of including assets determined by location is to overstate measured inequalities in household wealth. However, given that inclusion of multiple asset types as explanatory variables in regression analysis will undermine precision of estimates where the assets are strongly correlated, it may be sensible to construct separate indices for “household assets” and “amenities,” where the latter includes locational items (Houweling and others 2003).⁸
- The distinction between “direct” and “indirect” determinants of a variable of interest is important. Only indicators having indirect effects (via “wealth”) on a dependent variable, say, health status, should be included as components of an index used to measure in-

come/wealth in causal analyses. This is crucial for policy analysis because, given public resource constraints, policy may be more effective where it focuses on improving those assets that are the main drivers of change, rather than on measures to raise general wealth. The effect of including assets that have direct impacts on a variable of interest is to overstate estimated socioeconomic inequalities in that variable.⁹ For example, Houweling and others (2003) estimate that inclusion of water and sanitation variables in the index augments socioeconomic inequalities in under-five mortality, in 50 percent of cases, by attributing some of the impact of these direct factors to economic status. Excluding water and sanitation from a full index that also includes consumer durables, housing quality, and access to electricity supply, reduces inequality in under-five mortality in Chad and Malawi by more than 30 percent of the inequality index; in Brazil, Indonesia, and Pakistan, estimated inequality is reduced by between 10 and 30 percent; in Bolivia, Kenya, Tanzania, and Uganda, inequality is insignificantly affected; and in Cameroon estimated inequality rises. Excluding housing quality, water, and sanitation from the full index reduces estimated inequality by over 10 percent in 3 out of 10 (Brazil, Indonesia, and Kenya) but increases inequality in a further 3 (Chad, Malawi, and Tanzania). Additional exclusion of electricity produces estimates of inequality that are lower than the estimate produced by the full index in Bolivia, Brazil, Indonesia, Kenya, and Pakistan but higher in Chad, Malawi, and Tanzania.

The cost of excluding certain asset types is that it may prevent sufficiently continuous ranking of households, particularly in poor countries where ownership of durables may be highly limited among poorer segments of the population, or where the survey only collects limited information on durables.¹⁰ Houweling and others (2003) demonstrate for DHS data that the lack of the variation in the data may necessitate inclusion of more types of assets in order to stratify poorer households evenly into wealth quintiles.

Including the same list of assets for different countries, and weighting these assets by the same amounts, enables comparison across countries—this is relatively easy with DHS data since the questionnaire has a standard asset list included in all surveys. However, some assets may represent household wealth to different extents across countries (or even regions/sectors within countries) and thus are not strictly comparable; e.g., bicycles do not represent the prestige in Asia, where in many countries they are relatively common and cheap, that they do in most African nations, where they are rare and expensive. Sahn and Stifel (2000) generate separate weighting schemes for 11 African countries to reflect local preferences for each asset. But by doing so they lose the ability to make inter-country comparisons. Young Lives (2002), on the other hand, uses a common weighting system across countries, though it allows for two locally specific assets for each country, and retains comparability across countries by rescaling (see endnote 4).

Determining Weighting Coefficients

The weightings assigned to the individual asset components of the index can be determined using a variety of methods (Filmer and Pritchett 1998). First, weights may be assumed to be equal across assets, thus the ψ in equation (1) are equal to one and the index is simply the sum of individual asset indicators. However, as Filmer and Pritchett (1998: 4) state, this method “has as its *only* appeal not seeming as arbitrary as it really is.” In addition, equal weighting will give the least possible categories of the asset ownership index, making it more difficult to distinguish households, which is likely to be problematic given the limited number of assets on which data are usually collected; more complex weighting systems give more thorough rankings.

Second, assets may be assigned unequal weights using more arbitrary, but reasonable, methods, e.g., by making reasoned choices based on other information sources. In Young Lives (2002), the wealth index is calculated as the average of three components: (1) housing quality, which is the average of number of people per room (rescaled to vary from zero to one), and

dummies indicating finished floor, walls, and roof; (2) the sum of household ownership of a range of durable assets as a continuous variable scaled between zero and one; (3) the average value of dummy variables indicating access to drinking water, toilet, electricity, and cooking fuel. A particular advantage of the index is that by scaling total durables ownership between 0 and 1, it does not give undue weight to durables (of which there are more variables) over house quality and services, which an equally weighted index would.

Third, where data are available, weights could be determined by asset prices. Many surveys, such as the DHS, do not collect data on asset prices, though average national or regional relative prices could potentially be used instead (Young Lives 2002). However, it is not clear whether a weighting scheme based on prices is necessarily ideal—though this would create an index analogous to household income and consumption measures. Only under a highly restrictive set of assumptions do prices reflect what is measured at a theoretical level, namely marginal utilities in consumption. For example, where there exist imperfect competition, production, or consumption externalities, missing markets or imperfections in information—i.e., nearly all of the time, and especially in developing countries—prices are likely to deviate substantially from marginal utility.

Fourth, assets can be linearly aggregated into a single index using econometric methods. Filmer and Pritchett (1998, 1999) and Gwatkin and others (2000) used principal components analysis (PCA) to generate asset weightings, while Sahn and Stifel (2000) used factor analysis (FA). PCA/FA are data reduction methods that examine the statistical relationships between variables.¹¹ The first principal component/factor is the linear combination of variables explaining the maximum possible variation in the data, the second principal component/factor is that accounting for the largest of the remaining variation, etc. In analyses of asset ownership the characteristic that explains the largest variation (that estimated by the first principal component/factor) is assumed to be “wealth.” However, there are problems in attributing socioeconomic meaning to higher order components/factors, thus, only the

first is used to generate the asset index. Sahn and Stifel (2000) argue that, since FA accounts for the variation in assets using a smaller number of covariates, it is therefore preferable to PCA for generating the wealth index.¹² However, the methods tend to produce highly similar asset rankings.

There are two main concerns with PCA/FA. First, using an index may entail losing a lot of information regarding variance of asset ownership across households, but this is the case with any index. Second, PCA/FA replaces arbitrariness with statistical credibility, but does not necessarily estimate weightings that are meaningful in a socioeconomic sense—a more meaningful index may be created using an “arbitrary” but reasonable weighting system, as in Young Lives (2002). In Gwatkin and others (2000) index derived from PCA, a household that works its own or its family’s land is assigned negative weighting, which obviously reflects location (urban residents are more likely to have access to water, sanitation, and electricity utilities—as well as, therefore, other electrical assets—but less likely to work in agriculture). However, in rural areas it is the wealthier who own land (and the landless that are usually poorest), therefore the coefficient on land ownership, if indicating wealth, should enter positively. The negative coefficient on land is therefore a good example of why using productive assets can be misleading.

Another way to determine weightings econometrically is to use the coefficients estimated from a fully specified regression of the dependent variable of interest, say, height-for-age z-score, on the asset variables and all other explanatory variables (Lensink and White 2000). Given estimated coefficients, the asset index can be calculated and the regression re-estimated, replacing the individual assets with the asset index, in order for the impact of “wealth” to be determined.

Table C2.11 indicates some of the different ways in which assets have been aggregated in previous research. The studies were selected mainly, though not exclusively, as relating to DHS data and estimation of child mortality and nutrition. Madise and others (1999) use indicator variables for whether the household owns any one or more than one of a set of items. Many re-

cent studies have used an asset index similar to the one of Filmer and Pritchett (1998), which aggregates household and community-level assets into a single index. Some separate the two, e.g., Kishor and Parasuraman (1998) and Madise and others (1999), showing that amenities (water and sanitation) have separate impact on mortality, on top of household wealth. Often, the asset index is used to categorize households into asset quintiles,¹³ and/or is used to calculate socioeconomic measures of health inequality.

Controlling for Household Size and Economies of Scale

Since data are often collected at household level, an issue arises as to whether adjustment should be made for the size and possibly composition of the household. In the case of wealth, where durable goods are used as a store of value, e.g., as insurance to be sold in the event of an adverse household shock, the size of the household determines the amount of insurance per capita a given asset can provide.¹⁴ For other household goods, such as bicycle ownership, their utility under normal conditions also implies adjustment for household size.

Other assets are consumed publicly by the household—i.e., economies of scale in household consumption entail that assets can be used by one member without reducing consumption possibilities for another—and therefore adjustment for household size is not necessary. Indeed, many items in the asset index can be considered public goods, although to varying degrees, e.g., consumption of some utilities such water (unless metered), electricity (for heating), aspects of housing quality such as ceiling, wall, and floor materials, some durable goods (e.g., refrigerators, radio/TV), and to a lesser extent dwelling size (until congestion is reached) (White and Masset 2003).

The following formula is used to adjust household-level variables for household size and economies of scale:

$$A_i^* = A_i / n^{1-\alpha} \quad (2)$$

where A_i^* represents the adjusted value of the asset index, A_i the unadjusted index at household level, n the composition-adjusted number

of members, and α the scale coefficient, which ranges between 0 and 1. Zero represents no economies of scale (thus the adjusted index is simply equal to the unadjusted index expressed in per capita, or per equivalent adult, terms) and one indicates perfect economies of scale (household size does not alter the asset index). Empirical estimates of the scale coefficient usually are between 0.15 and 0.3 in developing countries (White and Masset 2003). For richer countries where economies of scale are likely to be greater (given the lower proportion of food in the household budget, and greater coverage of utilities), the coefficient will be higher; e.g., Wagstaff (2000) cites a study of OECD countries that implied the coefficient of scale was equal to around 0.6 (Buhmann and others 1988).

Deciding on an overall scale coefficient may be problematic, given that different assets are likely to have different scale economies, which may also depend on the particular use the index is trying to capture. However, it is possible to adjust some components of the asset index selectively.

Most studies using an asset index (e.g., Gwatkin and others 2000) do not indicate that they adjust the asset index for household size, which means that, implicitly, the scale coefficient is assumed to equal 0 (i.e., it is assumed that adding another person to the household does not alter the value of the index, nor the weights attached to specific variables in the index). Wagstaff (2000) takes an intermediate stance by setting the scale coefficient as equal to 0.5 for expenditure data—in his estimation for nine developing countries, assuming zero economies of scale tended to reduce socioeconomic inequalities in mortality. Filmer and Pritchett (1999) and Sahn and Stifel (2000) report that results are insensitive to different assumptions about scale coefficients.

Constructing a Wealth Index for Bangladesh

The preceding sections examined major conceptual and practical issues regarding the generation of an asset index. The aim is to provide recommendations for generating an asset index using DHS survey data for Bangladesh, to be used subsequently in bivariate and multivariate

analyses of child mortality and nutritional outcomes. First, given the importance of separating assets that are likely to be direct determinants of health outcomes from those that have indirect effects, by representing “wealth,” the paper recommends that direct factors should be excluded, and estimated separately from wealth. Direct determinants of child nutrition and survival are drinking water source, toilet facility, and floor material of the household. This constrains which assets can be included in the index, and therefore the possibilities for differentiating between households, especially given that other items that, while having arguably less direct effects on health and nutrition, are representative of location, such as electricity, and therefore should be excluded. We are left with household durables—ownership of almirah (wardrobe), clock, cot, bicycle, radio, table, TV—and housing quality as measured by wall and roof material.¹⁵

A wealth index is constructed based on weightings that are both intuitive and in part determined by the variation in the data. The structure of the index follows from Young Lives (2002), in that it is the average of household durables ownership and housing quality, where total durables owned and housing quality are scaled comparably. However, unlike Young Lives (2002), which calculates all weights arbitrarily, the weightings in the index for Bangladesh for durable goods are partly chosen by the variation in the data—specifically, individual weights for each durable owned is set equal to $(1-P)$ where P is the proportion of households owning the item; total ownership of durables is rescaled accordingly. This scheme allows much greater differentiation between households, as compared, for example, to a scheme based on equal weightings of durables.¹⁶ Weightings for housing quality are determined intuitively, though the weightings chosen are not incontrovertible, since choice of building material is partly determined by location.¹⁷ Each household’s asset score comprises the mean of:

- (1) Household durables: weighted sum of dummy variables for seven goods, scaled to range between 0 and 1.¹⁸

- (2) Housing quality, which is itself the mean of two components:

- Wall material dummy (weight): natural/other (0), wood (0.25), tin (0.5), brick (1).
- Roof material dummy (weight): natural/other (0), tin (0.5), cement/tile (1).

Since the weightings are time insensitive, this enables comparisons between households over time, to determine whether wealth poverty is increasing or decreasing over time. Due to the arguably strong public good aspects of the non-durables items, and given that the DHS do not collect data on number of items owned of each household durable, no adjustment is made for household size or composition—the economies of scale in the household are assumed perfect.

Table C2.4 shows the actual weighting ascribed to each variable in the asset index, as well as the raw Eigenvectors calculated using principal components analysis. In order to make comparisons between each, the scores are rescaled to lie in the interval (0, 1). PCA ranks housing

TABLE C 2 . 4		Weights for Bangladesh DHS 1993/94, 1996/97, and 1999/2000		
Asset variable	Scores for intuitive index	Score for PCA index	Rescaled scores	
			Intuitive	PCA
Household durable goods				
Radio	0.081	0.273	0.326	0.887
TV	0.101	0.314	0.403	0.948
Bicycle	0.095	0.176	0.379	0.743
Wardrobe (Almirah)	0.086	0.334	0.344	0.978
Table	0.050	0.326	0.201	0.966
Clock	0.062	0.349	0.247	1.000
Cot	0.025	0.259	0.100	0.866
Housing quality				
Jute/bamboo/mud wall	0.000	−0.325	0.000	0.000
Wooden wall	0.063	0.024	0.250	0.517
Tin wall	0.125	0.137	0.500	0.685
Brick wall	0.250	0.316	1.000	0.951
Other wall	0.000	−0.061	0.000	0.392
Bamboo/thatch roof	0.000	−0.284	0.000	0.061
Tin roof	0.125	0.149	0.500	0.703
Cement/tile roof	0.250	0.235	1.000	0.830
Other roof	0.000	−0.040	0.000	0.423

quality similarly to the intuitive index, with the exception of “other” categories. The main difference between the indices is that, as expected, PCA generates scores that give greater weight to durable goods. However, the indices are strongly correlated: the Spearman’s rank correlation between the indices is equal to 0.99, while the correlation coefficient is 0.98.

The index enables a fairly consistent ranking of households, meaning that they can be identified into nearly equal wealth quintiles. Table C2.5 shows the distribution of households by quintile for each year. As with remaining calculations, survey weights have been rescaled so that households have an equal chance of being in each year. The results indicate that household wealth is improving over time on average and for all quintiles. The Bangladesh PRSP (Bangladesh/IMF 2003) indicates that consumption poverty declined from 59 to 50 percent between 1991/92 and 2000. A poverty line is, therefore, chosen to define 50 percent of households as poor based on their wealth in 1999/2000, which is equal to a value of the wealth index of 0.30. Table C2.6 shows poverty estimates and standard errors for “poverty” as defined here, using the poverty index due to Foster and others (1984).¹⁹ Wealth poverty is estimated to be declining much faster than income poverty. Asset ownership among the poorest is also growing fastest, so that overall inequality is decreasing; in contrast, income inequality, as measured by the Gini index, rose considerably during the 1990s from about 25 to 30 percent (Bangladesh/IMF 2003). This fall in

inequality in the asset index makes intuitive sense since households attain assets as they get richer (and once attained, assets are kept), whereas the index is made up of a limited range of assets, and is unable to distinguish multiple ownership of the same asset. Table C2.7 breaks down the change in the asset index over time into shares made up by changes in ownership of individual assets, where each asset is weighted with its asset score, indicating that more than three-quarters of the change in the average value of the index comprises increases in ownership of TVs, clocks, tin walls and roofs, brick walls and cement roofs.

Linking Wealth and Expenditure

Wealth indices are used as a measure of economic well-being, partly as DHS collects data on assets but not expenditure. But economic analysis may tell us the income/growth effect of an intervention. Is a 5 percent increase in income the same as a 5 percent increase in the wealth index?

Thinking about how the wealth index is constructed suggests it is probably not so. Wealth indices are constructed on variables such as whether the household owns a radio and a bicycle (or sometimes the number of such items). Once households receive a certain level of income they will own these things, but not buy more as income rises. The wealth index is thus unlikely to pick up increases in income above a certain level. At the very least this argument suggests that the elasticity of the wealth index with respect to household expenditure should be

TABLE C2.5

Shares of Population in Wealth Quintile and Mean Value of Index by Year

Quintile	1993/94		1996/97		1999/2000	
	Percent of population	Mean value of wealth index	Percent of population	Mean value of wealth index	Percent of population	Mean value of wealth index
Q1	24	0.010	22	0.022	20	0.058
Q2	17	0.100	19	0.131	20	0.185
Q3	19	0.179	20	0.227	20	0.304
Q4	21	0.323	20	0.380	21	0.462
Q5	19	0.641	20	0.693	19	0.752
Total	100	0.244	100	0.287	100	0.350

Note: Households are ordered into quintiles separately by year.

less than one, and probably that it will be lower at higher levels.

This issue was examined using data from the Bangladesh Household Income and Expenditure Survey (HIES) of 2000-01. HIES and DHS questionnaires do not cover the same household assets, but are very similar. The variables used by DHS, and the corresponding variables used from HIES are in table C2.8.

Table C2.9 shows the expenditure elasticity of the wealth index, obtained from a regression of the log of the wealth index on the log of per capita household expenditure. The estimated elasticity is 0.8. Two other models are presented in the second and third column. The first includes a quadratic term of the log of per capita expenditure. The second includes an intercept and slope dummy for the top 25 percent of the per capita expenditure distribution. As expected, the expenditure elasticity is decreasing at higher levels of household expenditure.

TABLE C2.6		Estimates of Poverty and Inequality		
	Headcount index	Poverty gap index	Squared poverty gap	Gini index
1993-94	0.690 (0.011)	0.430 (0.011)	0.335 (0.011)	0.515 (0.007)
1996-97	0.616 (0.012)	0.353 (0.010)	0.261 (0.010)	0.467 (0.006)
1999-2000	0.501 (0.013)	0.247 (0.010)	0.169 (0.008)	0.400 (0.007)

Note: Standard errors in parantheses.

TABLE C2.7		Contribution to Change in Asset Index Over Time		
Asset variable	Percent owning asset			Percentage contribution to change in wealth index 1993-99
	1993	1996	1999	
Radio	25	32	32	5.3
TV	7	11	17	9.5
Bicycle	16	19	20	3.6
Almirah (wardrobe)	22	27	26	3.2
Table	49	55	63	6.6
Clock	35	46	55	11.7
Cot	72	78	84	2.8
Jute/bamboo/mud wall	73	71	61	0.0
Wooden wall	3	2	3	0.0
Tin wall	9	12	19	11.8
Brick wall	11	12	17	14.2
Other wall	4	3	0	0.0
Bamboo/thatch roof	37	29	20	0.0
Tin roof	53	59	70	20.0
Cement/tile roof	5	6	9	9.4
Other roof	6	6	0	0.0

TABLE C2.8

**Asset Variables
Collected by DHS
and Similar
Variable of HIES**

<i>DHS</i>	<i>HIES</i>
Radio	Radio
TV	TV
Bicycle	Bicycle
Almirah (wardrobe)	Drawing room furniture
Table	Dining room furniture
Clock	Clock
Bed	Bedroom furniture
Wall material	Wall material
Finished	Brick/cement
Rudimentary	Cl sheet/tile/wood
Natural	Hemp/hay/bamboo
Roof material	Roof material
Finished	Cement
Rudimentary	Cl sheet/tile/wood
Natural	Hemp/hay/bamboo

TABLE C2.9

**Expenditure
Elasticities of the
Wealth Index
(standard errors
in brackets)**

Variable	Double arithmetic	Log- quadratic	Top intercept and slope
Log of expenditure	0.81*** (0.02)	2.21*** (0.27)	0.87*** (0.04)
Log of expenditure squared		-0.10*** (0.02)	
Intercept of top 25%			2.53*** (0.38)
Slope of top 25%			-0.35*** (0.05)
Constant	-6.94*** (0.14)	-11.7*** (0.95)	-7.34*** (0.24)
Observations	6887.81***	6887.77***	6887.53***
R-squared	0.30	0.31	0.31

FIGURE C2.1

**Plot of the Predicted
Values Against per
Capita Expenditure
from the 3 Models
Above**




TABLE C2.10

Asset Score for Bangladesh DHS 1996-97, Generated by Principal Components Analysis as Reported in Gwatkin and others (2000)

Asset variable	Asset factor scores	Household score if:		Ownership (1=yes, 0=no)		Asset score	
		Has asset	Does not have asset	Household A	Household B	Household A	Household B
Has electricity	0.118	0.214	-0.065	1	0	0.214	-0.065
Has radio	0.071	0.104	-0.049	1	1	0.104	0.104
Has television	0.125	0.347	-0.045	1	0	0.347	-0.045
Has bicycle	0.032	0.065	-0.016	1	1	0.065	0.065
If household works own or family's agric. land	-0.022	-0.042	0.012	0	1	0.012	-0.042
If piped drinking water in residence	0.104	0.526	-0.021	0	0	-0.021	-0.021
If piped drinking water outside residence	0.020	0.179	-0.002	1	0	0.179	-0.002
If has tubewell for drinking water	-0.063	-0.022	0.186	0	0	0.186	0.186
If has a surface well	-0.009	-0.066	0.001	0	1	0.001	-0.066
If rain for drinking water	-0.001	-0.023	0.000	0	0	0.000	0.000
If uses river, canal or surface water for drinking	-0.009	-0.049	0.002	0	0	0.002	0.002
Other source of drinking water	0.000	0.005	0.000	0	0	0.000	0.000
If uses septic tank or toilet	0.127	0.370	-0.043	1	0	0.370	-0.043
If has pit latrine	-0.016	-0.041	0.006	0	1	0.006	-0.041
If uses a water-sealed or slab latrine	0.017	0.033	-0.009	0	0	-0.009	-0.009
If has open latrine	-0.033	-0.059	0.019	0	0	0.019	0.019
If uses a hanging latrine	-0.012	-0.054	0.003	0	0	0.003	0.003
If uses bush, field as latrine	-0.054	-0.092	0.031	0	0	0.031	0.031
If has other type of latrine	-0.001	-0.031	0.000	0	0	0.000	0.000
If has earth or bamboo as principal floor in dwelling	-0.152	-0.057	0.401	0	0	0.401	0.401
If has wood, plank principal floor in dwelling	0.010	0.146	-0.001	0	1	-0.001	0.146
If has cement principal floor	0.153	0.415	-0.056	1	0	0.415	-0.056
If has other type of flooring	0.003	0.135	0.000	0	0	0.000	0.000
If has cane, palm, trunks for walls	-0.108	-0.074	0.158	0	0	0.158	0.158
If has rudimentary walls	-0.004	-0.021	0.001	0	1	0.001	-0.021
If has tin for walls	0.012	0.034	-0.004	0	0	-0.004	-0.004
If has bricks, cement blocks, concrete walls	0.143	0.361	-0.057	1	0	0.361	-0.057
If has other material for walls	-0.010	-0.054	0.002	0	0	0.002	0.002
If has natural material roofing	-0.061	-0.095	0.039	0	0	0.039	0.039
If has rudimentary roofing	0.003	0.003	-0.004	0	1	-0.004	0.003
If has finished roof	0.125	0.498	-0.031	1	0	0.498	-0.031
If has other roofing	-0.014	0.053	0.004	0	0	0.004	0.004
Total asset score						3.379	0.659

TABLE C2.11

Studies Using Assets to Measure SES

Author	Description of study	Country	Asset/income measure	Asset type			
				Household consumer durables	Household dwelling characteristics	Amenities	Non-housing (production) items
Hill and others (1996)	Child mortality, DHS	Kenya	Asset index (PCA), generating wealth quintiles	Radio, TV, refrigerator, bicycle, motorbike, car	Quality of floor, roof	Electricity	
Filmer and Pritchett (1998)	Educational enrollment, NFHS	Indian states	Asset index (PCA), generating wealth quintiles	Clock/watch, bicycle, radio, TV, sewing machine, refrigerator, car, motorcycle	Number of rooms, building materials, cooking source	Toilet facilities, drinking water, electric lighting	HH owns more than 6 acres of land
Filmer and Pritchett (1999)	Educational attainment, DHS	33 countries across the world, including Bangladesh	Asset index (PCA), generating wealth quintiles	Clock/watch, bicycle, radio, TV, sewing machine, refrigerator, car, motorcycle	Number of rooms, building materials, cooking source	Toilet facilities, drinking water, electric lighting	HH owns more than 6 acres of land
Gage, Sommerfelt, and Piani (1996)	Immunization and diarrhea, DHS	11 Sub-Saharan African countries	Asset index (asset dummies summed—equal weightings)	Radio, motorcycle, car	Non-mud floor	Some toilet facility, piped drinking water, electricity	
Gwatkin and others (2000)	Health, nutrition and population, DHS	Bangladesh	Asset index (PCA), generating wealth quintiles	Radio, TV, bicycle	Floor material, wall material, roof material	Toilet facilities, drinking water, electricity	HH works own or family's agric. land
Kishor and Parasuraman (1998)	Infant and child mortality, NFHS	India	HH asset index (ownership dummies summed—weights chosen apparently subjectively), toilet and water facilities index (ranges from 0-2)	Asset (weight in parentheses): clock (1), sewing machine (2), sofa (2), fan (2), radio (2), bicycle (2), refrigerator (3), TV (3), scooter (3), car (4)		No toilet or water, either toilet or water, both toilet and water.	
Madise, Matthews, and Margetts (1999)	Nutrition, DHS	6 in Sub-Saharan Africa	Asset dummies: variables for hh items (at least one item, two or more items) and community items entered separately	'Modern hh items': working radio, TV, bicycle, motorcycle, car	Floor material, wall material	Electricity, pit/bucket toilet, flush toilet	
Masset and White (2002)	Infant and child mortality, NFHS	India	HH asset index—non-weighted, scaled to one	Radio, TV set, refrigerator, bicycle, motorcycle, car, land, live-stock, clock and sewing machine			
Sahn and Stifel (2000)	'Poverty' across countries and time	11 in Sub-Saharan Africa	Asset index (FA)	Radio, TV, refrigerator, bicycle, car, motorcycle	Floor quality	Toilet facilities, drinking water	Years of education of household head (or indicator variable where years education is not available)
Sastry (1996)	Child mortality, DHS variant	Brazil	Asset dummies measured at community level (proportions per population) (hh level SES measured using hh income)			Water supply, sanitation, electricity, garbage collection/public cleaning service (also health facilities and education per population and number of TV stations)	
Wagstaff and Watanabe (2002)	Nutrition, LSMS	19 countries	Income, asset index (PCA)	Radio, refrigerator, TV, motorcycle	Rooms per person, floor quality	Water supply, sanitation	

APPENDIX C.3: Socioeconomic Inequality in Mortality in Bangladesh During the 1990s

This annex presents estimates of intertemporal movements in socioeconomic inequality in mortality among under-fives in Bangladesh. Socioeconomic status is measured using the wealth index, which is calculated as the rescaled weighted sum of household ownership durables and indicators of housing quality (wall and roof materials). Mortality estimates are calculated from three rounds of the Bangladesh DHS (1993/94, 1996/97, and 1999/2000); data are pooled across surveys, so estimates become more precise as we move back in time, as they are based on a larger sample. Period mortality rates are estimated using the synthetic cohorts probabilities method (Rutstein 1984).

Examining health outcomes by income poverty, to which ill-health is often closely linked, it is possible to decompose changes in mortality by poor and non-poor. Total mortality in each period is equal to the weighted sum of mortality among the poor and mortality among the non-poor:

$$MR_t = p_t \cdot MR_t^p + (1 - p_t) \cdot MR_t^n \quad (1)$$

where MR_t is the mortality rate period t , p , and n superscripts denote poor and non-poor, and p equals the share of the population living in poverty. Table C3.1 provides some indication that the improvement in survival has been achieved among both “poor” and “non-poor.”

The poverty line is defined as the value of wealth that divides the population of households in half in the 1999/2000 survey round, which corresponds to the national estimate of expenditure poverty in 2000 of 50 percent (Government of Bangladesh 2002). As shown in the table, under-five mortality fell slightly faster for the poor than the non-poor, with the largest difference being for reductions of neonatal mortality, and in fact reversed for postnatal.

In order to examine changes over time between periods 1 and 2, equation (1) can be decomposed into the change in mortality rates among the poor, the change in mortality among the rich, and the change in poverty:¹

$$\Delta MR = \Delta MR^p \cdot p_2 + \Delta MR^n \cdot (1 - p_2) + \Delta p \cdot (MR_1^p - MR_1^n) \quad (2)$$

That is, the change in mortality is the weighted sum of the change in mortality for the poor and non-poor, plus the product of proportion of people moving out of poverty times the mortality differential for poor and non-poor. Table C3.2 presents the results of the decomposition,² indicating that, for each age group, the reduction in mortality over time has been largely due to the reduction in mortality among the poor. In addition, virtually all the change is explained by reduction of mortality within groups, and very little as a result of households moving

TABLE C3.1 Estimates of Mortality (per 1,000) by Poverty Status

	Neonatal		Postnatal		Child		Under-five	
	Poor	Non-poor	Poor	Non-poor	Poor	Non-poor	Poor	Non-poor
1980–84	101	70	55	41	91	42	233	148
1985–89	71	60	52	29	76	37	190	123
1990–94	67	50	38	33	60	28	158	109
1995–99	44	38	33	19	38	19	112	75
Percentage change	-56.4	-45.7	-40.0	-53.7	-58.2	-54.8	-51.9	-49.3

TABLE C3.2
**Mortality Rate
Decomposition by
Poverty Status**

	Decomposition $\Delta(\% \text{ of total } \Delta MR)$			ΔMR
	$\Delta MR \text{ poor}$	$\Delta MR \text{ non-poor}$	$\Delta \text{Poverty}$	
Neonatal	64.9	32.2	2.9	100.0
Postnatal	53.1	44.8	2.1	100.0
Child	69.4	27.7	2.9	100.0
Under five	63.4	34.2	2.4	100.0

out of poverty. These findings support the argument made in Annex B that factors other than income growth are the main drivers of mortality reduction in Bangladesh.³

The preceding analysis has examined some distributional characteristics of health status among children, but only by looking at two groups, “poor” and “non-poor,” and therefore abstracting from a large amount of distributional information. It is possible to utilize information on the entire wealth distribution by using concentration curves and concentration indices. There are various ways of measuring socioeconomic inequalities in health (and other welfare outcomes) (see Mackenbach and Kunst 1997), though a common and easily interpreted measure is the concentration index, which is calculated from the concentration curve in the same way as the Gini coefficient is obtained from the Lorenz curve. The concentration curve is similar to the Lorenz curve, except that where the Lorenz curve ranks observations by the variable in which inequality is being measured, the concentration curve ranks them according to another variable (e.g., to generate socioeconomic inequalities in health, the concentration index ranks households by SES and calculates the resultant inequality in health status. Unlike Gini coefficient, the concentration index, C , can take negative values: $C < 0$ indicates poorer socioeconomic groups have worse health status; $C = 0$ indicates perfect equality in health status by socioeconomic status; $C > 0$ indicates poorer socioeconomic groups have better health status.

The concentration index can be calculated from the following formula (Kakwani 1980):

$$C = 2 \operatorname{cov}(y_i, R_i) / \mu \quad (3)$$

where μ is the mean for variable y whose inequality is being measured and cov is the covariance between each individual i 's value of y and the individual's fractional rank in the wealth distribution, R_i . Unweighted standard errors are calculated from:






$$\operatorname{var}(C) = 1/n \left[\sum_i a_i^2 - (1 + C)^2 \right] \quad (4)$$

$$\text{where } a_i = y_i / \mu (2R_i - 1 - C) + 2 - q_{i-1} - q_i \quad (5)$$

$$\text{and } q_i = 1/(\mu n) \sum_i y_i \quad (6)$$

(Kakwani and others 1997). Examination of concentration curves for the 1990s for under-five, infant, and child mortality indicates that mortality is more concentrated among the poor during childhood than infancy (figure C3.1). This is expected since socioeconomic characteristics determine survival to a greater extent as the child grows older and biological factors become less important. Concentration curves have also, generally, moved toward the line of equality, with much of the reduction in inequality arising from inward movements of the lower half of the distribution of wealth, particularly for neonatal infants, showing that improvements in survival prospects have been faster for newborns of poorer people. Table C3.3 presents concentration indices and t-statistics, confirming these trends. Inequality in under-five mortality has decreased over the period, mainly caused by a reduction in mortality inequality among children and among neonates (table C3.4). However, differences between concentration indices over time are not significant at reasonable levels, with the exception of neonates, for whom the concentration index fell significantly between 1980-84 and 1995-99.⁴

Examining socioeconomic inequalities in mortality risk by child's gender, the conclusion emerges that inequalities between rich and poor are smaller for boys than for girls, and the inequalities among boys have also been declining faster (table C3.5). By differentiating be-

FIGURE C3.1**Concentration Curves for Infants, Children, and Under-Fives, all Children and by Gender***Concentration curves for under-five mortality**Concentration curves for neonatal mortality**Concentration curves for child mortality
(Children aged 12-59 mos.)**Concentration curves for postnatal mortality**Concentration curves for infant mortality*

Sources: Adapted data from WDI; graph, bars from DHS.

TABLE C3.3**Concentration Indices for Infants, Children, and Under-Fives**

	Infant	t stat.	Child	t stat.	Under five	t stat.
1980–84	–0.10	–9.72	–0.22	–14.56	–0.15	–17.38
1985–89	–0.08	–7.90	–0.21	–13.83	–0.14	–16.25
1990–94	–0.09	–7.29	–0.21	–11.89	–0.13	–13.25
1995–99	–0.08	–4.52	–0.20	–6.93	–0.12	–7.45

TABLE C3.4**Concentration Indices for Neo- and Postnates**

	Neonatal	t stat.	Postnatal	t stat.
1980–84	–0.08	–6.24	–0.12	–7.12
1985–89	–0.06	–4.89	–0.11	–5.95
1990–94	–0.08	–5.10	–0.11	–5.41
1995–99	–0.04	–2.82	–0.14	–3.88

tween infants and children (table C3.6) and then, among infants, between neonates and postnates (table C3.7), it becomes clear that all of the apparent decline in mortality inequality between “poor” and “non-poor” girls has occurred for the neonatal period—postnatal and child mortality inequality has remained constant or deteriorated among girls. However, tests show that differences between girl concentration indices over time are not statistically significant, with the exception of girl neonates for whom the reduction in inequality in mortality risk between 1980–84 and 1995–99 is marginally significant at the 10 percent level. For boys,

TABLE C3.5**Concentration Indices for Under-Five Mortality by Child’s Gender**

	Male	t stat.	Female	t stat.
1980–84	–0.14	–11.67	–0.16	–12.97
1985–89	–0.13	–11.03	–0.14	–11.81
1990–94	–0.13	–8.99	–0.14	–9.89
1995–99	–0.09	–4.16	–0.14	–6.47

much of the decline in socioeconomic inequality in mortality occurred for children and, to a lesser extent, for neonates, while inequalities in postnatal mortality widened during the mid-1990s, though none of these differences are statistically significant.

It is possible to decompose socioeconomic inequalities in mortality risk into inequalities in the determinants of mortality; similarly changes in inequalities can be decomposed into changes in inequality of the determinants.

Following Wagstaff (2000), the concentration index can be expressed as the weighted sum of the concentration indices c_i of the n variables determining mortality:

$$C^* = y_1 c_1 + y_2 c_2 + \dots + y_n c_n \quad (7)$$

where the weights are calculated by multiplying the coefficient estimate by the mean of that variable and dividing by the linear prediction of the regression model at the means of the data. Note that what is really being examined in this analysis is socioeconomic inequality in the underlying latent variable in the case of probit regressions (for neonatal and postnatal mortality) and the log of the hazard ratio in the case of the survival model (for child mortality). The change in the concentration index over time t can also be decomposed into the proportional contribution of the change in concentration indices for each variable i :

$$\Delta = y(\Delta c_i)/C^* \quad (8)$$

Table C3.8 presents a summary of the decomposition of the changes in concentration indices over time for determinants of mortality used in subsequent regressions, with the full results, including concentration indices for each variable, given in table C3.9. Socioeconomic inequalities in determinants of mortality have generally declined over time; this is true for factors such as wealth, education attainment, and maternal mobility. It is not true for the measures of health: there has been no change for antenatal care and a worsening for vitamin A and having no vaccination. For neonatal mortality, fertility-related changes, such as birth order have played

TABLE C3.6		Concentration Indices by Child's Gender						
	Infant				Child			
	Male	t stat.	Female	t stat.	Male	t stat.	Female	t stat.
1980–84	–0.09	–7.00	–0.11	–6.63	–0.20	–8.69	–0.23	–11.51
1985–89	–0.10	–6.77	–0.05	–4.16	–0.18	–7.95	–0.23	–11.44
1990–94	–0.10	–5.63	–0.07	–4.55	–0.18	–6.90	–0.22	–9.89
1995–99	–0.08	–3.12	–0.08	–3.29	–0.15	–3.52	–0.24	–6.16

TABLE C3.7		Concentration Indices by Child's Gender						
	Neonatal				Postnatal			
	Male	t stat.	Female	t stat.	Male	t stat.	Female	t stat.
1980–84	–0.09	–4.83	–0.08	–3.96	–0.10	–4.83	–0.15	–5.23
1985–89	–0.07	–4.14	–0.04	–2.60	–0.14	–5.27	–0.07	–3.07
1990–94	–0.07	–3.74	–0.08	–3.43	–0.15	–4.48	–0.08	–3.13
1995–99	–0.06	–2.58	–0.03	–1.52	–0.12	–2.03	–0.17	–3.71

TABLE C3.8		Summary of Contribution to Changes in Concentration Index		
	Neonatal	Postnatal	Child	
Wealth	0.012	0.001	−0.164	
Education	−0.019	0.002	−0.085	
Fertility related	−0.149	−0.017	0.023	
Electricity	−0.069	0.000	0.005	
Water and sanitation	−0.015	0.000	0.003	
Gender related	−0.014	−0.001	0.061	
Locational	−0.003	−0.004	0.010	
Other	0.121	0.001	0.060	
Total	−0.136	−0.018	−0.087	

the largest role, followed by a reduction in the inequality in access to electricity. The decompositions for postnatal mortality are unrevealing, due to the small estimated change in postnatal mortality during the period. Finally, for children

the most important single variable contributing to reduction in socioeconomic inequality in mortality is wealth, followed by maternal education, which accounts for just over half the change accounted for by wealth.

TABLE C3.9		Decomposition of Changes in Socioeconomic Inequality in Mortality by Age Group			
	Early 1990s	Late 1990s	Neonatal	Postnatal	Child
Wealth	0.514	0.412	0.012	0.001	-0.164
Electricity	0.593	0.478	-0.069	0.000	0.005
Water & sanitation	0.899	0.846	-0.015	0.000	0.003
Mother primary ed	0.030	-0.049	-0.011	0.000	-0.048
Mother finished primary	0.313	0.182	-0.027	0.000	-0.011
Mother secondary ed	0.552	0.425	0.025	0.001	-0.025
Mother finished second or higher	0.763	0.741	-0.006	0.001	-0.001
Contraception	0.041	0.037	-0.020	-0.001	-0.003
Multiple birth	0.112	-0.087	0.102	0.000	
Age at birth	-0.002	-0.003	-0.027	-0.001	-0.004
Age at birth sq	-0.004	-0.008	0.042		
Prev child died	-0.126	-0.117	-0.005	0.000	-0.001
Prec interval <15	0.004	-0.064	0.043	0.000	
Prev child died x prec higher interval	-0.011	-0.128	-0.028	0.000	
Birth order	-0.034	-0.058	-0.398	-0.009	0.030
Birth order sq	-0.048	-0.097	0.297	-0.006	
Female child	-0.008	-0.005	0.004	0.000	0.004
Female x birth order	-0.036	-0.066		0.000	0.052
Female head of hh	0.003	0.004	0.002	0.000	0.000
Mother's mobility	0.026	0.017	-0.015	0.000	-0.001
Mother divorced	-0.258	-0.083	0.001	0.000	-0.004
Mother remarried	-0.009	-0.016	-0.006	-0.001	0.010
Antenatal visits	0.304	0.304	-0.001		
Chittagong	0.010	-0.026	-0.011	0.000	-0.017
Dhaka	0.058	0.069	0.003	0.000	0.007
Khulna	-0.046	0.104	0.035	0.001	0.059
Rajshahi	-0.062	-0.107	-0.009	-0.001	-0.033
Rural	-0.048	-0.073	-0.021	-0.004	-0.006
Born In Feb-Mar x rural	-0.082	-0.065	0.004	0.000	
Born In Apr-May x rural	-0.029	-0.083	-0.029	0.000	
Born In Jun-Jul x rural	-0.064	-0.037	0.003	0.000	
Born In Aug-Sep x rural	-0.069	-0.076	-0.003	0.000	
Born In Oct-Nov x rural	-0.053	-0.099	-0.008	0.000	
Mean_vitamin A	0.005	0.076		0.001	0.037
Mean_no_vaccine	-0.035	-0.086			0.024

ANNEX D. CHILD MALNUTRITION DURING THE 1990S

Bangladesh has made good progress in reducing rates of malnutrition among children, particularly in the 1990s, when malnutrition is estimated to have fallen by almost one-third (Deolalikar 2002; Annex A). However, malnutrition remains a big problem. Stunting was estimated at nearly 50 percent in 2000 (see figure D.1), among the highest rates in the world. Of countries with available data, only a dozen others, mainly in South Asia and Sub-Saharan Africa, have worse stunting rates (UNICEF 2002, table 2). Similarly, underweight prevalence in Bangladesh, though falling, remains one of the world's highest. Bangladesh seems to perform much better compared with other countries in incidence of wasting, though it should be borne in mind that, due to seasonal variation in nutrition, it is difficult to make sound comparisons of short-term indicators across surveys that may well have been conducted at different times of year.

This paper examines the determinants of nutritional status among children, in order to ascertain the contribution of various factors to nutrition differentials across children and improvements in mean outcomes over time. Bangladesh experiences higher malnutrition than its income level alone would predict (see Annex B), and discrimination against girls in nutrition and health care has been well-documented (for a relatively early example, see Chen and others 1981).

Modeling Child Nutritional Status

Nutritional status is measured anthropometrically,¹ using three indicators: height for age (stunting), weight for height (wasting), and weight for age (underweight). Height for age is an indicator of long-term malnutrition because it


measures nutrition cumulatively over time. Weight for height is sensitive to short-term fluctuations in nutritional state. Weight for age is a composite measure of weight for height and height for age; it is criticized due to its inability to distinguish children who are truly thin from those who weigh less because they are shorter. However, it has been shown to be more closely related to mortality risk in Bangladesh than other indicators (Chen and others 1980).²

Child nutritional status is determined by comparing the anthropometric indicator for each child in the survey population to the expected measurement for a child of the same age/height and sex from a healthy reference population.³ This comparison can be done in a variety of ways, the most effective of which is to normalize the anthropometric measure by calculating z scores:

$$z_{i(y,g)} = \frac{x_i - \mu_{(y,g)}}{\sigma_{(y,g)}}$$

where x_i indicates the anthropometric measure for individual i and μ and σ are the respective median and standard deviation of the anthropometric measure for the appropriate age or height y and gender g in the reference population.⁴

The basic model of child nutritional status is due to UNICEF (1990), in which nutrition depends proximately on dietary intake and health status. Thus, high rates of child malnutrition are likely to be closely linked to the underlying household food insecurity and the exposure to infectious disease such as diarrhea, due to poor sanitary environment and inadequate medical treatment. A third underlying factor is the quantity and quality of child care, which determines

FIGURE D.1 Malnutrition Prevalence in Bangladesh

Source: Calculated from DHS data.

how effectively income is converted into nutrition and the share allocated to children (and, importantly, which children receive the most), as well as the healthiness of the home and community environments. Note that these three underlying factors may be complements or substitutes in the production of child health.

The meta-analysis conducted as a background paper for this study (Charmarbagwala and others 2004) provides a summary of the literature modeling nutritional status, as indicated by height for age z score. In multivariate analysis, nutritional status can be modeled as a function of child-specific factors, d , household-specific factors, b , and community factors, e :

$$z = n(d, b, e).$$

Child-specific factors include demographic variables such as child's age, gender, birth order and birth interval, and behavioral factors such as breastfeeding and immunization. The relationship between age and nutritional status depends on whether the anthropometric indicator measures short-term or long-term malnutrition. For

short-term measures, age should exert a non-linear concave impact on nutritional status: nutrition deteriorates in the period in which children are most susceptible to disease—from the onset of weaning (which should be at age 6 months) until age 24 months—and improves thereafter. For long-term measures the relationship between age and malnutrition is also likely to be concave, though because stunting is often permanent, the function is unlikely to fall after 24 months of age. Gender-based discrimination in South Asia means that the coefficient on a female dummy may be negative; girls who have older sisters or who are competing with the first-born male are particularly likely to be discriminated against (Croll 2001).⁵ Surprisingly, the meta-analysis found that boy children were significantly more likely to be malnourished than girls (Charmarbagwala and others 2004). However, this finding may be due to mortality selectivity, which was not allowed for in previous studies (see below).

More generally, children of high birth order (born latest) and children born closely together

are more likely to have lower nutritional states, due to socioeconomic factors (resource competition) and biological factors (e.g., physically depleted mothers may give birth to low birth-weight babies and may be unable to breastfeed). Mozumder et al. (2000) find that short subsequent birth interval impacts negatively on weight for age, but no statistically significant impact of preceding interval, which they attribute to the likelihood that “the new infant holds an advantageous position with the mother, compared with any older siblings, because of breastfeeding” (p. 295). One issue concerning use of household composition and demographic variables is that they may be endogenous to decisions on child health, leading to biased regression estimates. As Horton (1986) argues, parents are likely to make joint decisions on child quantity and child quality (demonstrated, for example, by nutritional status). Regression estimation should account for this source of bias, e.g., through use of instruments or by dropping endogenous demographic variables to assess robustness of the other coefficient estimates.

Child immunization is an important variable to control for—vaccination reduces chances of contracting debilitating disease. A mother’s decision to immunize her child depends partly on availability of health services and income, but also her preference regarding modern medical care, which is unobservable.⁶ Due to unobservable heterogeneity, behavioral variables such as immunization may be correlated with the error term in the regression equation, leading to biased regression estimates (Rosenzweig and Schultz 1983; Thomas and others 1991; Alderman and Garcia 1994). Consistent estimation requires use of instrumental variables (the “production function” approach).

In addition to the above, determinants of long-term malnutrition (height for age) should control for genetic health endowment and birth weight, which can be an important determinant of ill-health and height throughout childhood and onwards. Commonly used indicators of health endowment are (log of) mother’s height and, due to seasonal variation in food availability, month of birth may be a good proxy for birth

weight; DHS round 2000 also provides data on mother’s recollection of low birth weight.

Household-specific factors include parental or family resources such as income, household size and composition, parental education, mother’s age and mobility, and sex of the household head; these are proxies for food security and the quality and quantity of care provided to children.

Income—broadly defined to include imputed own production—is a key determinant of household nutritional intake. In the health production function literature, income is considered jointly determined with nutrition and health, leading to biased estimates of the coefficient on the income variable in standard regression analysis (see Annex C). The extent to which current income and child health are endogenously determined can be questioned for extreme cases such as long-term malnutrition (stunting) and mortality (see Chamarbagwala and others 2004). However, estimation techniques usually instrument for income, a suitable instrumental variable being wealth.⁷ The meta-analysis found income/wealth to be strongly correlated with child nutrition, with the clear majority of studies finding a significantly positive effect (see Chamarbagwala and others 2004).

Food availability is highly seasonally dependent in rural areas of Bangladesh (see Annex J). Models of short-term nutritional status should therefore include the month that measurement was taken (interacted with a rural sector dummy). The wet season in July–October, which occurs before the main rice harvest, *aman*, is a critical time of year for child health, due to greater prevalence of water-borne disease (diarrhea, malaria) and food scarcity (Muhuri 1996; Annex I). DHS data were collected between November and March, limiting the possibilities to explore seasonal variation in short-term malnutrition to this 5-month period. DHS 2000 round does, however, provide data on the household’s (self-reported) food availability during the year.

Household size is likely to be positively correlated with child nutrition, reflecting availability of replacement caretakers, including older children and grandparents (people who may other-

wise be considered economic “dependents”). The meta-analysis found evidence for a significantly positive effect of household size on nutrition, and a significantly negative effect of presence of young children in the household (Charmarbagwala et al. 2004). Estimation should account for potential endogeneity of household composition and demographic variables, as explicated above.

Maternal education usually has a positive effect on child nutrition, as indicated by the meta-analysis, which found education to be positively correlated with child nutritional status, particularly of mothers. The relationship between maternal education and child well-being reflects a number of factors: greater educational attainment means greater income earning opportunities; schooling may impart knowledge of modern caring techniques directly; literacy means better ability to assimilate new information from newspapers; exposure to new environments due to schooling makes women more receptive to modern medical treatment; education improves self-confidence and therefore decision-making ability in the family; and schooling provides the opportunity to form social networks (Alderman et al. 2003). Some of the effects of education can therefore be broken down by controlling for income, father’s education, mother’s literacy, and knowledge of health and family planning.⁸

Other indicators of women’s power in the household likely to determine child nutritional status include women’s agency, mobility, and age and whether the household head is female. The share of economic resources devoted to children is often greater in households where women have greater say in decision making, though in the case of female-headed households (FHHs) the positive impact on child health and nutrition may be counter-balanced by the greater likelihood of both monetary and time poverty.

Community factors include location (urban/rural, division of residence) and environmental resources including access to clean drinking water, adequate sanitation, electricity and health services. Despite the apparent logic that greater food availability in the countryside would favor

rural child nutritional status over urban, urban location was found to exert a positive impact on child nutritional status in the meta-analysis, possibly due to better access to health facilities in urban areas and other factors relating to better communications and physical infrastructure (Charmarbagwala and others 2004).

Variables such as water and sanitation are key complements to food availability in determining child nutrition, because diseases such as diarrhea diminish the body’s nutrient intake; the meta-analysis found that water and sanitation were positively correlated with nutrition (ibid.). These are often termed “community variables” because access is often determined by location and because there are likely to be positive spillovers from one person’s consumption of clean water and sanitation to another person (e.g., by reducing exposure to contagious diseases).⁹ However, it may be so that those households with direct access to a facility, such as electricity, derive greater benefit than do other members of the community.

Mortality selectivity, a final estimation issue, concerns the lower censoring of malnutrition data due to child death. Malnourishment is associated with increased mortality risk (Gomez et al. 1956; Briend and others 1986), the risk of death from malnutrition being greatest between ages 6 and 36 months in one Matlab study (Fauveau et al. 1990). Since anthropometric data cannot be collected on children who are dead and these children are more likely to have been malnourished in life, the sample of live children is unlikely to be random (Lee and others 1997; Charmarbagwala and others 2004). In consequence, the nutrition model should be estimated conditional on survival probability; this is possible because DHS collect complete fertility and mortality histories of eligible mothers, enabling survival probability to be estimated. Lee and others (1997) do not find evidence for non-random selectivity in survival with respect to anthropometric measures.¹⁰

Data and Model

The data used in this study are from the nationally representative Demographic and Health Surveys (DHS) collected in 1996/97 and 1999/

2000. DHS compile complete fertility histories of ever-married women aged 10-49, as well as data on health status and health care, knowledge of health and family planning, socioeconomic factors such as education, household asset ownership, water and sanitation, and information on facilities operating in the community. In the survey rounds analyzed here, DHS also collected data on height and weight of children aged less than 5 years old, enabling calculation of anthropometric indicators.

Multivariate analysis of z scores is usually carried out on individual child-level data using ordinary least squares (OLS):

$$z_i = \beta X_i + e_i$$

where X is the vector of explanatory variables, β the vector of coefficients to be estimated and e a normally distributed random error. However, as noted above, the error term in OLS nutrition equations is unlikely to be random due to selectivity in survival; if the unobserved factors determining survival subsumed into the OLS error are correlated with the X s, OLS estimates of β will be biased. They will also be biased for any X that is correlated with e for reasons of endogeneity.

An appropriate model that accounts for sample selection is Heckman's two-step procedure. In the first stage the non-random selection variable—the probability of survival S —is estimated using a probit model:

$$\text{Prob}(S_i = 1) = \Phi(\gamma W_i)$$

where γ is the set of coefficients estimated on W explanatory variables and Φ indicates the cumulative standard normal density. From the fitted values of the probit model, the inverse Mill's ratio is calculated and included as an explanatory variable in the second-stage regression model of nutrition determinants:

$$z_i = \beta X_i + \lambda \frac{\phi(\gamma W_i)}{\Phi(\gamma W_i)} + u_i$$

ϕ being the probability density function of the normal distribution and λ the estimated coefficient on the inverse of Mill's ratio.¹¹ The procedure produces consistent estimates assuming that the error distribution of selection and re-

gression equations is bivariate normal (see Greene 2000). Unlike linear simultaneous equations systems, which require some difference in the variable set in each equation for identification, the Heckman two-step model is identified simply by the non-linearity of the selection probit. However, there should also be a theoretical justification of the model, which comprises variables determining selection probability but exogenous to child nutritional status. Variables satisfying this criterion are the community (non-self) means of attended births and mortality rates; means are estimated by mother's education level and sex of child, in order to improve accuracy of the instrument.

However, it would be incorrect to use a standard Heckman procedure incorporating a first-stage probit on the full sample of dead and live children, since mortality/survival data are right-censored (see Annex C). The solution adopted in this paper is to model the selection mechanism using survival analysis, the particular method being Cox's proportional hazards model (Annex C). The convenience of survival analysis is that the predictions of the regression model are the individual hazard rates, which are the same as inverse Mill's ratios. Note, however, that the estimated hazard rate is equal to the inverse Mill's ratio in the case of upper censoring (see Greene 2000):

$$\vartheta_i = \frac{\phi(\gamma W_i)}{1 - \Phi(\gamma W_i)}$$

whereas the correct Mill's ratio for the nutrition model is the lower-censoring case (above). The solution is to model non-selection (mortality) using a hazards model, generating the hazard rate which can be shown to be equal to the negative of the inverse of Mill's ratio in the case of lower-censoring for the selection model.¹²

Results

Table D.1 presents descriptive statistics of child nutritional status, as measured by anthropometric variables, by various correlates.¹³ On the whole, children have better mean anthropometric scores where they live in households that are non-poor (measured using a threshold value of

TABLE D.1

Weighted Means of HAZ, WAZ, and WHZ for Regression Sample by Various Characteristics

	HAZ score mean		WAZ score mean		WHZ score mean	
	1997	2000	1997	2000	1997	2000
All children	-2.31	-1.99	-2.24	-2.01	-1.11	-1.04
Non-poor	-1.94	-1.72	-1.95	-1.78	-0.99	-0.94
Poor	-2.53	-2.22	-2.41	-2.21	-1.18	-1.13
Household head male	-2.30	-1.99	-2.24	-2.01	-1.11	-1.04
Household head female	-2.37	-1.82	-2.22	-1.91	-1.05	-1.02
No electricity	-2.45	-2.15	-2.34	-2.15	-1.15	-1.10
Electricity	-1.83	-1.59	-1.88	-1.68	-0.99	-0.90
Unsanitary toilet	-2.48	-2.17	-2.35	-2.16	-1.13	-1.11
Sanitary toilet	-2.05	-1.81	-2.08	-1.86	-1.09	-0.98
Non-piped water	-2.34	-2.01	-2.26	-2.03	-1.12	-1.06
Piped water	-1.29	-1.32	-1.46	-1.33	-0.84	-0.67
Mother no schooling	-2.52	-2.20	-2.40	-2.19	-1.17	-1.12
Mother some primary education	-2.33	-2.12	-2.22	-2.12	-1.07	-1.10
Mother completed primary	-2.19	-1.87	-2.11	-1.89	-1.01	-0.97
Mother some secondary educ	-1.77	-1.66	-1.89	-1.74	-1.03	-0.92
Mother completed secondary	-1.42	-1.26	-1.64	-1.44	-1.00	-0.80
Mother higher educated	-0.98	-0.98	-1.25	-1.18	-0.79	-0.72
Mother mobility low	-2.30	-2.06	-2.24	-2.07	-1.12	-1.06
Mother mobility medium	-2.38	-1.97	-2.27	-1.99	-1.11	-1.03
Mother mobility high	-1.76	-1.79	-1.86	-1.87	-1.01	-1.02
Mother agency low		-2.10		-2.10		-1.07
Mother agency medium		-1.92		-1.96		-1.02
Mother agency high		-1.86		-1.91		-1.03
Female child	-2.34	-2.03	-2.27	-2.05	-1.09	-1.03
Male child	-2.27	-1.94	-2.20	-1.97	-1.13	-1.05
Barisal Division	-2.42	-2.06	-2.15	-2.10	-0.90	-1.12
Chittagong Division	-2.44	-2.06	-2.35	-2.04	-1.18	-1.04
Dhaka Division	-2.31	-1.97	-2.23	-1.97	-1.09	-1.00
Khulna Division	-2.00	-1.77	-2.04	-1.82	-1.08	-0.96
Rajshahi Division	-2.18	-1.93	-2.18	-2.03	-1.15	-1.11
Sylhet Division	-2.61	-2.25	-2.40	-2.20	-1.08	-1.08
Rural	-1.79	-1.62	-1.85	-1.73	-0.98	-0.96
Urban	-2.36	-2.06	-2.27	-2.06	-1.12	-1.06

ously? Multivariate regression analysis enables this question to be answered, while also accounting for censoring and direction of causality. Table D.2 shows results of the Cox proportional hazards models for under-fives. Deaths of other children in the community are a significantly positive determinant of risk of death of the index child; the coefficient on (non-self) share of children delivered by a doctor or nurse has a negative sign but is estimated imprecisely; other determinants were identified and discussed elsewhere (Annex C). The results are discussed for determinants of height for age z score (HAZ), weight for age z score (WAZ), and weight for height z score (WHZ) separately. In each case, standard OLS regressions and two-step regressions controlling for survival selectivity are presented. Modeling is undertaken for children aged between 6 and 60 months,¹⁴ and separately for 1997 and 2000 data. A Chow test indicated that the coefficients from separate regressions for each year were jointly significantly different from one another.

the wealth index), in which the head is female, which have electricity, sanitary toilet facilities and piped water into the dwelling, and which are located in rural areas and outside of Chittagong and Sylhet Divisions. Children fare better, on average, the higher their mother's level of education, the higher their mother's mobility and agency, and if they are of a non-Muslim religious group.

To what extent are these trends reproduced when controlling for many factors simultane-

Height for Age

The HAZ regressions provide a reasonably good fit of the data, as indicated by the values of R -squared (tables D.3-D.4). The OLS regressions indicate that:

- Household wealth exerts a significantly positive effect on nutritional status.
- Services are important determinants of HAZ, as indicated by the positive effect of access to

electricity and, in 1997, use of a sanitary toilet facility. In contrast, although the estimated coefficient is positive, access to piped water into the dwelling does not appear to significantly improve HAZ.

- Maternal education at secondary or higher levels results in significantly better child height. Children also have greater HAZ where their mothers have more say in household decisionmaking regarding cooking, purchases, and health care, measured by the agency index. Children of shorter and younger mothers have lower HAZ.
- Girls under five are significantly shorter than boys, controlling for other factors, particularly in poorer households, as indicated by the interaction between gender and household wealth.
- Of the fertility-related factors, children of higher birth order and those born less than 24 months after the preceding child tend to be shorter. Children living in households with more adults relative to children fare better nutritionally. These variables were omitted in order to test for presence of bias on other coefficients, the effect being to increase the magnitude and significance of the coefficient on the inverse Mill's ratio (results not reported).
- Children born in rural areas in the lean season are shorter on average, as are children born in Barisal, Chittagong, and, especially, Sylhet, the poorest division.
- Results from 2000 indicate that children living in households with food deficit during part or all of the year and low birth weight children are significantly more likely to be short for their age.
- It was possible to estimate the effect of whether the household resides in a cluster covered by the Bangladesh Integrated Nutrition Project (BINP) in 2000.¹⁵ The BINP variable is insignificant alone, but significantly positively correlated with HAZ when included alongside the interaction between BINP and maternal education, which itself is significantly negative, suggesting that the effect of BINP is stronger for mothers with low levels of education.

TABLE D.2	Under-Five Mortality: Cox Regression Results			
	1997		2000	
	HAZ Ratio	z	HAZ ratio	z
Wealth	0.834	-0.63	0.711	-1.15
Electricity	0.836	-1.22	0.845	-1.15
Water & sanitation	0.383	-1.56	0.569	-1.18
Mother primary ed	0.951	-0.45	0.860	-1.23
Mother lower secondary	0.822	-1.02	0.974	-0.15
Mother higher secondary	0.741	-0.56	0.269**	-2.18
Contraceptive knowledge	0.924***	-2.65	0.880***	-3.57
Multiple birth	7.254***	12.45	6.534***	9.53
Mother age at birth	0.977*	-1.80	0.988	-0.82
Previous child dead	1.095	0.70	1.022	0.13
Preceding interval <15	2.163***	4.74	2.789***	4.56
Prev dead * prec interval			0.511	-1.58
Female child	0.973	-0.32	0.879	-1.39
Birth order	0.868**	-2.00	0.726***	-3.79
Birth order sq	1.209***	3.59	1.343***	5.08
Female head	1.032	0.19	1.010	0.05
Maternal mobility	0.837**	-2.51	0.856*	-1.78
Maternal agency			1.009	0.53
Chittagong Division	0.824	-1.21	0.702*	-1.79
Dhaka Division	0.823	-1.28	1.013	0.07
Khulna Division	0.716*	-1.76	0.594**	-2.31
Rajshahi Division	0.823	-1.27	0.867	-0.74
Sylhet Division	0.988	-0.07	1.180	0.82
Rural	0.747*	-1.91	0.745**	-2.18
Mean—deaths in community	3.988***	2.62	3.406***	2.61
Mean—doctor/nurse delivery	0.768	-0.72	0.928	-0.24
Food deficit			0.988	-0.10
Low birth weight			1.414***	3.29
BINP			1.306	0.92
BINP * mother educ			0.413***	-3.01
# Obs	6332		6974	
# Deaths	610		522	
Wald chi-squared	319.333***		350.888***	
Log-likelihood	-4984.2		-4400.6	

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

Table D.3 also shows results of HAZ regressions controlling for selectivity in survival using the two-step method. Estimated coefficients on selection terms (LAMBDA) in each equation are significantly positive, providing statistical evidence for selectivity in survival, and indicating that the unobserved characteristics determining mortality risk and malnutrition are positively correlated—in other words, children who are more likely to die are estimated to be those more

TABLE D.3 Results of OLS and Two-Step Regressions of HAZ Score

	1997 (OLS)		1997 (2-step)		2000 (OLS)		2000 (2-step)	
	Coeff.	t	Coeff.	t	Coeff.	t	Coeff.	t
Wealth	0.525***	3.00	0.502*	2.86	0.481***	3.28	0.401***	2.73
Finished floor	0.057	0.59	0.058	0.60	-0.023	-0.29	-0.013	-0.17
Electricity	0.228***	3.25	0.214***	3.05	0.164***	2.98	0.145***	2.62
Sanitary toilet	0.102**	1.96	0.100*	1.92	0.033	0.73	0.031	0.68
Piped water	0.165	1.16	0.141	0.98	0.031	0.32	-0.007	-0.07
Mother primary educ	0.072	1.24	0.062	1.08	0.036	0.73	-0.003	-0.05
Mother lower secondary	0.316***	3.92	0.294***	3.62	0.243***	3.58	0.222***	3.25
Mother higher secondary	0.687***	4.23	0.659***	4.04	0.505***	4.15	0.413***	3.32
Adult/child ratio	0.071***	2.93	0.070***	2.89	0.014	0.79	0.012	0.70
Mother age	0.023***	3.59	0.021***	3.20	0.025***	4.42	0.023***	4.10
Female head	-0.043	-0.44	-0.038	-0.38	0.063	0.71	0.067	0.74
Mother mobility	0.019	0.53	0.008	0.21	0.035	1.03	0.008	0.23
Mother agency					0.014*	1.89	0.013*	1.83
Log mother height	0.968***	3.86	0.971***	3.85	0.743***	3.49	0.743***	3.50
Female child	-0.154**	-2.32	-0.160**	-2.42	-0.115*	-1.86	-0.146**	-2.39
Female*wealth	0.237	1.37	0.248	1.44	0.195	1.41	0.221	1.62
Birth order	-0.113**	-2.56	-0.120***	-2.75	-0.164***	-3.83	-0.198***	-4.54
Birth order square	0.037	0.89	0.050	1.23	0.075*	1.87	0.110***	2.66
Prec interval<24	-0.126**	-2.37	-0.115**	-2.16	-0.220***	-4.36	-0.192***	-3.76
Child's age	-0.071***	-10.89	-0.069***	-10.46	-0.045***	-7.95	-0.042***	-7.42
Child's age sq/10	0.083***	8.39	0.081***	8.15	0.051***	6.04	0.048***	5.66
Rural	0.155*	1.74	0.139	1.55	-0.050	-0.72	-0.084	-1.18
Born Feb-Mar*rural	-0.139*	-1.86	-0.145*	-1.95	-0.251***	-3.41	-0.253***	-3.47
Born Apr-May*rural	-0.137*	-1.71	-0.142*	-1.77	-0.078	-1.07	-0.082	-1.14
Born Jun-Jul*rural	-0.149*	-1.83	-0.152*	-1.87	-0.015	-0.20	-0.017	-0.23
Born Aug-Sep*rural	-0.012	-0.16	-0.016	-0.21	0.040	0.54	0.035	0.49
Born Oct-Nov*rural	0.006	0.08	0.003	0.04	0.042	0.64	0.042	0.65
Chittagong Division	-0.026	-0.31	-0.041	-0.49	-0.052	-0.67	-0.091	-1.15
Dhaka Division	0.144*	1.86	0.126	1.62	0.044	0.57	0.056	0.72
Khulna Division	0.437***	4.93	0.408***	4.54	0.190**	2.41	0.134*	1.67
Rajshahi Division	0.370***	4.69	0.350***	4.41	0.196**	2.48	0.177**	2.23
Sylhet Division	-0.130	-1.40	-0.117	-1.26	-0.087	-1.00	-0.024	-0.27
Mean-measles vacc.	0.046	0.15	0.050	0.16	0.139	0.62	0.144	0.63
Food deficit					-0.073	-1.57	-0.077*	-1.66
Low birth weight					-0.424***	-8.23	-0.358***	-6.78
GINI					0.241*	1.92	0.266**	2.09
GINI*educ					-0.183***	-2.91	-0.231***	-3.56
Constant	-6.753***	-5.27	-6.613***	-5.13	-5.448***	-4.97	-5.121***	-4.67
Lambda			0.815**	2.00			1.993***	3.70
# Obs	4,258		4,258		4,719		4,719	
F-statistic	24.7***		24.1***		21.0***		21.2***	
R-squared	0.159		0.160		0.153		0.158	

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

likely to be malnourished. The correct specification is therefore the two-step method, which is reported for all subsequent HAZ regressions.

The routes through which education affects child anthropometry can be examined using indicators of women's literacy and knowledge (table D.4). Maternal literacy and, to a lesser extent, contraception knowledge (a proxy for knowledge of health and sanitary practices) have positive impacts on HAZ of children when used independently of education. However, inclusion of schooling variables (specification 2) knocks out the significance of literacy, while leaving the impact of contraception knowledge virtually unaffected. This suggests the beneficial impact of maternal schooling on child nutritional status may operate partially through improving literacy, though there are likely to be other important factors at work.

The preceding analysis attempted to control for child immunization using the (non-self) share of children receiving measles vaccination, which, though a positive sign, as expected, was insignificant in all regressions. We now restrict analysis to children aged over 12 months in order that the individual child's immunization status can be used. Table D.5 shows results of regression for children aged over 12 months, including the indicator of whether the child received measles vaccination. In two-step regressions (specification 1), the effect of measles is significantly positive in 2000. However, due to the possibility of upwards bias on the measles vaccination coefficient due to unobserved heterogeneity, the second columns (specification 2) for each year report regression results accounting for the endogeneity of vaccinations (as well as the survival selectivity).¹⁶ The estimated coefficients on RHO indicate that the errors are correlated between the equations (though the correlation is marginally insignificant), while the estimated coefficient on measles becomes insignificant. Thus, though evidence is not conclusive, the results appear to provide indication of upwards bias in the estimated effect of vaccination due to unobserved heterogeneity.

Weight for Age

Results for weight for age (table D.6) are broadly similar to those for height for age. Wealth, elec-

tricity, and maternal education (particularly secondary) have significantly positive effects on child weight.¹⁷ Age of the child has a significantly convex relationship with WAZ. The survival selection terms (LAMBDA) are again significant, indicating that the two-step regressions accounting for non-random selection bias are the better specification.¹⁸

However, there are some differences with height for age results. For example, piped water in 2000 has a significantly positive effect on weight for age. On top of the effect of household wealth, children living in households with finished, as opposed to mud and wooden, floors are estimated to be heavier for their age, consistent with a hypothesis that cleanable floors can help reduce the spread of pathogens. Children measured during the interviews carried out in November and December were also found to have the lowest WAZ scores on average. Measles vaccination, measured using the (non-self) share of children vaccinated in the community has a positive effect on WAZ, which is significant in 2000.

Weight for Height

In contrast with HAZ and WAZ, the WHZ score regressions fit much more poorly (table D.7). Significantly positive impacts on WHZ are found for wealth, finished flooring, piped water (though sanitary toilet is estimated to have a perverse negative impact in 1997), and measles vaccinations in 2000. The selection terms in the two-step regressions do not provide evidence for non-random sample selection bias; we may expect mortality to be less closely related to a short-term malnutrition indicator such as WHZ than longer-term one (HAZ or WAZ).

Decompositions

Decompositions examining the contribution of different variables to the improvements in anthropometric outcomes over time are presented in table D.8, which shows the means of the explanatory variables used in the calculations (which are restricted to those variables which are statistically significant in regression analysis), and the percentage change in anthropometric measure due to each variable. The most impor-

TABLE D.4
Impact of Maternal Education on HAZ score (OLS and Two-Step Regressions)

	1997 (1)		1997 (2)		2000 (1)		2000 (2)	
	Coeff.	t	Coeff.	t	Coeff.	t	Coeff.	t
Wealth	0.536***	3.05	0.491***	2.76	0.423***	2.89	0.392***	2.65
Finished floor	0.087	0.89	0.061	0.63	0.022	0.29	-0.012	-0.16
Electricity	0.214***	3.05	0.215***	3.06	0.138**	2.50	0.144***	2.60
Sanitary toilet	0.099*	1.91	0.095*	1.83	0.039	0.88	0.031	0.69
Piped water	0.208	1.45	0.145	1.01	0.015	0.15	-0.008	-0.08
Mother primary educ			0.052	0.81			-0.019	-0.34
Mother lower secondary			0.271**	2.37			0.174*	1.89
Mother higher secondary			0.628***	3.49			0.368***	2.72
Literacy	0.171***	2.66	0.019	0.22	0.151***	2.86	0.054	0.76
Contraception	0.021	1.29	0.019	1.17	0.002	0.10	0.000	-0.01
Adult/child ratio	0.074***	3.04	0.071***	2.92	0.012	0.72	0.012	0.68
Mother age	0.023***	3.59	0.021***	3.24	0.025***	4.56	0.023***	4.09
Female head	-0.038	-0.38	-0.036	-0.36	0.051	0.57	0.063	0.71
Mother mobility	0.013	0.37	0.003	0.07	0.017	0.51	0.008	0.24
Mother agency					0.014*	1.85	0.013*	1.83
Log mother height	0.987***	3.85	0.966***	3.82	0.748***	3.49	0.741***	3.49
Female child	-0.164**	-2.47	-0.159**	-2.40	-0.142**	-2.32	-0.145**	-2.37
Female*wealth	0.269	1.55	0.250	1.45	0.221	1.61	0.220	1.61
Birth order	-0.136***	-3.12	-0.121***	-2.77	-0.210***	-4.86	-0.198***	-4.54
Birth order square	0.060	1.46	0.051	1.25	0.117***	2.85	0.110***	2.67
Prec interval<24	-0.106**	-1.99	-0.113**	-2.13	-0.183***	-3.61	-0.191***	-3.76
Child's age	-0.069***	-10.42	-0.070***	-10.51	-0.042***	-7.40	-0.042***	-7.39
Child's age sq/10	0.081***	8.10	0.082***	8.21	0.048***	5.60	0.048***	5.64
Rural	0.133	1.47	0.142	1.58	-0.087	-1.23	-0.084	-1.19
Born Feb-Mar*rural	-0.144*	-1.94	-0.146**	-1.96	-0.254***	-3.48	-0.253***	-3.47
Born Apr-May*rural	-0.134*	-1.67	-0.143*	-1.78	-0.080	-1.10	-0.083	-1.15
Born Jun-Jul*rural	-0.155*	-1.91	-0.153*	-1.89	-0.019	-0.26	-0.018	-0.25
Born Aug-Sep*rural	-0.012	-0.15	-0.017	-0.22	0.037	0.50	0.035	0.48
Born Oct-Nov*rural	0.006	0.08	0.002	0.02	0.045	0.69	0.043	0.66
Chittagong Division	-0.022	-0.26	-0.026	-0.31	-0.068	-0.86	-0.087	-1.09
Dhaka Division	0.139*	1.80	0.134*	1.72	0.074	0.96	0.061	0.78
Khulna Division	0.419***	4.67	0.413***	4.60	0.145*	1.81	0.138*	1.71
Rajshahi Division	0.354***	4.49	0.357***	4.49	0.204***	2.56	0.182**	2.29
Sylhet Division	-0.094	-0.98	-0.090	-0.94	-0.002	-0.02	-0.021	-0.23
Mean-measles vacc.	0.309	1.01	0.045	0.14	0.269	1.24	0.131	0.58
Food deficit					-0.086*	-1.86	-0.077*	-1.67
Low birth weight					-0.359***	-6.78	-0.357***	-6.73
GINI					0.251**	1.96	0.269**	2.11
GINI*educ					-0.223***	-3.36	-0.232***	-3.57
Constant	-6.840***	-5.23	-6.673***	-5.16	-5.217***	-4.72	-5.106***	-4.66
Lambda	0.795*	1.91	0.739*	1.77	1.978***	3.57	1.994***	3.52
# Obs	4,258		4,258		4,719		4,719	
F-statistic	23.6***		22.8***		20.9***		20.2***	
R-squared	0.157		0.160		0.155		0.158	

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

TABLE D.5

Impact of Measles Vaccination on HAZ Score, Children Aged 12-59 Months

	1997				2000			
	(1)		(2)		(1)		(2)	
	Coeff.	t	Coeff.	t	Coeff.	t	Coeff.	t
Dep var. = HAZ score								
Wealth	0.606***	3.82	0.600***	3.78	0.507***	3.62	0.519***	3.70
Finished floor	0.107	1.02	0.109	1.04	-0.021	-0.27	-0.022	-0.28
Electricity	0.221***	2.93	0.209***	2.73	0.112**	1.99	0.121**	2.14
Sanitary toilet	0.124**	2.25	0.124**	2.24	0.049	1.03	0.051	1.07
Piped water	0.074	0.49	0.075	0.49	-0.076	-0.77	-0.074	-0.75
Mother primary educ	0.035	0.59	0.025	0.41	-0.017	-0.32	-0.006	-0.12
Mother lower secondary	0.282***	3.44	0.271***	3.25	0.245***	3.63	0.268***	3.86
Mother higher secondary	0.602***	3.71	0.588***	3.61	0.468***	3.82	0.490***	3.99
Adult/child ratio	0.091***	3.51	0.091***	3.52	0.018	0.93	0.017	0.91
Mother age	0.022***	3.32	0.021***	3.11	0.021***	3.57	0.022***	3.73
Female head	-0.060	-0.59	-0.065	-0.62	0.087	0.93	0.086	0.92
Mother mobility	0.005	0.13	0.002	0.05	0.024	0.70	0.028	0.81
Mother agency					0.012	1.53	0.013*	1.68
Log mother height	0.986***	4.33	0.986***	4.33	0.670***	3.28	0.673***	3.29
Female child	-0.110**	-2.44	-0.104**	-2.29	-0.104***	-2.64	-0.111***	-2.78
Birth order	-0.118***	-2.69	-0.115***	-2.61	-0.195***	-4.28	-0.200***	-4.36
Birth order square	0.039	0.98	0.040	1.00	0.119***	2.75	0.119***	2.76
Prec interval<24	-0.109**	-1.96	-0.110**	-1.97	-0.181***	-3.40	-0.182***	-3.43
Child's age	-0.017*	-1.85	-0.017*	-1.86	0.000	0.05	0.000	0.06
Child's age sq/10	0.015	1.19	0.015	1.20	-0.006	-0.53	-0.006	-0.54
Rural	0.180*	1.89	0.182*	1.90	-0.055	-0.75	-0.061	-0.83
Born Feb-Mar*rural	-0.254***	-3.16	-0.254***	-3.16	-0.372***	-4.73	-0.372***	-4.74
Born Apr-May*rural	-0.280***	-3.18	-0.280***	-3.18	-0.222***	-2.91	-0.222***	-2.90
Born Jun-Jul*rural	-0.209**	-2.40	-0.209**	-2.40	-0.108	-1.36	-0.108	-1.35
Born Aug-Sep*rural	-0.041	-0.50	-0.041	-0.51	-0.001	-0.01	0.000	-0.01
Born Oct-Nov*rural	0.024	0.32	0.024	0.32	0.045	0.68	0.046	0.69
Chittagong Division	-0.048	-0.54	-0.038	-0.42	-0.059	-0.73	-0.061	-0.74
Dhaka Division	0.172**	2.09	0.180**	2.19	0.126	1.57	0.109	1.35
Khulna Division	0.481***	5.00	0.470***	4.82	0.194**	2.34	0.197**	2.38
Rajshahi Division	0.406***	4.82	0.401***	4.73	0.267***	3.21	0.262***	3.15
Sylhet Division	-0.050	-0.51	-0.036	-0.36	0.042	0.44	0.022	0.23
Measles vacc.	0.018	0.30	0.135	0.86	0.105**	2.03	-0.068	-0.50
Food deficit					-0.097**	-1.99	-0.097**	-1.98
Low birth weight					-0.349***	-6.28	-0.347***	-6.24
BNP					0.257*	1.89	0.274**	2.01
BNP*mother educ					-0.230***	-3.58	-0.233***	-3.63
Constant	-7.733***	-6.59	-7.798***	-6.62	-5.647***	-5.30	-5.556***	-5.21
Lambda	0.606***	3.82	0.689*	1.78	1.812***	3.34	1.862***	3.40
Dep var. = Measles vaccination								
Wealth			0.173	1.04			0.214	1.42
Electricity			0.398***	4.61			0.158***	2.28
Previous child died			-0.058	-0.71			0.085	1.04
Prec interval < 15			0.153	1.10			-0.192	-1.57
Birth order			-0.109***	-4.94			-0.091***	-4.40

(Continued)

TABLE D.5

Impact of Measles Vaccination on HAZ Score, Children Aged 12-59 Months (*continued*)

	1997				2000			
	(1)		(2)		(1)		(2)	
	Coeff.	t	Coeff.	t	Coeff.	t	Coeff.	t
Dep var. = HAZ score								
Measles vacc.	0.018	0.30	0.135	0.86	0.105**	2.03	-0.068	-0.50
Mother's age at birth			0.036***	4.78			0.027***	4.01
Female child			-0.170***	-3.43			-0.145***	-3.04
Female head			0.144	1.24			0.020	0.17
Mother primary educ			0.287***	4.18			0.165***	2.69
Mother lower secondary			0.371***	3.58			0.473***	5.26
Mother higher secondary			1.206***	3.84			0.974***	3.56
Contraception			0.038**	2.02			0.055***	2.99
Maternal mobility			0.085*	1.89			0.063	1.40
Maternal agency							0.021**	2.28
Mother remarried			-0.113	-1.06			-0.467***	-4.22
Community TV			0.063	0.88			0.187***	2.65
Distance to Thana hq			-0.004	-0.65			-0.026***	-4.22
Chittagong Division			-0.270***	-2.59			0.026	0.26
Dhaka Division			-0.229**	-2.39			-0.282***	-2.97
Khulna Division			0.481***	3.68			0.138	1.27
Rajshahi Division			0.192*	1.92			-0.060	-0.60
Sylhet Division			-0.332***	-2.94			-0.262**	-2.44
BINP							0.298**	2.45
Constant			-0.179	-0.71			0.335	1.45
Rho			-0.053	-0.83			0.088	1.36
Chi-squared (rho = 0)			0.69				1.95	
Prob > chi-squared			0.41				0.16	
# Obs	3,779		3,779		4,243		4,243	
F-statistic	18.3***				19.3***			
R-squared	0.135				0.153			
Wald chi-squared			590.4***				721.2***	
Log likelihood			-8204.6				-8857.8	

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

tant factors explaining mean improvements in anthropometry over time are general increases in wealth, secondary education of mothers (particularly up to Grade 10) and electricity. The large estimated effect of changes in birth order over time on HAZ and WAZ indicate that reductions in fertility have also played an important role in improving nutrition.

Conclusion

The conclusion that emerges from this analysis is that improvements in wealth and maternal secondary education had strong effects on reducing child malnutrition in the late 1990s in

Bangladesh. Presence of BINP in the community has been beneficial to the nutritional status of children of lesser educated mothers. This is in contrast to the effect of BINP on survival (Annex C), which was estimated to be beneficial for better educated mothers only. One explanation for these seemingly contradictory results is that certain aspects of BINP contribute to malnutrition reduction, e.g., supplementary feeding and nutrition education, because they are most beneficial to less educated mothers; in contrast, BINP reduces mortality most effectively where mothers are more willing to seek modern medical care, which is more likely among the educated.

TABLE D.6

Results of OLS and Two-Step Regressions of WAZ Score

	1997 (OLS)		1997 (2-step)		2000 (OLS)		2000 (2-step)	
	Coeff.	t	Coeff.	t	Coeff.	t	Coeff.	t
Wealth	0.620***	5.51	0.61***	5.40	0.489***	4.78	0.444***	4.30
Finished floor	0.151*	1.89	0.15*	1.90	0.090	1.40	0.095	1.49
Electricity	0.171***	2.91	0.16***	2.72	0.115***	2.73	0.102**	2.42
Sanitary toilet	-0.014	-0.35	-0.02	-0.40	0.014	0.41	0.013	0.38
Piped water	0.049	0.40	0.03	0.25	0.172*	1.92	0.145	1.61
Mother primary educ	0.073*	1.68	0.07	1.51	0.012	0.31	-0.015	-0.40
Mother lower secondary	0.130**	2.13	0.11*	1.84	0.155***	2.96	0.141***	2.70
Mother higher secondary	0.474***	3.13	0.45***	2.98	0.335***	3.13	0.270**	2.50
Adult/child ratio	0.051***	2.94	0.05***	2.89	0.018	1.23	0.017	1.16
Mother age	0.014***	2.94	0.01**	2.55	0.013***	3.15	0.012***	2.84
Female head	0.017	0.23	0.02	0.29	0.025	0.38	0.027	0.42
Mother mobility	0.000	-0.01	-0.01	-0.32	0.013	0.52	-0.006	-0.23
Mother agency					0.007	1.32	0.007	1.28
Log mother height	0.858***	4.73	0.86***	4.72	0.466***	3.70	0.467***	3.72
Female child	-0.068**	-2.13	-0.07**	-2.19	-0.053*	-1.89	-0.069**	-2.44
Birth order	-0.063*	-1.78	-0.07**	-1.96	-0.064**	-2.01	-0.088***	-2.70
Birth order square	0.009	0.28	0.02	0.58	0.018	0.59	0.042	1.36
Prec interval<24	-0.066	-1.60	-0.06	-1.40	-0.099***	-2.63	-0.080**	-2.10
Child's age	-0.031***	-6.06	-0.03***	-5.67	-0.022***	-4.76	-0.020***	-4.34
Child's age sq/10	0.045***	5.84	0.04***	5.59	0.030***	4.47	0.028***	4.15
Rural	0.175**	2.43	0.16**	2.19	0.072	1.37	0.045	0.85
Interview Feb*rural	-0.079	-1.31	-0.08	-1.29	-0.102*	-1.95	-0.103**	-1.99
Interview Mar*rural	-0.004	-0.04	-0.01	-0.11	-0.114	-1.45	-0.110	-1.40
Interview Nov*rural	-0.127**	-2.48	-0.13**	-2.49	-0.094	-1.64	-0.084	-1.46
Interview Dec*rural	-0.189***	-3.50	-0.18***	-3.42	-0.106*	-1.78	-0.101*	-1.70
Chittagong Division	-0.186***	-2.69	-0.20***	-2.85	0.012	0.16	-0.008	-0.11
Dhaka Division	-0.056	-0.90	-0.07	-1.12	0.026	0.36	0.041	0.57
Khulna Division	0.165**	2.28	0.14*	1.94	0.197***	3.19	0.159**	2.54
Rajshahi Division	0.033	0.48	0.02	0.27	0.088	1.39	0.078	1.23
Sylhet Division	-0.108	-1.40	-0.10	-1.30	-0.051	-0.74	-0.005	-0.07
Mean-measles vacc.	0.238	1.17	0.24	1.18	0.376	1.87	0.378	1.88
Food deficit					-0.053	-1.53	-0.056	-1.62
Low birth weight					-0.434***	-11.49	-0.388***	-10.01
BINP					0.225***	2.64	0.243***	2.82
BINP*educ					-0.076	-1.45	-0.109**	-2.04
Constant	-6.611***	-7.15	-6.50***	-6.99	-4.495***	-6.87	-4.282***	-6.53
Lambda			0.61*	1.78			1.393***	3.81
# Obs	4,258		4,258		4,719		4,719	
F-statistic	16.5***		16.0***		19.3***		19.2***	
R-squared	0.121		0.122		0.152		0.156	

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

TABLE D.7 Results of OLS and Two-Step Regressions of WHZ Score

	1997 (OLS)		1997 (2-step)		2000 (OLS)		2000 (2-step)	
	Coeff.	t	Coeff.	t	Coeff.	t	Coeff.	t
Wealth	0.376***	3.32	0.37***	3.29	0.205**	2.29	0.193**	2.13
Finished floor	0.171**	2.08	0.17**	2.08	0.142**	2.53	0.143**	2.56
Electricity	0.027	0.47	0.02	0.40	0.037	0.95	0.033	0.85
Sanitary toilet	-0.097**	-2.42	-0.10**	-2.44	-0.008	-0.25	-0.008	-0.26
Piped water	-0.059	-0.50	-0.07	-0.55	0.204**	2.55	0.196**	2.43
Mother primary educ	0.048	1.08	0.05	1.02	-0.007	-0.21	-0.015	-0.43
Mother lower secondary	-0.055	-0.82	-0.06	-0.90	0.032	0.64	0.028	0.56
Mother higher secondary	0.134	0.83	0.13	0.78	0.047	0.47	0.029	0.29
Adult/child ratio	0.011	0.61	0.01	0.60	0.015	1.12	0.015	1.10
Mother age	0.002	0.32	0.00	0.20	-0.002	-0.42	-0.002	-0.53
Female head	0.061	0.84	0.06	0.86	-0.006	-0.11	-0.005	-0.10
Mother mobility	-0.008	-0.27	-0.01	-0.37	-0.003	-0.13	-0.009	-0.36
Mother agency					-0.001	-0.31	-0.002	-0.32
Female child	0.054*	1.66	0.05	1.63	0.034	1.31	0.029	1.13
Birth order	-0.014	-0.42	-0.02	-0.48	0.034	1.19	0.028	0.93
Birth order square	0.001	0.03	0.00	0.14	-0.028	-1.10	-0.022	-0.80
Prec interval<24	-0.005	-0.13	0.00	-0.06	0.014	0.42	0.020	0.57
Child's age	-0.008	-1.44	-0.01	-1.33	-0.010**	-2.38	-0.010**	-2.24
Child's age sq/10	0.016**	2.09	0.02**	2.02	0.017***	2.73	0.017***	2.62
Rural	0.106	1.45	0.10	1.36	0.159***	3.39	0.152***	3.17
Interview Feb*rural	0.032	0.52	0.03	0.53	-0.072	-1.53	-0.072	-1.54
Interview Mar*rural	-0.040	-0.45	-0.04	-0.47	-0.087	-1.31	-0.086	-1.30
Interview Nov*rural	-0.125**	-2.34	-0.12**	-2.35	-0.199***	-3.77	-0.196***	-3.70
Interview Dec*rural	-0.170***	-3.03	-0.17***	-3.01	-0.207***	-3.88	-0.205***	-3.85
Chittagong Division	-0.280***	-3.72	-0.28***	-3.77	-0.036	-0.54	-0.042	-0.62
Dhaka Division	-0.194***	-2.87	-0.20***	-2.94	-0.044	-0.69	-0.040	-0.62
Khulna Division	-0.134*	-1.70	-0.14*	-1.78	0.127**	2.25	0.116**	2.00
Rajshahi Division	-0.281***	-3.77	-0.29***	-3.82	-0.010	-0.19	-0.013	-0.24
Sylhet Division	-0.086	-1.02	-0.08	-0.99	0.007	0.13	0.021	0.34
Mean-measles vacc.	0.285	1.22	0.29	1.22	0.439**	2.50	0.440**	2.51
Food deficit					-0.011	-0.35	-0.012	-0.38
Low birth weight					-0.257***	-7.76	-0.244***	-6.91
GINI					0.141	1.61	0.146**	1.66
GINI*educ					0.019	0.39	0.009	0.19
Constant	-1.119***	-5.97	-1.08***	-5.53	-1.140***	-7.82	-1.078***	-6.88
Lambda			0.21	0.75			0.394	1.11
# Obs	4,258		4,258		4,719		4,719	
F-statistic	3.88***		3.75***		6.98***		6.82***	
R-squared	0.029		0.029		0.053		0.054	

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively. %.

TABLE D.8	Decompositions for HAZ, WAZ, and WHZ Regressions				
	Mean (x_i)		Percentage change x_i		
	1997	2000	HAZ	WAZ	WHZ
Wealth	0.28	0.33	8.30	9.19	3.99
Finished floor				1.70	2.56
Electricity	0.23	0.29	3.56	2.51	
Sanitary toilet	0.41	0.51	1.18		-0.31
Piped water				0.41	0.56
Mother lower secondary	0.13	0.19	4.44	2.82	
Mother higher secondary	0.02	0.03	1.53	1.00	
Adult/child ratio	1.22	1.32	0.48	0.66	
Mother's age	26.07	26.19	1.02	0.52	
Log mother's height	5.00	5.00	0.51	0.32	
Female child	0.50	0.49	0.42	0.20	
Female child*wealth	0.14	0.16	2.06		
Birth order	3.14	2.94	14.38	6.38	
Birth order sq	1.44	1.28	-6.59	-2.52	
Prec interval <24	0.41	0.42	-0.65	-0.27	
Child's age	32.16	32.06	1.67	0.79	0.38
Child's age sq/10	12.77	12.71	-1.21	-0.70	-0.41
Rural	0.91	0.83	2.32	-1.25	-4.21
Born Feb/Mar	0.16	0.13	2.80		
Born Apr/May	0.13	0.13	0.17		
Born Jun/Jul	0.12	0.11	0.10		
Born Aug/Sep	0.13	0.13	0.00		
Born Oct/Nov	0.17	0.17	-0.04		
Interview Feb				-1.45	-1.01
Interview Mar				-1.24	-0.98
Interview Nov				1.76	4.10
Interview Dec				3.75	7.64
Chittagong	0.24	0.22	0.87	0.08	0.40
Dhaka	0.31	0.31	0.05	0.04	-0.04
Khulna	0.10	0.11	0.04	0.05	0.03
Rajshahi	0.21	0.23	1.06	0.47	-0.08
Sylhet	0.06	0.07	-0.08	-0.02	0.07
Mean-measles vacc.				0.79	0.92

Note: % change gives the share of the total change explained due to each variable x_j , that is:

$$\beta_{j2}(x_{j2}-x_{j1})/(\sum_j(\beta_{j2}(x_{j2}-x_{j1})) + \sum_i(x_{i1}(\beta_{i2}-\beta_{i1}))).$$

ANNEX E: WOMEN'S AGENCY, HOUSEHOLD STRUCTURE, AND HEALTH OUTCOMES

Bangladesh is often characterized as a country in which women face many restrictions. This is certainly so: recent data from DHS show that just over one in five women are solely responsible for decisions relating to their own health, and the large majority is not permitted to travel outside their household unaccompanied. But it is also easy to overstate the case, and to ignore that things are changing, and changing quite rapidly.

This paper begins by presenting data from the three DHS surveys from Bangladesh on women's role in decisionmaking (women's agency) and women's mobility. These issues are linked to household structure, which is shown here to be a critical determinant of female autonomy. Other determinants that may lay behind the changes that took place in the 1990s, such as increased women's work outside the home and the spread of secondary education, are also analyzed.

A related issue is age at marriage. The data show that the age at marriage is increasing. In most countries, raising the marriage age has been a critical factor in reducing fertility. However, it is argued that this is not likely to be the case in Bangladesh.

Household Structure

Household surveys contain a household roster in which basic information on all household members is collected. In DHS, as in most other surveys, all people who spend the night before the survey are included in the roster, with a further question asking if that person is a "usual resident" of the household. The roster also includes people deemed to be household members but absent the previous night. Not collecting this information means that female-headed households are "over-identified," that is, a household

in which the adult male is absent for prolonged reasons for work is not recorded in the roster, so that the wife may appear as the household head.

The first question asked about each person in the roster is their relationship to the household head. The responses to this question are tabulated in table E.1, for all 54,627 respondents of the DHS 1999/2000 survey and, in the second column, excluding the 3,190 respondents who were not usual residents. The final two columns tabulate the share for usual residents by sex.

The "core" of each household is the head, their husband or wife and children. Just under half of household members are children, around one-third of men are household heads, and one-third of women are the wife of the head. As explored below, extended households are not uncommon, as shown by 4.0 percent who are parents or parents-in-law and the 2.5 percent who are a son or daughter-in-law. The gender disaggregation shows these to be daughters-in-law: it is uncommon for a man to live with his parents-in-law (only 0.4 percent of men are the son-in-law of the head compared with the 5 percent of women who are daughters-in-law). Comparing the shares for the whole roster for those of "usual residents" shows that grandchildren, siblings, and other relatives are the most likely visitors. The small percentage of "not related" are generally not visitors, so these people are most likely to be live-in servants. The category co-spouse received zero responses. This does not mean there are no polygamous households, since the relationship shown is that to the household head; rather, it means there are no polygamous households in which one of the wives was identified as the head. For just eight people in total, the relationship was not known or not stated.

TABLE E.1		Relationship to Household Head (percent)		
	Whole roster	Usual residents only		
		Total	Male	Female
Head	18.0	19.1	34.7	3.3
Wife or husband	15.8	16.7	0.2	33.5
Son/daughter	43.9	45.3	50.6	40.0
Son/daughter-in-law	2.8	2.7	0.4	5.0
Grandchild	5.0	4.1	4.3	3.9
Parent	3.5	3.6	1.3	6.0
Parent-in-law	0.5	0.3	0.1	0.6
Brother/sister	3.4	3.1	3.9	2.4
Co-spouse	0.0	0.0	0.0	0.0
Other relative	5.3	3.2	2.9	3.6
Adopted/foster child	0.3	0.3	0.3	0.2
Not related	1.5	1.5	1.4	1.6
Don't know/missing	0.0	0.0	0.0	0.0
Absolute number	54,627	51,437	25,877	25,560

Source: DHS 1999/2000.

The data in the household roster can be used to identify five household types, where this classification is fully exhaustive and mutually exclusive (that is, it covers all households with no overlaps):¹

- *Extended*: a household including either a parent, parent-in-law, or son or daughter-in-law of the household head. White (1992) distinguishes between extended and joint households, where the latter live in the same compound but separate production and consumption (crucially, they do not eat together, which is the definition of household adopted

TABLE E.2		Household Types by Income Group (percent)			
		Income group			Total
		Low	Middle	High	
Extended		24.6	37.1	41.3	33.5
Nuclear		64.8	54.6	50.9	57.5
Female head		8.8	6.6	6.4	7.4
Single male headed		1.0	0.8	0.5	0.8
Polygamous		0.8	0.9	0.9	0.8
Absolute number of households		3,639	3,865	2,350	9,854


Source: DHS 1999/2000.

in some household surveys). It is not possible with DHS data to distinguish these two household types,² and White notes that the distinction is often more nominal than real.

- *Nuclear*: a household including the spouse of the head, but no parent, parent-in-law, or son or daughter-in-law
- *Female-headed household*: we do not simply take those households for which a woman is listed as the head, as some of these women have husbands or live with another adult male (table E.1). Rather, we classify as FHH those cases where a woman is identified as the head and there is no spouse, male parent, or son-in-law listed in the roster.
- *Single-male-headed household*: where there is no spouse listed.
- *Polygamous*: where the head has more than one wife.

The last column of table E.2 shows the breakdown of the 9,853 households covered by DHS 1999/2000 into these five categories. The majority of households are nuclear, though just over one-third are extended. The share of female-headed households is likely overstated for the reasons given above: that is, many of these households will have an absent male (husband) who supports the family and visits occasionally. A woman left alone after death or desertion of the spouse is more likely to return to her parents' household.³ The share of female-headed households can also be overstated where co-spouses live in separate households, which would also explain why the share of polygamous households is low, at less than one percent. Unsurprising is that the share of single-male-headed households is low, since men typically remarry on death of, or separation from, the spouse (White 1992; Kabeer 2004).

Also unsurprising is the correlation between household size and structure: the majority of households between two to six people are nuclear, but larger households are more likely to be extended (figure E.1). The largest household in the dataset is an extended family of 25 people consisting of an 80-year-old man and his 60-year-old wife, their four sons with their wives, one unmarried son aged 30, and 14 grandchildren.⁴

FIGURE E.1 Household Size and Household Structure

There is also a relationship between household structure and economic well-being as measured by an asset index based on housing quality and ownership of consumer durables (see Appendix C.2 on the construction of the asset index); see table E.2. Extended families are more common among higher-income groups, with the converse being true for nuclear families. Given that there is also the link between household size and structure, this finding means that large (extended) households are better off: a fact that runs counter to “standard economics poverty profiles,” in which household size is correlated with poverty, but is of no surprise to anthropologists, who report that households with resources need and attract labor, mostly through the family.⁵ For example, White (1992) attributes the likelihood of a household disintegrating to its lack of resources.

Patterns of Women's Agency and Mobility

Measurement

The Bangladesh Demographic Health Surveys contain a number of questions on women's

agency. The most recent survey contained a number of questions on women's say in decisionmaking. Specifically, if the woman had a say in decisions relating to:

- Her own health care
- Child health care
- Large household purchases
- Household purchases for daily needs
- Visits to family, friends, or relatives
- What food to cook each day.

For each question respondents could answer whether they make the decision alone, whether the husband makes it, “someone else” makes it, or whether they decide “jointly with husband” or “jointly with someone else.”

Both DHS 1999/2000 and the earlier surveys asked questions on women's mobility—that is, their ability to go to the market or health center either alone or with other family members. However, only one question (frequency of going shopping) has been asked the same way in all three surveys (table E.3).

TABLE E.3**Questions on Women's Mobility Contained in Different DHS Questionnaires**

	DHS93	DHS96	DHS99
If do go outside village/town/city alone		X	X
If can go outside village/town/city alone	X	X	X
If go to health center/hospital alone		X	X
If can go health center/hospital alone	X	X	X
Frequency of going shopping	X	X	X
With whom do they go shopping		X	X
Frequency of going outside village/town/city	X ^a	X	X

a. In 1993 the question was whether they go to another part of their village/town/city rather than if they leave it.

For questions on frequency of going shopping or going outside village/ town/city, respondents could answer “never,” “several times a year,” “once a month or more.” Answers were ranked/ordered accordingly. For questions on whether they go or can go outside village, town, or city or to health center, respondents could choose to answer one of the following: “don’t go,” “alone,” “with husband,” “with children,” or “other.” For our purposes, going “with husband” and “other” were pooled and the responses

ranked/ordered with “don’t go” as lowest ranking, followed by going “with others” (including husband), with children, and finally alone.

Women's Agency: Descriptive Analysis

Table E.4 summarizes the six aspects of women's agency collected by DHS 1999/2000. These data are tabulated for three groups: all respondents (which are ever-married women aged 11-49), currently married women in male-headed households (since we argue below that women in female-headed households largely make decisions by default), and women who live with their mother-in-law, since the literature suggests that these women will have less say in decisionmaking. The data indeed show that women living with their mother-in-law have a diminished role in decisionmaking compared with other women, whereas “someone else” (presumably the mother-in-law) is much more likely to have a say—and implicitly that women in female-headed households decide more than those in male-headed ones. We return to this issue below.

Other than the decision about what to cook, fewer than one in five women are solely responsible for the decisions shown, but over half are

TABLE E.4**Women's Say in Decisionmaking by Position in Household (percent)**

	Own health	Child's health	Large purchases	Daily purchases	Visits	Cooking
All respondents (n = 9,716)						
Respondent	20.6	19.2	10.3	19.5	14.0	66.8
Joint	36.1	47.1	51.0	43.8	48.3	20.4
Husband	36.9	26.9	29.4	27.2	29.5	4.6
Someone else	6.4	6.9	9.3	9.6	8.2	8.2
Currently married women in male-headed households (n = 8,706)						
Respondent	15.7	14.3	6.1	15.5	9.3	66.8
Joint	36.8	50.0	53.6	46.1	51.1	20.4
Husband	41.0	29.9	32.5	30.2	32.7	4.6
Someone else	5.1	5.8	7.7	8.2	6.9	8.2
Living with mother-in-law (n = 2,017)						
Respondent	12.2	10.3	3.2	9.8	5.2	42.8
Joint	36.2	48.5	50.5	45.2	48.1	31.3
Husband	39.5	27.1	26.1	22.8	26.7	3.3
Someone else	12.2	14.5	20.12	22.2	20.0	22.6

involved in the decision either solely or jointly. Women are least commonly solely responsible for the decision regarding large purchases; but husbands are also less commonly solely responsible for this decision, which is the most common joint decision.

Women's involvement in decisionmaking on matters related to their own health and that of their children is low. Forty-three percent of women have *no say* in the decisions on their own health; this number rises to 52 percent for women living with their mother-in-law. With respect to decisions related to child health, these figures are 34 and 42 percent, respectively.

Using the DHS 1999/2000 data, an index of women's agency (*AGENCY*) was constructed in two stages. First, the response to each of the six questions was recoded so that the respondent deciding by herself = 2, with husband or someone else = 1, and someone else decides = 0. Second, the responses to the six questions were summed together to give a 13-point scale from 0 to 12. A score of zero indicates that the respondent has no say in any of the six decisions, and 12 means she is solely responsible for all of them. For purposes of presentation, three categories of *AGENCY* were created: low for *AGENCY* between 0 and 4, medium for 5 to 7, and high for values of *AGENCY* between 8 and 12 inclusive.

Table E.5 tabulates the categorical version of the agency variable against the status of the woman in the household. The majority of the women in the sample (8,706 out of 10,543; 83

percent) are married women and usual residents of a male-headed household. Over one-third of these have a low level of agency, though nearly one-half have a medium level. We can distinguish the just over one-fifth of these women who live with their mother-in-law from those who do not. There is a significant difference in agency between these two groups: women living with their mother-in-law have a significantly lower say in decisionmaking, with nearly half having low agency, and less than 10 percent having high agency.

Agency is highest among the various categories of female-headed households. Nearly 95 percent of unmarried women heading their households with no man present have high agency. Perhaps more surprising is the high agency of female-headed households where the woman is married but the man absent. Closer inspection of the data shows that in these cases, the woman makes day-to-day decisions by herself, but the decision on large purchases is in many cases made either jointly or just by the husband. The third category comprises female-headed households living with at least one adult male. These are women who are divorced or widowed who live with either or both a parent or an adult son (defined as age greater than 15). These women also have a high level of agency.

An implication of this analysis is that when looking at the correlates of agency, it makes sense to focus on married women in male-headed households, who are the bulk of the

TABLE E.5**Women's Agency by Status in the Household**

Woman's marital status	Household type	Women's agency (percent)			Number of observations
		Low	Medium	High	
Married	Male-headed households	37.9	48.1	14.0	8,706
	o/w Living with mother-in-law	49.2	43.4	7.5	2,015
	Other	34.5	49.5	16.0	6,691
Unmarried	Female-headed (husband not present)	3.1	12.9	84.0	136
	FHH with no adult male	2.7	3.3	94.0	162
	FHH with adult male	3.3	16.9	79.9	112
	Other	35.7	31.3	33.0	574
Total		36.9	45.0	18.2	10,543

sample. Including women in female-headed households would undermine the power of the results, since these women are the decision-makers by virtue of their status, regardless of their other characteristics (wealth, religion, and the like).

Bivariate tabulations are shown for the former group of women against various characteristics in table E.6. The relationship between the variables shown is statistically significant in all cases. The main area where there is no difference is the similar distribution of agency between Muslim and Hindu women, although the small group of women from other religions do have higher

agency. A women's agency increases with household wealth, her age, whether she has had a child or not (with a son counting for more than a daughter), and if she is working (with working for cash having more of an impact). A slight kink in the pattern is the lopsided U-shaped relationship between high agency and asset quintile, with women in the lowest quintile having slightly greater agency than those in the second and third. This result may reflect that poorer households are more likely to be nuclear, and women have higher agency in these households. Multivariate analysis is used below to separate out these effects.

TABLE E.6	Agency by Selected Characteristics of Married Women in Male-Headed Households				
	Agency (percent)			Total	
	Low	Medium	High	Percent	Number of observations
Asset quintile (poorest = 1)					
1	41.9	44.7	13.4	100.0	1,980
2	40.2	49.0	10.8	100.0	1,374
3	40.3	47.2	12.5	100.0	1,552
4	37.5	49.2	13.3	100.0	1,697
5	29.0	51.3	19.8	100.0	2,103
Religion					
Muslim	38.3	47.3	14.4	100.0	7,553
Hindu	36.5	52.7	10.9	100.0	1,047
Other	21.4	61.1	17.5	100.0	106
Number and sex of children					
No children	53.2	42.1	4.7	100.0	840
Daughter(s) only	40.6	46.7	12.8	100.0	1,295
At least one son	35.5	49.1	15.4	100.0	6,571
Age group					
11–19	53.8	40.3	5.9	100.0	1,318
20–29	39.3	48.8	11.9	100.0	3,280
30–39	31.5	50.2	18.3	100.0	2,547
40–49	31.3	50.1	18.6	100.0	1,561
Work status					
Not working	40.5	46.6	13.0	100.0	6,993
Working (not for cash)	35.0	49.4	15.6	100.0	175
Working (for cash)	27.3	54.5	18.2	100.0	1,538
Total	37.9	48.1	14.0		8,706

Women's Mobility: Descriptive Analysis

An analysis of women's mobility using the same methodology as above reveals that mobility and agency are related to the same factors, even though, as shown below, the two are only imperfectly correlated.

Women's mobility is most restricted when it comes to going shopping, indicating that *pardah* and social norms continue to render the marketplace off-limits to women. Thus, a whole 68 percent of all women and three-fourths of women who live with their mother-in-law never go shopping. In comparison, only 4 to 5 percent of women cannot leave their village, town, or city of residence or go to the health center.

As is the case for intra-household agency, women who live with their mother-in-law have less mobility than other women. Fewer of these women can leave their village, town, or city alone, and more can only go if accompanied by their husband or someone else. The pattern is the same for going to the health center and going shopping.

An index of women's mobility (*MOBILITY*) was constructed in the same way as the agency index, using DHS 1999 data. As before, the response to each of the three questions was re-coded so that the respondent going out alone = 2, with husband or someone else = 1, and not at all = 0. Second, the responses to the questions were summed together to give a 10-point scale from 0 to 9. A score of zero indicates that the respondent cannot go out for any of the three purposes, and 12 means she can go out alone for all

of them. Again, three categories of *MOBILITY* were created: low for agency between 0 and 3, medium for 4 to 6, and high for values 7 to 10.

Table E.8 illustrates women's levels of mobility by their status in the household. Only one in eight married women who live in male-headed households have high mobility, almost half have medium mobility, and 40 percent have low mobility. As is the case for women's agency, women who live with their mother-in-law have significantly lower mobility than others, with over half having low mobility and only 7 percent having high mobility. Similarly, a much larger share of women whose husbands are away have high mobility (over one-third). Unsurprisingly, unmarried female heads of household with no adult male present are those with the highest level of mobility, with half having high mobility.

Analyzing women's mobility by selected characteristics for married women in male-headed households (table E.9) reveals a pattern similar to that of women's agency. There is a U-shaped relationship between assets and mobility, with women in wealth quintiles 1 and 2 having greater mobility than the women in quintile 3, and women in the poorest quintile also having greater mobility than women in quintile 4. Such a relationship is consistent with the notion that the poorest women in Bangladesh often are those who adhere least to *purdah* because their financial situation forces them to venture outside to work: the largest percentage of women who undertake paid work belong to the poorest

TABLE E.7	Women's Mobility (percent)		
	Can leave village/town	Can go to health center	Go shopping
All respondents (n = 9,716)			
Alone	17.0	30.0	13.8
With children	49.5	24.2	6.4
With husband or someone else	29.2	40.6	11.7
Cannot go	4.3	5.2	68.1
Currently married women in male-headed households (n = 8,704)			
Alone	14.8	28.4	11.5
With children	50.1	23.4	6.1
With husband or someone else	30.7	43.0	12.2
Cannot go	4.3	5.2	70.2
Living with mother-in-law (n = 2,017)			
Alone	10.2	22.6	6.6
With children	38.3	16.4	3.9
With husband or someone else	46.2	54.3	15.2
Cannot go	5.3	6.7	74.4

asset quintile,⁶ while, as illustrated in table E.8, a larger proportion of women who work have high mobility. This is consistent with the notion that women from better-off households can more easily afford to adhere to *purdah* and hence to have more restricted mobility.

The difference between Hindu and Muslim women is small. Just over 2 percent more from the former group have high mobility, whereas the percentage is 40 percent higher for the small

TABLE E.8		Women's Mobility by Status within the Household			
Woman's marital status	Household type	Women's mobility (percent)			Number of observations
		Low	Medium	High	
Married	Male-headed households	41.5	46.2	12.3	8,706
	o/w Living with mother-in-law	55.1	37.4	7.6	2,015
	Other	37.4	48.8	13.7	6,691
Unmarried	Female-headed (husband not present)	14.0	54.7	31.3	136
	FHH with no adult male	8.7	40.7	50.5	162
	FHH with adult male	5.7	55.3	39.1	112
	Other	26.0	45.0	29.0	574
Total		40.0	45.7	14.3	10,543

TABLE E.9**Mobility by Selected Characteristics for Married Women in Male-Headed Households**

	Mobility (percent)			Total
	Low	Medium	High	Number of observations
Asset quintile (poorest = 1)				
1	42.6	45.7	11.6	1,980
2	42.1	47.7	10.1	1,374
3	46.1	45.2	8.7	1,552
4	43.0	45.6	11.4	1,697
5	33.8	46.8	19.4	2,103
Religion				
Muslim	42.1	46.5	11.5	7,553
Hindu	40.1	46.0	13.9	1,047
Other	16.9	28.3	54.8	106
Number and sex of children				
No children	73.5	19.7	6.8	840
Daughter(s) only	48.6	39.0	12.4	1,295
At least one son	36.1	50.9	13.0	2,135
Age group				
11–19	69.4	26.5	4.1	1,318
20–29	42.5	45.7	11.8	3,280
30–39	32.8	51.5	15.7	2,547
40–49	29.2	55.7	15.1	1,561
Work status				
Not working	44.1	46.2	9.7	6,993
Working (not for cash)	27.6	57.3	15.2	175
Working (for cash)	32.5	44.6	22.9	1,538
Total	40.0	45.7	14.3	10,543

sample of women of other religions. Whether a woman has children—and whether at least one child is a son—has a significant impact on women's mobility, to an even greater extent than it impacts women's agency. Hence, low mobility is characteristic of nearly three-quarters of women with no children, just under half of women with only daughters, and just over a third of women who have at least one son.

Agency and Mobility

Although agency and mobility are associated with the same factors, they are highly, but not

perfectly, correlated. Table E.10 shows the cross-tabulation of the two. If there were no relation, then the entry in each cell would be 1/3 (33 percent), which is far from the case. Applying a chi-squared test shows the two to be highly related. This finding is somewhat contrary to that of some anthropologists, who argue that women's physical mobility does not necessarily reflect their intra-household agency, and that activities outside the homestead such as paid employment usually serve to further the household interest (White 1992, p. 79). The imperfect correlation, however, could indicate that this is true for some women but not all.

Changes over Time

Women's overall mobility has increased significantly over just a few years (see tables E.11a and E.11b). Almost a quarter of women could not leave their village, town, or city at all in 1993, but only 4 percent could not do so in 1999. Similarly, women who could not go to the health center decreased from over a fifth to less than 6 percent. At the same time, however, we see a very large increase in the percentage of women who can go with "others" and a decrease in those who can go alone—a change that may be partially explained by a slight difference in the way these questions were asked in 1993 and in the two later surveys. Finally, the proportion of women who never go shopping has decreased from over 80 percent to under two-thirds. This is concurrent with an increase in the proportion of women who go shopping once a month or more, which increased from less than one in 20 in 1993 to almost one in 5 in 1999. Thus, while going shopping is the area where women's mobility continues to be most severely restricted, there has been quite a significant change over a period of only six years, indicating a change of social norms and the practice of *purdah*. This does not mean that *purdah* is disappearing, but, as suggested by Kabeer (2002), it is adapting to changing socioeconomic conditions.

Multivariate Analysis

Regression analysis was performed for the agency and mobility indices and the health decision variables. The indices are discrete variables

in the ranges 0-12 and 0-9, so might appear to be able to be treated as continuous variables using ordinary least squares. However, a tabulation shows both series to be right censored (observations are bunched at the upper limit), so that a probit model is appropriate. The recoded agency and mobility variables related to health can be used as dependent variables in an ordered probit model. The sample in all cases is married women in male-headed households.

By and large, the results support the patterns shown in bivariate tabulations. The following results are particularly striking:

- Both age and age squared are significant, indicating that both agency and mobility increase with age, but then decrease after a certain threshold.⁷ The coefficients show this threshold to be in the early forties (see table A.3). This variable may also pick up a “time effect,” where older households afford women less agency and mobility. Having a child also increases agency, with the effect being stronger if the child is male.
- Both agency and mobility increase with educational status (the reference category is no education), though the effect of primary is not always significant. Given that there is a strong correlation between education and the level of the asset index (table A.2), it is striking that both sets of variables are frequently significant, showing that the impact of education can operate independently of that through wealth.
- The three media variables, especially TV, have a strong effect, though reading a paper is not always significant.⁸
- Daughters-in-law have a strongly significant lower level of agency and mobility. Mobility is higher for mothers-in-law, but not agency.
- The inverted U with respect to wealth is evident for agency but not mobility.
- Working has a strong positive effect, as does being a member of a credit organization.
- Regional effects are high, with women in Sylhet and Barisal faring worst.⁹ Rural women also have lower mobility and agency.
- The small number of women from other religions (the reference category is Muslim)

TABLE E.10		Women's Agency and Mobility (as percent of agency category)			
		Agency			Total
		Low	Medium	High	
Mobility	Low	48.6	51.3	22.7	45.5
	Medium	46.0	30.6	42.4	36.2
	High	5.4	18.1	34.9	18.3

TABLE E.11a		Changes in Mobility, 1993–99 (percent)				
		Can go outside village/town/city			Can go to health center	
		1993	1996	1999	1993	1996
No		24.9	15.6	4.3	22.0	13.1
With other		4.0	7.8	30.9	5.6	40.4
With children		34.0	49.0	47.2	38.1	25.8
Alone		37.2	27.6	17.6	34.3	20.7
Number of observations		7,308	7,308	10,543	9,639	9,127

TABLE E.11b		Changes in Mobility, 1993–99 (percent)		
		Frequency of shopping		
		1993	1996	1999
Never		81.9	78.8	64.4
Once/yr or less		6.7	6.2	5.2
Several times/yr		7.2	8.8	11.6
Once/month or more		4.1	6.2	18.9
Number of observations		9,639	9,127	10,543

have more agency and mobility. The Hindu dummy is negative but not significant.

In summary, the policy implications are that:

- It is true that women often do not have a say—so health and nutrition messages need to reach a wider audience.
- But agency and mobility can change over time, and are doing so quite rapidly in Bang-

ladesh. Policy relevant factors that can affect this rate of change are communications (IEC), and education.

Age at Marriage in Bangladesh

The age at marriage is a well-known determinant of maternal and child health outcomes. Children born to young mothers are more likely to die, so that raising the age of marriage, especially in a country such as Bangladesh, in which births outside of marriage are extremely rare, will reduce infant mortality. Traditionally, rising age at marriage has been a driver of fertility reduction, though there are reasons to think that this effect may be muted in Bangladesh, where age at marriage is indeed low, but fertility has already been substantially reduced (Annex F). Data from the Demographic and Health Survey show surprisingly large changes in the age at marriage over a very short period. This paper analyzes the determinants of age at marriage, and thus identifies factors behind these changes.


Marriage in Bangladesh

Virtually all women in Bangladesh get married. Data on over 110,000 women over the age of 10

from the three DHS surveys show that 99.4 percent of women over 30 have been married at some point (that is, are currently or formerly married). Figure E.2(a) shows the marital status for women in different age groups.¹⁰ This graph shows that, over 30, only a tiny proportion of women have never been married. It also shows that the vast majority of women are married by their early 20s, and that virtually all are married by the time they reach 25. As might be expected, older women are more likely to be formerly married, since their husbands predecease them. Husbands dying before their wives is not only a result of the shorter life expectancy of males, but also of the large age differential that is common in Bangladesh. Figure E.2(b) shows that men marry later than women, with the majority marrying by the time they are 30, rather than by their early 20s, as is the case for women. Most men under 20 are not married, whereas most women are.

However, the age of marriage is changing. Despite the short period covered by the DHS surveys, which might be expected to be too short to pick up significant social trends, women in Bangladesh are getting married later. As

FIGURE E.2 Marital Status by Age



shown in table E.12, the proportion of women married before they are 22 fell from 86 percent in the 1993 survey to 80 percent in 1999. The change is not among the youngest group—just under 5 percent of 10-15 year olds are married in all three surveys—but among those aged 16-21.

The assertion that the age at marriage is rising is contrary to the DHS reports, which show only a slight rise in the median age at marriage for women aged 20-49, from 14.1 to 14.7 years. However, this calculation is based on the age of marriage for women who have married. If a larger proportion of women are marrying later, then they simply drop out of this calculation, creating a downward bias in the sample estimate of the age of marriage. The presentation in table E.12 both avoids this problem, and shows that there is indeed a problem in basing the calculation simply on the age of marriage of those who have married. Alternatively, the median age at marriage can be calculated using the whole sample of women, rather than the ever-married sample.

A first look at why the age at marriage may be falling comes from bivariate tabulations of age at marriage and various individual and household characteristics. There is a problem in using this approach, as DHS collects detailed data only from ever-married women. Hence, the younger women in the sample are not representative, as they are married, and so, for example, are less likely to be at school than unmarried girls in the same age range. For this reason, girls under 15 are excluded from the tables. This problem can be tackled more satisfactorily in a multivariate setting, as seen below.

Looking at age at marriage by educational attainment (DHS 1999), we see that women's level of education affects their age at marriage positively. As illustrated in table E.14, women of higher educational status marry later, especially those with higher education. But notably more women with secondary education get married in their 20s, compared with women with less education. The regional pattern is not very strong, except that women in Sylhet appear to get married older.

Multivariate analysis may be done in two ways. First is a Cox hazards model (where the hazard is getting married). This model takes care of the

TABLE E.12		Median Age at Marriage					
	20-24	25-29	30-34	35-39	40-44	45-49	20-49
1993/94	15.3	14.8	14.2	13.9	13.6	13.6	14.1
1999/2000	16.1	15.4	14.9	14.5	14.0	13.8	14.7

TABLE E.13		Percentage of Women Ever Married		
		1993	1996	1999
10-15		4.9	4.7	4.9
16-18		30.7	30.4	28.1
19-21		50.6	48.9	47.1
Total married before 21		86.3	84.0	80.1

Source: Calculated from DHS 1993 and 1999.

TABLE E.14		Age at Marriage by Selected Characteristics (percent)			
	Age range				Number of observations
	16–18	19–21	22–25	>26	
Education status					
None	76.7	16.6	4.9	1.8	1,033
Primary	76.9	17.4	4.1	1.6	827
Secondary	69.4	21.2	7.9	1.5	1,228
Higher	31.9	39.3	21.8	7.0	514
Asset quintile					
1	78.3	14.9	5.5	1.3	457
2	73.3	19.1	5.7	1.9	419
3	74.3	18.0	5.7	2.1	529
4	70.3	22.1	6.1	1.5	787
5	59.1	25.5	11.8	3.5	1,410
Region					
Barisal	69.3	21.0	6.8	2.9	309
Chittagong	68.9	21.5	7.9	1.7	836
Dhaka	67.1	23.9	7.2	1.8	930
Khulna	71.6	21.3	5.2	1.9	536
Rajshahi	69.8	18.1	9.7	2.4	507
Sylhet	60.3	21.5	13.0	5.2	484
Total	2,444	777	294	87	

censoring problem. But the DHS does not have much data on unmarried women. An alternative is to use Heckman, first estimating a selection equation for if a woman is married or not, and then at what age those who are married got mar-

ried, using a sample selection bias term from the first-stage equation. The selection equation should contain at least one variable not in the age at marriage equation. In this case identification is easy: age is the overwhelming determinant of whether a woman is married or not, and that is all we need to enter in the selection equation.

The results (table E.15) show the expected relationship with education. There is a U-shape with assets: poorer and richer women tend to get married later, presumably as the poor need time to save for a dowry (and are considered less desirable brides and so a “last choice”) and the rich get married later for social reasons.

TABLE E.15		Regression Results for Age at Marriage		
	Heckman model		Cox Hazards model	
	Coefficient	z statistic	Hazard ratio	z statistic
No education	−5.46	−41.53***	5.06	33.40***
Primary	−4.93	−37.68***	3.86	27.71***
Secondary	−3.51	−27.56***	2.01	14.63***
Quintile 2	−0.15	−1.71*	1.08	2.15**
Quintile 3	−0.17	−1.86*	1.07	1.90*
Quintile 4	−0.07	−0.75	1.05	1.43
Quintile 5	0.02	0.17	1.01	0.36
Intercept	19.96	138.64***		
	Selection equation	t statistic		
Age	0.21	68.22***		
Intercept	−3.84	−65.76***		
/athrho	−0.52	−16.26***		
/lnsigma	1.02	141.33***		
Rho	−0.48			
Sigma	2.76			
Lambda	−1.33			
No. of observations	15,773		15,573	
Of which censored (Heckman)	5,230			
Of which failures (Cox)			10,543	

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

TABLE E.16

Women's Agency and Mobility: Survey-Ordered Probit Regression

Own	Coefficient	t	Child	Coefficient	t	Cangohealth	Coefficient	t
age	0.048	4.38***	age	0.035	2.98***	age	0.072	6.7***
age_sq	-0.001	-3.43***	age_sq	0.000	-2.35**	age_sq	-0.001	-5.26***
primary	-0.013	-0.36	primary	0.045	1.26	primary	0.033	0.99
secondary	0.010	0.22	secondary	0.068	1.61	secondary	0.151	3.51***
tertiary	0.226	3.18***	tertiary	0.224	3.39***	tertiary	0.359	4.64***
child	0.058	0.95	child	0.065	1.22	child	0.299	5.54***
child_son	-0.015	-0.38	child_son	0.091	2.6***	child_son	0.077	2.08**
listen_radio	-0.026	-1.7*	listen_radio	-0.007	-0.44	listen_radio	0.021	1.39
watch_tv	0.023	1.44	watch_tv	0.066	4.28***	watch_tv	0.066	4.17***
read_paper	0.015	0.87	read_paper	0.038	2.24**	read_paper	0.058	3.1***
wealth2	-0.049	-1.14	wealth2	-0.015	-0.37	wealth2	-0.025	-0.55
wealth3	-0.046	-1.01	wealth3	-0.029	-0.64	wealth3	-0.103	-2.21**
wealth4	0.014	0.31	wealth4	0.080	1.8*	wealth4	-0.055	-1.16
wealth5	0.118	2.1**	wealth5	0.176	3.37***	wealth5	-0.080	-1.5
DIL	-0.100	-2.74***	DIL	-0.145	-4.22***	DIL	-0.135	-4.39***
MIL	0.021	0.33	MIL	0.016	0.23	MIL	0.003	0.05
belong credit	-0.059	-1.47	belong credit	0.009	0.23	belong credit	0.229	5.56***
other org	-0.014	-0.33	other org	-0.022	-0.6	other org	0.105	2.5***
work	0.175	4.98***	work	0.182	5.04***	work	0.191	4.98***
rural	-0.105	-2.21**	rural	-0.098	-2.64***	rural	-0.157	-3.77***
Hindu	0.011	0.22	Hindu	-0.021	-0.52	Hindu	-0.076	-1.22
other_rel	0.229	2.49**	other_rel	0.269	2.4**	other_rel	0.422	2.35**
Chittagong	0.117	2.04**	Barisal	0.082	1.44	Sylhet	-0.134	-1.18
Sylhet	-0.141	-1.59	Chittagong	0.075	1.43	Khulna	0.075	1.27
Rajshahi	0.082	1.72*	Sylhet	-0.154	-1.99*	/cut1	-0.106	-0.7
/cut1	0.767	4.68***	/cut1	0.445	2.6***	/cut2	1.572	10.08***
/cut2	1.902	11.68***	/cut2	1.923	11.32***	/cut3	2.227	14.09***

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

TABLE E.17

Women's Agency and Mobility: Tobit Estimates

Agency	Coefficient	t	Mobility	Coefficient	t
age	0.206	8.48***	age	0.172	10.5***
age_sq	-0.003	-6.74***	age_sq	-0.002	-8.11***
primary	0.129	1.72*	primary	-0.016	-0.31
secondary	0.184	1.88*	secondary	0.259	3.92***
tertiary	0.680	4.21***	tertiary	0.968	8.84***
child	0.433	3.4***	child	0.437	5.09***
child_son	0.146	1.65*	child_son	0.131	2.2**
listen_radio	-0.019	-0.57	listen_radio	0.047	2.06**
watch_tv	0.174	5.22***	watch_tv	0.183	8.09***
read_paper	0.071	1.78*	read_paper	0.138	5.16***
wealth2	-0.080	-0.82	wealth2	-0.042	-0.63
wealth3	-0.089	-0.92	wealth3	-0.221	-3.39***
wealth4	-0.037	-0.37	wealth4	-0.168	-2.53**
wealth5	0.239	2.13**	wealth5	-0.158	-2.08**
DIL	-0.755	-10.08***	DIL	-0.397	-7.86***
MIL	-0.049	-0.35	MIL	-0.011	-0.12
belongcredit	0.087	1	belongcredit	0.261	4.45***
otherorg	0.112	1.38	otherorg	0.348	6.3***
work	0.619	8.08***	work	0.700	13.46***
rural	-0.392	-5.35***	rural	-0.643	-12.94***
Hindu	-0.058	-0.63	Hindu	-0.061	-0.98
other_rel	0.646	2.38**	other_rel	1.773	9.45***
Sylhet	-0.529	-5.35***	Chittagong	-0.038	-0.65
_cons	1.149	3.29***	Sylhet	-0.124	-1.71*
			Khulna		0.465***
_se	2.728	(Ancillary parameter)	Rajashahi		-0.082
			_cons		0.687**
			se	_1.838533	(Ancillary parameter)

Note: P<0.1 ** P<0.05 *** p<0.01, DIL= daughter-in-law, MIL= mother-in-law

Agency:

Number of observations = 8,706

LR chi2(23) = 992.60

Prob > chi2 = 0.0000

Log likelihood = -20,858.077 Pseudo R2 = 0.0232

Mobility:

Number of observations = 8,704

LR chi2(26) = 2,040.09

Prob > chi2 = 0.0000

Log likelihood = -17,428.529

Pseudo R2 = 0.0553

ANNEX F: FERTILITY

This annex presents analysis on fertility trends and determinants in the 1990s and strategies for achieving continued fertility decline to accelerate achievement of a stable population. The first issue addressed is that of fertility trends in the 1990s. This issue is of policy importance, since there is a widespread perception that fertility decline leveled off in the 1990s, suggesting that the hitherto successful family planning program had run out of steam. But it is shown here that fertility decline continued throughout the 1990s, albeit at a slower rate than previously.

The regression model results feed directly into strategy formulation for maintaining fertility decline. First, the results clearly indicate that son preference is a barrier to reducing fertility. Experience in Bangladesh and elsewhere shows that son preference exerts itself more strongly as fertility falls, so that policies to address the problem, while not obvious, are required. The regression results show that, contrary to normal international patterns, age at marriage has limited potential to reduce fertility, and why this is so is discussed. While desirable for other reasons, strategies to raise the age at marriage will have, at best, limited impact on fertility. It is proposed instead that attention should focus on high-fertility households.

What Has Been Happening to Fertility?

Following the rapid decline during the 1980s, DHS data suggest that fertility leveled off in the 1990s, declining slightly from 3.4 to 3.3 between 1993 and 1996, and then remaining at that level until 1999.

It has been suggested that this slackening pace of fertility reduction comes from a tempo effect. Suppose that during the 1980s, women began postponing births, while maintaining the number of children they had at the same level.

Cohort fertility—the average number of children born to women of a particular age—would remain unchanged. But the total fertility rate is calculated using the synthetic cohort method, which is an average of the age-specific fertility rates of women currently alive. Hence, TFR will temporarily fall as a result of postponed births (because of the reduction in the age-specific fertility rate among the women postponing births), only to return to its original level when these women give birth in the next period (see box F.1). It is plausible that such a tempo effect in Bangladesh accelerated fertility decline in the 1980s, but served to counteract a continuing underlying decline in the early 1990s. If this argument is correct, then the fact that TFR stagnated rather than rose in the 1990s indeed implies that there were other changes taking place to exert a downward pressure on the TFR.

However, while there is evidence of such a tempo effect, there is no need to rely on that to explain the claimed fertility plateau for a number of reasons. First, it has not been usual to rely on direct estimates of fertility to estimate fertility in the past, so it is not clear why they should be the reference point for the 1990s. Second, even if the direct estimates are used, there are reasons to believe that fertility was underestimated in 1993/94 on account of displacement in birth reporting. But, third, if indirect estimates are used, they show both that the direct estimates underestimate fertility (particularly in 1993/94, reinforcing the previous point) and that fertility did continue to decline in the 1990s. These arguments are considered in turn.

Direct Versus Indirect Estimates

Table F.1 summarizes direct fertility estimates from various surveys, together with indirect es-

BOX F.1**The Tempo Effect from Birth Spacing****Number of Births per 100,000 Women**

	Period 1	Period 2	Period 3	Period 4
15–19	40,000	40,000	20,000	20,000
20–24	50,000	50,000	50,000	70,000
25–29	20,000	20,000	20,000	20,000
30–34	8,000	8,000	8,000	8,000
35–39	5,000	5,000	5,000	5,000
40–44	1,200	1,200	1,200	1,200
45–49	400	400	400	400

Age-Specific Fertility Rate

	1	2	3	4
15–19	0.40	0.40	0.20	0.20
20–24	0.50	0.50	0.50	0.70
25–29	0.20	0.20	0.20	0.20
30–34	0.08	0.08	0.08	0.08
35–39	0.05	0.05	0.05	0.05
40–44	0.01	0.01	0.01	0.01
45–49	0.00	0.00	0.00	0.00
TFR	6.23	6.23	5.23	6.23

The table shows the number of births per 100,000 women by five-year age cohorts. These numbers are used to calculate the age-specific fertility rates shown in the bottom part of the table.

The fertility rate is the sum of these age-specific rates (multiplied by five, as each rate covers a five-year period). This is different from the cohort fertility rate, which is the sum of the age-specific rates for an age cohort across its reproductive years. For example, Cohort A is aged 15–19 in period 1, 20–24 in period 2, and so on.

In the example shown, the cohort aged 15–19 in period 3 decides to postpone 2,000 births from age 15–19 until they are aged 20–24. There is no change in the number of children they will have over their life span, so the cohort fertility rate is unchanged.

But the fertility rate is not measured in this way (as it could only be observed for women who have completed their child bearing years). The total fertility rate temporarily drops in period 3, as the age-specific fertility rate for 15–19s has declined. But in the next period the rate for 20–24s rises, as the postponed births take place.

TABLE F.1**Direct and Indirect Fertility Estimates from Various Sources**

Year	Survey	TFR estimates	
		Direct	Indirect*
1974	BRSFM	4.8	7.3
1975	BFS	5.4	7.4
1983	CPS	4.9	7.0
1985	CPS	4.6	6.5
1989	CPS	4.9	5.9
1989	BFS	4.6	5.4
1991	CPS	4.2	n.a.
1993/94	BDHS	3.4	4.3
1996/97	BDHS	3.3	4.0
1999/2000	BDHS	3.3	3.7

Note: 1974–89 from Cleland and others (1994) and 1993–99 from OED analysis.

timates made using data from these surveys. Two points can be noted. First, the direct estimates are systematically lower than the indirect ones. Second, the trend in fertility decline is far less from the direct estimates than from the indirect ones. For example, both CPS 1983 and 1989 yielded direct TFR estimates of 4.9. But these results have not been used to argue that there was a fertility plateau in the 1980s. To the contrary, this is seen as the period of most rapid decline.

Underestimation of TFR in DHS 1993/94

The argument that the direct estimate of TFR in the 1993/94 DHS was an underestimate is based on an apparent displacement of births, with births that were reported as taking place more than three years ago, actually being more recent. The evidence for this phenomenon comes from comparing the number of births reported in the

three years preceding the survey with the number reported for the three years before that (that is, 4-6 years before the survey). Since they cover similar periods, the ratio of these two amounts will be one, except that declining fertility or postponed births raise it above one.

Table F.2 shows this analysis. The ratio for the 1993/94 survey is greatly in excess of that for the other surveys. In that survey, 26 percent more births were reported for 4-6 years before the survey, compared with those reported for within 3 years before the survey. This ratio compares with just 4 percent for 1996/97 and a 5 percent reduction in 1999/2000. Yet comparing the ratio for five-year periods there is little difference between the 1993/94 and 1996/96 surveys. These data thus show that in 1993/94 children under three were being reported as being older. The effect of this birth displacement is exactly the same postponement of births shown in box F.1. That is, it will reduce reported fertility. But since, in this case, it is believed that births are being incorrectly attributed to a later period, this finding represents a downward bias of the direct fertility estimate.


TABLE F.2			Number of Births in Selected Periods Before DHS			
	Three-year intervals		Ratio	Five-year intervals		Ratio
	Births in X years preceding the survey			Births in X years preceding the survey		
	X = 3 years	X = 4–6 years		X = 5 years	X = 6–10 years	
1993/94	3,800	4,795	1.26	6,947	8,226	1.18
1996/97	3,655	3,799	1.04	6,187	7,102	1.15
1999/2000	4,167	3,940	0.95	6,852	6,864	1.00

Indirect Fertility Measures

Appendix F.1 presents indirect estimates of fertility using three different approaches, using BFS 1989 and the three DHS. One set of estimates are also made using 1991 and 2001 census data. Figure F.1 summarizes the results.

The data shown here confirm that direct methods overestimate fertility. This gap is largest for 1993/94, which supports the argument that birth displacement exacerbated un-

FIGURE F.1 Indirect Estimates of Total Fertility Rate



derestimation in that year. The results also show that fertility continued to decline during the 1990s, albeit at a slower rate than had been the case from 1989-93.

Summary

Three arguments have been presented to counter the commonly held view that there was a fertility plateau in Bangladesh in the 1990s. This finding matters, since the perceived plateau gave weight to the view that family planning services were becoming moribund and could not make additional

inroads into fertility. That fertility was being reduced points to the continuing success of these programs. Nonetheless, it is the case that the rate of fertility reduction was slowing down, suggesting that thought does need be given to strategy for attaining the goal of a stable population.

Modeling Fertility Determinants

Table F.3 presents the results of regression models of two dependent variables that proxy for fertility: (1) whether a woman desires to have more children and (2) whether a woman has had a

TABLE F.3

Logistic Regression Results of Fertility Proxies

	Dependent variable		
	Desire more children	Desire more children	Birth in the last three years
Current age >= 25 years	-0.84***	-0.84***	-1.81***
Age at first marriage >= 15 years	0.25***	0.25***	0.53***
Division (Ref Sylhet)			
Barisal	-0.72***	-0.72***	-0.34***
Chittagong	-0.09	-0.09	-0.1*
Dhaka	-0.64***	-0.65***	-0.33***
Khulna	-1.00***	-1.00***	-0.54***
Rajshahi	-0.912***	-0.92***	-0.47***
Rural (ref urban)	0.28***	0.31***	0.01
Education (ref No education)			
1-5	0.08	0.06	-0.05
6-9	-0.00	-0.05	0.09**
10+	-0.13	-0.19**	0.22***
Non-Muslim	-0.35***	-0.356***	-0.23***
Number of Living Children (Ref None)			
1	-1.62***	-1.62***	
2	-3.71***	-3.7***	
3+	-4.82***	-4.81***	
All living children are females			0.46***
A living son			-0.26***
Number of children* all living children female			-0.19***
Male child died* all living children female			0.04
Respondent currently working	-0.18***	-0.19***	-0.21***
Watch television every week	-0.14**		-0.12***
Listen radio every week	-0.06		-0.08***
Access to electricity	0.02	-0.043	-0.23***
Survey (Ref BDHS 93)			
BDHS 96	-0.02	-0.03	-0.05
BDHS 99	-0.05	-0.06	0.03
Constant	4.07	4.03	-0.1
-2 Log likelihood	16,763	16,785	
Model chi-square	11,158	11,171	

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

birth in the last three years. Some of the expected patterns emerge here: all regions have lower fertility than Sylhet, non-Muslims have lower fertility, women with more children are less likely to desire another child, and older women are less likely to have had a child in the last three years or to desire another. Working women have lower fertility, which may reflect the effect of women's agency. Some of the other results require more comment.

First, the age at marriage is significantly positively related to the two fertility measures, a result that is robust to different model specifications. This finding is contrary to the conventional wisdom that increasing age at marriage is a driver of lower fertility, as indeed has been the case in many countries (including, quite probably, Bangladesh in the past). However, as discussed in more detail below, since fertility is already low, there is limited possibility for increasing age at marriage to reduce fertility—the effect is rather to postpone births (which will, as explained above, create a tempo effect that temporarily accelerates the rate of fertility decline). Hence, women who have married later are more likely to want more children and to have had a child in the last three years.

Second, the level of education has the expected effect on the desire for more children—more educated women are less likely to want another child. But there is the opposite relationship for women who have had a child in the last three years—those women having reached at least grade 6 are more likely to have had a child in the last three years, and those at grade 10 and above even more likely to have done so. This result once again reflects the marriage postponement effect—women attending school after the onset of their reproductive years are postponing births rather than having fewer than the (already low) average.

Third, there is considerable evidence of son preference. This preference is picked up by the four interactive variables used in the final regression. A woman is more likely to have had a child in the last four years if all her children are girls (although this effect is mitigated somewhat if she has a lot of them), and less likely to if she already has a son.

Fourth, access to electricity has an unexpected, though insignificant, positive impact on the desire to have more children. However, the coefficient becomes negative (though still insignificant) once the media variables are removed. This result shows that the effect electricity has on lowering fertility is by facilitating access to media, notably TV, which has a significant fertility-reducing effect when included. This variable is thus a measure of the impact of IEC through one specific channel.

Finally, it might be expected that the survey dummy coefficients would be negative, showing an “autonomous” downward trend in fertility by factors not included in the survey. This pattern is present, though not significant, for the “desire more children regression.” But the result does not appear for the child in last three years result. Rather than evidence of such a downward trend, this result reinforces the argument of birth displacement in the 1993/94 survey.

Characteristics of High-Fertility Women

Although fertility has fallen, there remain a significant number of women bearing high numbers of children. The following groups of women were defined as having “high fertility”: (1) those aged 35 or older with five or more children; (2) those aged 35 or older with four or more children; (3) those aged 25 or older with four or more children; and (4) those aged 25 or older with four or more children, and who want more children.

Age 35 was chosen as the cutoff age after which very few (around 3 percent) of the births take place. The problem with this cutoff is that it cannot capture fertility behavior of women currently in their reproductive years. Hence, the lower cutoff of 25 was also used.

The percentage of women in the high-fertility category using the various definitions is shown in table F.4. Except under the fourth definition, a significant minority of the population is identified as having “high fertility.” The fact that many women with four children do not want more children, but many will go on to have them, represents either unmet demand for contraception, incorrect usage, contraceptive failure, or that it is not solely the woman's decision. By all defini-

TABLE F.4**Women in High Fertility Categories by Year and Definition (percent)**

Year	1	2	3	4
1999	17.2	22.4	34.1	1.8
1996	18.2	22.3	37.3	2.0
1993	23.3	29.0	39.2	2.4

Note: See text for definitions.





tions there is a clear drop in women in the high-fertility category between 1993 and 1999.

Figure F.2 shows the relationship between high fertility and selected characteristics from the 1999 DHS data. Although there are variations by region, religion, and wealth, these are not that marked, in that there is a substantial proportion of high-fertility women in each group. There is a slight deterioration in the share by wealth quintile (though by other definitions and in other years there is an inverted U, with the

highest share in the middle quintiles). The proportion of high-fertility women is highest in Sylhet and lowest in Khulna. There is a more marked pattern for education. Close to half of women with no education have a high level of fertility, compared with just 6 percent of those with higher education.

Strategies for Fertility Reduction

The evidence is clear that Bangladesh has achieved substantial fertility reduction, in excess of what is expected from a country at its level of socioeconomic development by virtue of its effort in family planning (see Annex B). Although fertility reduction has slowed, it is of course necessary to continue to disseminate family planning messages and support contraceptive supply in order to preserve gains to date and seek additional progress. Given the slowdown in fertility reduction, which is not unexpected, what policies are most likely to prove successful for mov-

FIGURE F.2**Correlates of High Fertility**

ing toward the goal of bringing fertility down to the replacement level? It is first argued that raising the age at marriage, while it has other benefits, is unlikely to make much of an impact on fertility. Rather, attention should be paid to the three interrelated areas of high-fertility families, son preference, and continuing reductions in under-five mortality.


Why Raising the Age at Marriage Will Have a Limited Impact on Fertility in Bangladesh

There is a well-established inverse relationship between the age of marriage and fertility, with an increasing age at marriage often being a driver of fertility reduction. DHS data from around the world, shown in figure F.3, bear this out. There is a clear negative relationship: the fitted regression line suggests that increasing the age of marriage by one year reduces fertility by one-third of a child.¹ However, Bangladesh is a clear outlier in this figure, with the three DHS surveys for that country laying in the lower left quadrant. Bangladesh has the lowest age at marriage of all countries shown, but well below-average fertility, with an age at first marriage typical for a country in which the median age at marriage is over 20, rather than 14-15. This evidence both supports the view that Bangladesh has already, through other means (that is, family planning programs), reaped the gains to be had from reducing the age at marriage, and suggests that there may be rather limited fertility gains from increasing the age at marriage. This latter point is further supported by the pattern displayed by data, with the data points being more or less horizontal. This pattern is in contrast to other countries, for example Egypt, Peru, and Senegal, for which the line joining the data points has the expected downward slope.

Figures F.4(a)-(c) show the age at marriage and at first birth, and the lag between them. During a short period in the 1990s there was a surprisingly large change in the age at marriage (see also Annex E). For example, in 1993, 26 percent of the women interviewed had been married by age 12, compared with just 14 percent in 1999.² These changes are shown by a shift of the cumulative distribution in figure F.4(a) to the right. But the age at first birth, in figure F.4(b),

FIGURE F.3


Bangladesh Is an Exception to the Normal Relationship Between Age at Marriage and Fertility



Source: ORC Macro 2004. MEASURE DHS STATcompiler. <http://www.measuredhs.com>, November 2 2004.

shows no such shift. Women are getting married later, but not having their first child at an older age than before. Thus, it should be the case that the lag between getting married and having the first child has shortened, which figure F.4(c) shows to be the case: in 1999, 17 percent of women had their first child within a year of marriage, compared to just 9 percent doing so in 1993.

Figure F.4(d), which shows the scatter plot of age at first birth against age at marriage, digs a bit deeper into what is going on here. The plot is fitted with a locally weighted regression line, which shows there is no relationship between age of marriage and age at first birth (that is, the regression line is flat) up to about 13 years of age.³ Close to 20 percent of ever-married women in the 1990s had married by 12, and 40 percent by 13. But, as the graph shows, increasing the age at marriage within this age range will have no effect at all on the age at first birth. Girls getting married below age 13 are generally considered to be below reproductive age and only exceptionally bear children (only 4 percent of births are to girls aged 13 and younger). Second, the

FIGURE F.4 Changes to Age at First Marriage and First Birth

Source: Calculated from DHS 1993 and 1999.

slope of the line, which is constant after the structural break at around 13 years, is less than 1 (it is 0.8, see footnote 3). Hence, an increase in the age of marriage of one year leads to a one less a one year increase. This means that as the age at marriage increases, the absolute gap between marriage and first birth declines. The regression results suggest this gap is on average 2.7 years for a woman marrying at 15, but only 1.8 years for one marrying at 20. This is precisely the phenomenon shown in panels (a) and (c) of figure F.4—as the age at marriage increases the interval to first birth declines.

The point being made above is that delaying marriage only has a limited effect of delaying the

onset of a woman's reproductive life.⁴ It has, on average, no effect at all for women marrying at 13 or younger. And it has a muted effect on older women.

It is usually argued that delaying marriage reduces fertility, since a woman starts reproduction later, and so has less time to achieve her reproductive potential. But in Bangladesh the majority of women are having three or four children during their reproductive lives. This is quite possible to achieve if childbearing starts at 20 rather than 15. Hence, even if delaying marriage delayed first births, which it does to only a limited extent, it still would not affect the total number of children they have during their lifetime.

Strategies to Reduce Fertility: Addressing High-Fertility Households, Son Preference, and High Mortality

As shown above, one-third of all women aged 25 have four or more children. There seems to be greater potential in persuading these families to have 3 children than in persuading those having 2-3 children to have fewer. That son preference is a barrier to fertility decline in Bangladesh has been argued elsewhere (Chowdury and Bairagi 1990; Bairagi 1996; and Khan and Khanum 2000), and is supported in the regression results presented above. Reducing son preference would hence have a beneficial impact on fertility. This is easier said than done, however: son preference is a deeply ingrained cultural tradition, ex-

acerbated by the spread and escalation of dowry.⁵ Strategies that may be adopted include Behavior-Change Communication, particularly including work with religious authorities, expanding female education, and employment and income-generating opportunities for women. These last two are areas of already sizeable programs in Bangladesh.

Bangladesh has followed international norms in that fertility decline has been linked to declining under-five mortality. There remain pockets of high mortality, especially in Sylhet where use of health services is low. Concerted efforts to increase immunization coverage and other interventions to reduce mortality can be expected to have an impact on fertility in these areas.

APPENDIX F.1: Indirect Estimates of Fertility in the 1990s

This appendix provides indirect estimates of fertility in Bangladesh for the 1990s using three methods: Rele method, Proximate Determinant method (Bongaarts decomposition), and the regression method. The data for the analysis are taken from BFS 1989 and the three DHS surveys, and, for one set of calculations, census data.

Rele Method

This method infers fertility levels from child-woman ratios (CWR), utilizing the close linear relationship between the CWR and the gross reproduction rate (GRR)¹ shown to exist at any given level of mortality (measured by life expectancy, e^0_0). Rele (1976) provides the parameters required for implementation of the approach.² Using these coefficients, GRR is estimated from CWR using suitable values of e^0_0 in a given population and then applying the sex-ratio at birth estimate to obtain TFR from GRR.

The method uses two types of CWR in estimating GRR—children aged 0-4 years divided by women aged 15-49 (CWR1), and children aged 5-9 divided by women aged 20-54 (CWR2). The e^0_0 value used corresponds to the period when the children in the CWR were born. The fertility estimates also refer to the same period viz., estimates obtained through using CWR1 produces an estimate for five years preceding the census/survey and that with CWR2 produces for 5 to 10 years preceding that.

This method has certain strengths and weaknesses. Among the former, the estimated fertility is insensitive to the errors in e^0_0 , so that an approximate value of e^0_0 gives a fairly accurate estimate of fertility. Among the weaknesses, the derived fertility estimates depend on the accuracy of CWR. In developing countries, due to errors in age reporting, notably exaggeration of age at young ages and underreporting of very young children, the reported age distributions are often distorted. To overcome this problem it was initially suggested to average the two rates

obtained from the two CWRs and arrive at a decadal estimate (Rele 1976). Also, one may use smoothed age data. But this procedure has some disadvantages as estimates with the smoothed age distribution are influenced by the particular method used in smoothing the age data.

Rele (1987) refines the method for situations more typical in some developing countries, including India. The refined method requires two or more successive censuses preferably with a 10-year gap between them and use of weights³ derived on the basis of extent of under-enumeration/over-enumeration of two types of CWRs to adjust preliminary estimates made under this method. If weights cannot be computed very accurately for some populations, it may not affect much the final or adjusted estimate for GRR/TFR. In other words, the estimates are not that sensitive to some variation in the weights.

Estimates for TFR are obtained using both survey and census data. The survey data are those from three BDHS conducted in the 1990s and the BFS-1989. In the case of population census, data from only 1991 and 2001⁴ census were used. The value of life expectancy at birth (e^0_0) is assumed on the basis BBS estimates for the same. They are given in tables F1.1 and F1.2, displaying the estimated TFR as obtained under this method. Un-smoothed age distribution data have been used for estimating TFR under this method.

TFR estimates applying the Rele method using survey data with overlapping periods are shown in table F1.2. Given the overlapping periods, a final estimate has been derived using the weights suggested in the refined Rele method (1987).

Proximate Determinant Method

Davis and Blake (1956) were the first investigators to develop a framework outlining the intermediate fertility variables. That seminal work

identified 11 intermediate variables through which fertility is affected in a population. Later on, Bongaarts (1978) revised this framework incorporating only 8 of them, which he termed proximate determinants of fertility. As he suggested, each of these eight variables directly influences the fertility and they together determine the level of fertility in a population. However, using the data from both developed and developing countries, Bongaarts and Potter (1983) observed that 96 percent of the variance in total fertility rates could be explained by four principal proximate determinants: marriage, contraception, induced abortion, and lactational infecundability. The other four, which can be disregarded, are: frequency of intercourse, sterility, spontaneous intrauterine mortality, and duration of the fertile period. These last four factors all relate to the natural fertility.

The key contribution of the Bongaarts framework is to set up the procedures for parceling out the effects on fertility of different proximate variables. The forces of each of the intermediate variables in the Bongaarts framework is measured by an index 'C' that varies from zero to one. When it is zero, none of the potential for fertility is translated into births. When it is 1.0, the controls exert no restraining effect whatsoever.

In Bongaarts' framework, the total fertility rate (TFR) in a population is expressed as the product of four indexes measuring the fertility inhibiting effect of four principal intermediate variables and the total fecundity rate (TF). The total fecundity rate is the average number of live births expected among women who during their entire reproductive period remain married, do not use contraception, do not have any induced abortion, and do not breastfeed their children. The TF does not vary much between populations, lying between 13 and 17 births per woman with a standard value of 15.3 (Bongaarts and Potter 1983).

The following equation summarizes the basic structure of the Bongaarts model by relating the fertility measures to the proximate determinants:

$$TFR = Cm \times Cc \times Ci \times Ca \times TF$$

Where Cm is the index of marriage, Cc is the index of contraception, Ca is the index of induced abortion, and Ci is the index of lactational

TABLE F 1. 1		Rele Estimates of GRR and TFR Obtained from Survey Data			
Period	e^0_0	Preliminary estimates		Final estimates	
		GRR	TFR	GRR	TFR
BFS-1989					
1979-84	50	3.26	6.69	3.09	6.30
1984-89	53	2.56	5.25		
BDHS-1993/94					
1984-89	53	2.99	6.12	2.86*	5.86*
1989-94	56	2.13	4.37		
BDHS-1996/97					
1987-92	55	2.73	5.61	2.55**	5.22**
1992-97	57	1.92	3.94		
BDHS-1999/2000					
1990-95	57	2.32	4.76	2.20***	4.51***
1995-2000	60	1.83	3.76	1.85	3.78

Notes: * for 1984-89; ** for 1987/89-92/94; and *** for 1990/92-95/97.

infecundability. Each index measures the extent to which fertility is reduced from the maximal levels by specified proximate determinants. The indexes are measured as follows:

$$C_m = \frac{m(a) \times g(a)}{\sum g(a)}.$$

Where $m(a)$ = age specific proportions currently married among female and $g(a)$ = age specific marital fertility rate,

$$Cc = 1 - 1.08 \times u \times e.$$

Where u = proportion currently using contraception among married women of reproductive age and e = average use effectiveness of contraception,

TABLE F 1. 2		Rele Estimates of GRR and TFR Obtained from Census Data			
Period	e^0_0	Preliminary estimates		Final estimates	
		GRR	TFR	GRR	TFR
1981-86	51.0	3.44	7.04	3.33	6.82
1986-91	54.0	2.93	6.01	2.91	5.97
1991-96	57.0	2.37	4.86	2.38	4.89
1996-01	60.0	1.91	3.91	1.90	3.89

$$C_i = \frac{20}{18.5 + i}.$$

Where i = average duration of post-partum infecundity caused by breastfeeding or post-partum abstinence,

$$C_a = \frac{TFR}{TFR + .4 \times (1 + u) \times TA}.$$

Where TA is total abortion rate.

C_m , C_c , and C_i are calculated for different years using the information from three BDHS of the 1990s and BFS of 1989. Data for C_a are not available, hence, the value of it has been assumed 1.0 all through. This assumption most likely produces an upward estimation of TFR for population that may practice some degree of induced abortion in Bangladesh (Begum 2003), although the extent is unknown. Such practice may also be on the rise over the years.

In estimating C_c the knowledge about the contraceptive use-effectiveness is required. There is no uniform view about this. Using the information of 1993/94 BDHS, Islam and others (1996) estimated this to be 0.93, but using village-level statistics (Matlab) Bairagi and other (1996) estimated a value of 0.81. However, the Matlab estimate is more akin to the experiences of the developing countries (Ross and Frankenberg 1993). Here C_c is estimated using both sources (table F1.3) giving two C_c estimates and, consequently, two TFR estimates, which are averaged to arrive at a single figure for TFR.

However, the BDHS have shown an increase in the discontinuation rate of contraceptive methods over the 1990s. Hence, the C_c estimates for 1996/97 and 1990-2000 using contraceptive use-effectiveness from 1993/94 or the period around that may overestimate the contraceptive effect on TFR in the former two periods. Thus, this may at least partly offset the opposing effect of induced abortion noted above.

The C_i has been estimated using direct estimates of the post-partum amenorrhea period by different surveys (table F1.4). This has been falling over time, exerting an upward pressure on fertility.

The values of C_m , C_c , and C_i as estimated for different periods are presented in table F1.5, together with the corresponding TFR. TFR is shown to have continued to fall during the 1990s, with the bulk of the decline coming from increased contraceptive usage. As noted, C_i exerted a countervailing tendency.

Regression Method

From international comparisons of the family planning program performance and fertility change, Blanc (1990) has derived a set of regression equations that can be used to predict TFR from the information on CPR. Blanc proposed five regression equations. However, to arrive at a single value of TFR the average of five TFR values is suggested. Past users of this method have argued that this method is likely to produce an underestimation of TFR in a country such as Bangladesh (Kantner and Frankenberg 1986; Kantner and Noor 1992), and suggested that estimates obtained through this procedure may be treated as the lower-bound estimates. The TFR estimation formulae under this method are as follows; to arrive at a single value for the TFR, five different estimates obtained under five different formulae are average out:

$$TFR = 7.30 - (6.42 * CPR)$$

$$TFR = 6.83 - (6.20 * CPR)$$

$$TFR = 7.38 - (7.20 * CPR)$$

$$TFR = 7.28 - (6.55 * CPR)$$

$$TFR = 7.15 - (6.56 * CPR).$$

The results are presented in the next section, alongside those of the other methods.

TABLE F1.3		Methods-Specific Use— Effectiveness of Contraception	
Method	BDHS 1993/94	Matlab	
Pill	0.96	0.74	
IUD	0.99	0.92	
Injectables	0.98	0.99	
Condom	0.87	0.57	
Sterilization	1.00	1.00	
Periodic abstinence	0.81	0.70	
Withdrawal	0.81	0.70	
Others	0.63	0.70	

Source: Islam, Mamun, and Bairagi 1996.

Estimated TFR

The estimates for TFR as obtained under the different methods are presented in tables F1.6 and F1.7. The Rele and Proximate Determinant methods are mainly relied upon to arrive at an estimate of TFR, as the Regression method is believed to produce an underestimate.

The estimates obtained using census and survey data under the Rele method and those obtained through Proximate Determinant methods are in close agreement, while the Regression method, as expected, produced consistently lower estimates for all the periods. Among the former two, the major discrepancy is for TFR estimates for the early 1980s. The Rele method, by using census data, produced an estimate of 6.82 for the TFR for the 1981-86 period and, using survey data, produced an estimate of 6.30 for the 1979-84 period. Both estimates are obtained using primarily the CWR2, which used the 5-9 age group as the numerator, hence it may be that there was some higher enumeration of this age group in the census, which is less of a problem in the survey data, which are collected with greater care and attention. Both in 1961 and 1974 the population censuses have excessively enumerated the 0-9 age and particularly the 5-9 age (Bangladesh 1961; Begum 1976, 1990).

For the 1986-91 period, the Rele method, using census data, produced an estimate of 6.0 for the TFR and, using survey data, produced an estimate of 5.9 for 1984-89 period. Hence, according to this method, the TFR in the country in the 1987-89 period was around 6 per woman,

TABLE F1.4		Estimated Duration of Post-Partum Amenorrhoea (months)
Period		Duration (months)
1989		11.9
1993/94		11.5
1996/97		10.9
1999-2000		9.5

Source: Survey reports.

while the Proximate Determinant method estimated the same level for the year 1989. Thus, on the basis of Rele and Proximate Determinants method, we can conclude that the fertility level in Bangladesh before 1990 was around 6 per woman. A similar result was found by Cleland and others (1994), who report a TFR of 5.86 for 1988-89 using CPS data, and Islam and others (1996), who report a TFR of 5.83 using BFS. As a crude measure, the average number of children born to women aged 35 or over in DHS 1992/93 was 5.9, which fits with these estimates.

The Rele method, using census data for the first half of 1990s viz., for 1991-96, produced an estimate of 4.9, while using survey data produced an estimate of 4.51 for 1990/92-1995/97 period. The Proximate Determinant method produced an estimate of 4.5 for 1993/94, which corresponds roughly to the mid-period of the above two periods noted for the Rele method. From these rates, it appears the TFR in 1993/94 in Bangladesh was around 4.5.

TABLE F1.5		Estimated Index Values of the Proximate Determinants				
Year	Cm	Cc		Ci	Cm x Cc x Ci*	TFR*
		Using BDHS use-effectiveness	Using Matlab use-effectiveness			
BFS 1989	.826	.695	.730	.658	.387	5.99
BDHS 1993/94	.761	.550	.610	.667	.294	4.50
BDHS 1996/97	.756	.503	.573	.680	.276	4.23
BDHS 1999-2000	.745	.466	.545	.714	.269	4.11
Percent change, 1989-99	-9.8	-32.900	-25.3	8.5	-30.500	-31.40
Share	32.200	95.6*	-27.900	100.000		

Note: * Calculated using average of two Ccs. Growth rate of TFR differs from that of product term due to rounding.

TABLE F1.6**Estimates of TFR Using Rele Method**

Period	Census estimates	Period	Survey estimates
1981–86	6.82	1979–84	6.3
1986–91	5.97	1984–89	5.86
1991–96	4.89	1987–89–1992–94	5.22
1996–2000	3.93	1990–92–1995–97	4.51
		1995–2000	3.78

For 1996/97 the Proximate Determinant method has produced an estimate of 4.23 and, for 1999–2000, 4.11. On the other hand, the Rele method using census data produced an estimate of 3.93 for the second half of 1990s viz., for 1996–2000, which corresponds to the mid-period of 1998, while using survey data produced an estimate of 3.78 for 1995–2000, corresponding to the year 1997. Hence, from these estimates we can conclude that the TFR in the country during 1996/97 was perhaps around 4.2 and that during 1999–2000 it was around 4.

Summary

Two major conclusions can be drawn from this analysis. First, national sample surveys conducted in the country consistently underestimated the true fertility level in the country. According to these estimates, of three BDHS conducted in the country in the 1990s, underestimation of TFR by highest margin has taken place in 1993/94.

Second, as these estimates show, the fertility did decline in the country over the 1990s albeit at a slower rate than that noticed between the late 1980s and early 1990s. For example, TFR declined by about 25 percent between 1989 and 1993/94, while during 1993/94 and 1999/2000 the magnitude of decline has been only about 11 percent.

The dynamic behind the slowing fertility decline in the second half of the 1990s can be discerned from the trend in index values of various proximate determinants over this time (table F1.8). As they suggest, for a big drop in TFR between 1989 and 1993/94, both marriage and contraception played a big role. Contraception use during that period increased 14 percentage points from 31 percent to 45 percent, while the never-married group in the 10–49 age group increased from 20 to 32 percent. Since 1993/94, marriage played only a marginal role in inhibiting fertility. Such role of marriage improved by less than 1 percent during 1993/94–1996/97 and by another 1.5 percent during 1996/97–1999–2000, while during the 1989–1993/94 period alone it improved by 8 percent.

Similarly, during the 1989–1993/94 period, while the fertility inhibiting effect of contraception increased by 19 percent, it improved by a much smaller degree after that. During 1993/94–1996/97 the fertility-inhibiting effect of contraception increased by 7 percent and by 6 percent during 1999–2000. While the lesser degree of increase in the contraception use after 1993/94 compared to 1989–1993/94 may in part be responsible for this, there is also evidence that in the latter part of the 1990s, contraception use relied more on relatively less-effective methods and the use-effectiveness of the method experienced some decline during this period.

Throughout this period the role of post-partum infecundability has been in the opposite direction viz., it favored fertility increase. Not only this, this role increased noticeably over time, offsetting part of the fertility-inhibiting effect exerted by other proximate determinants. During the 1989–1993/94 period the fertility inhibiting effect of *Ci* slackened by 1.4 percent, it slackened by another 1.9 percent during the

TABLE F1.7**Estimates of TFR Using Proximate Determinant Method and Regression Method**

Year	Proximate determinant method			Regression method
	Using <i>Cc</i> estimated from 1993/94 BDHS	Using <i>Cc</i> estimated from Matlab data	Average of two estimates	
1989	5.85	6.14	5.99	5.16
1993/94	4.27	4.74	4.50	4.25
1996/97	3.96	4.51	4.23	3.96
1999–2000	3.79	4.43	4.11	3.65

193/94–1996/97 period and another 5 percent during 1996/97–1999/2000.

The outcome of all these effects has been continuous weakening of collective strength of the fertility-inhibiting effect of various proximate determinants. During the 1989–1993/94 period mentioned above, three proximate determinants viz., *Cm*, *Cc*, and *Ci* together could improve their fertility-depressing effect by another 24 percent. While they could continue improving such an effect throughout the rest of 1990s, they could do so only to lesser degree. During 1993/94–1996/97 such improvement has increased another 6 percent, and during 1996/97 and 1999/2000 it improved by another 2.5 percent.

TABLE F 1.8		Percentage Change in TFR and Different Indices of Proximate Determinants Since 1989			
Period	<i>Cm</i>	<i>Cc</i>*	<i>Ci</i>	Combined effect of <i>Cm</i>, <i>Cc</i>, <i>Ci</i>	TFR
1989–1993/94	–8.0	–18.6	+1.4	–24.0	–17–22
1993/94–1996/97	–0.7	–7.2	+1.9	–6.1	–6–15
1996/97–1999/2000	–1.5	–6.0	+5.0	–2.5	–5–10

* Change for *Cc* has been calculated from average value.

ANNEX G: ANALYSIS OF BINP'S COMMUNITY-BASED NUTRITION COMPONENT

The Community-Based Nutrition Component, the central component of BINP, focused on growth monitoring, combined with nutritional counseling and food supplementation. The impact of the component has been studied in an independent evaluation conducted for the project (Karim and others 2003), by Save the Children in their report *Thin on the Ground* (Save the Children 2003), in an evaluation commissioned by GoB's Implementation, Monitoring and Evaluation Department (Haider and others 2004), and in research carried out under the supervision of Professor Mascie-Taylor at the University of Cambridge, U.K. (Mascie-Taylor 2004). For this report, further analysis has been conducted using existing data, but partly different approaches to those used in these other studies. For this study we have had access to the project evaluation dataset, the data collected by Save the Children, and data from Helen Keller International's Nutritional Surveillance Survey. Thanks are due to the respective agencies for making these data available, and to World Bank staff in the Nutrition Hub for facilitating access to the evaluation dataset.

This annex presents analysis relating to both child nutrition and low birth weight, whose prevalence is intended to be reduced through improved weight gain among pregnant mothers. The annex is structured to reflect the logic of project design. The key assumption behind the Community-Based Nutrition Component is that "bad practices" are responsible for malnutrition in Bangladesh. This point of view was strongly argued in the appraisal document for the project: "behaviors related to feeding of young children have at least as much (if not more) to do with the serious problem of malnutrition in

Bangladesh as poverty and the resultant household food insecurity do" (BINP Staff Appraisal Report, World Bank 1995, para 1.13, p.4). Therefore, changing bad practice to good will bring about nutritional improvements. There are a number of steps in the causal chain behind this approach:¹

- The right people (those making decisions with respect to undernourished children) are targeted with nutritional messages.
- These people participate in project activities, and so are exposed to these messages.
- Exposure leads to acquisition of the desired knowledge.
- Acquisition of the knowledge leads to its adoption (that is, a change in practice)
- The new practices make a substantial impact on nutritional outcomes.

A feeding program for malnourished children and pregnant women was implemented alongside growth monitoring. For this program to work:

- The target groups have to enroll in the program.
- The criteria have to be correctly applied in selecting those to receive supplementary feeding.
- Those selected for supplementary feeding have to attend sessions to receive the food.
- There can be no leakage (such as selling of food supplements) or substitution (reducing other food intake).
- The food has to be of sufficient quantity and quality to have a noticeable impact on nutritional status.

Project Coverage and Targeting

Participation in Growth Monitoring for Children

Under BINP, growth monitoring sessions are held monthly at which the Community Nutrition Promoter (CNP) weighs all children at the Community Nutrition Center (CNC).² Their weight is marked against age on a growth monitoring chart by the CNP. Mothers of children were asked whether or not they attended the growth monitoring sessions with their child, with possible responses of never, sometimes, and almost always at midterm, and never, sometimes, almost always, and always at endline. The results show that around 85 percent of mothers reply that they always participate (table G.1). There is little change in the overall average between endline and baseline, but considerable variation among thanas, with participation consistently lowest in Rajnagar.

A simple bivariate analysis shows that more educated mothers are more likely to attend than less educated, though this difference is no longer significant by the endline (table G.2). However, at midterm it appears that the poorer mothers (lower housing quality) are more likely to participate, but this pattern is reversed by endline. Multivariate analysis of participation was used to disentangle these effects, using both a logit specification of participate/don't participate and a multinomial logit of the level of participation.

The multivariate results, which are very similar for midterm and endline, show the following:

- There is no sex bias in participation. Participation has a non-linear relationship with the child's age, with those aged 9-10 months most likely to attend.
- Both younger mothers and daughters-in-law are less likely to attend (daughter-in-law meaning here a woman living with her mother-in-law). The daughter-in-law dummy alone is significantly negative (results not shown), but is no longer so when an interactive term is included for the daughter-in-law dummy and the dummy for the two more conservative thanas, Rajnagar and Sharasti.³ The significant negative coefficient on this interactive term shows the effect of restrictions on female mobility in attending growth monitoring sessions, although they only have an effect in two of the six thanas. If the mother is employed outside the household, which is positively associated with women's agency (Annex E), she is more likely to participate. It is also the case that Hindu women are more likely to participate than Muslim women.
- Qualitative analysis shows that remoteness (travel time) is also a constraint on participation. There is no direct measure of this variable in the dataset, but it is likely that type of water supply is a reasonable proxy for this variable (it is difficult to imagine a direct effect of this variable), the results confirming that inferior forms of water supply (indicating remoteness) significantly reduce participation.⁴
- The results confirm that the most educated women are significantly less likely to partici-

TABLE G.1

Participation in Growth Monitoring by Thana (percent)

Thana	Midterm			Endline		
	Never	Occasionally	Regularly	Never	Occasionally	Regularly ^a
Banaripara	2	3	94	6	4	91
Faridpur	8	13	80	4	6	90
Gabtohi	2	5	93	2	3	95
Mohammadpur	4	8	88	4	3	94
Rajnagar	15	15	70	16	9	75
Shahrasti	2	12	87	9	7	83
Total	5	9	85	7	5	88

a. Combines "almost always" and "always."

Source: BINP evaluation dataset.

pate. There is no income-bias in participation in the multivariate analysis, this result being robust to several different specifications (that is, models that do not have other variables correlated to income, such as water and sanitation and parent's education). However, at midterm the less poor are less likely to attend, based on both the housing quality index and the categorical income variable (based on the enumerator's assessment). By contrast, at endline those with better housing are more likely to participate, but women in households with sizeable landholdings less likely to do so (reflecting a time constraint for such women, which is shown to be important when it comes to putting knowledge into practice).

Of these factors, it is the more conservative nature of Rajnagar and Shahrasti that appears to account for lower participation in these thanas. The coefficients on the other variables are not such as to have such a large difference, and the differences in these explanatory characteristics between the thanas are not so great (see figure G.1).

The logic of the project is that attendance at growth monitoring provides a context for nutritional counseling, especially for women with malnourished children. The monitoring sessions are not a good setting for such counseling, but BINP provides other opportunities for this, including group sessions. As shown in table G.4, fewer women have participated in nutrition discussions than participate in growth monitoring. Save the Children data also show participation in nutritional counseling at just under 50 percent.

Targeting of Children's Supplementary Feeding

Growth monitoring is used to provide supplementary feeding to children who are either faltering in their growth or severely malnourished (third-degree level of malnutrition by the Gomez classification, which corresponds to approximately less than -4 SDs from the WAZ reference median). Table G.5 presents the official screening criteria, though field experience suggests that the actual criteria applied vary from thana to thana. The majority of the children ad-

TABLE G.2		Rele Estimates of GRR and TFR Obtained from Census Data		
Comparison for calculation of t-test	Non-participant	Occasional participants	Regular participants	
	Compared to occasional participant	Compared to regular participant	Compared to non-participant	
Midterm				
Mother's education	2.1	2.2***	2.8**	
House index	4.5***	4.1**	4.2***	
Endline				
Mother's education	4.0	4.1	4.1	
House index	4.5	4.7	4.7*	

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

TABLE G.3 (a)		Participation in Growth Monitoring: Midterm				
Variable	Logit		Multinomial logit			
	Participants		Occasional participants		Regular participants	
	Coeff.	t stat.	Coeff.	t stat.	Coeff.	t stat.
Child sex	-0.02	-0.12	0.12	0.59	-0.04	-0.26
Child age	0.19	1.77*	0.20	1.62*	0.20	1.79*
Child age squared	-0.01	-1.85*	-0.01	-1.39	-0.01	-1.92*
Mother's age	-0.04	-3.08***	-0.05	-2.60***	-0.04	-3.04**
Female-headed household	0.00	-0.01	-0.21	-0.39	0.04	0.08
Mother works	1.34	1.23	0.15	0.11	1.45	1.33
Daughter-in-law	-0.19	-0.54	-0.15	-0.36	-0.20	-0.56
Daughter-in-law in Rajnagar and Shahrasti	-1.13	-2.15**	-0.81	-1.37	-1.19	-2.22**
Widow	-0.45	-0.49	0.04	0.04	-0.56	-0.61
Number of pregnancies	0.18	1.63*	0.01	0.09	0.21	1.82*
Primary education	0.13	0.52	0.05	0.19	0.14	0.57
Secondary education	0.42	1.35	0.27	0.79	0.45	1.42
Higher education	-0.41	-0.96	-0.49	-0.91	-0.40	-0.94
Per capita income	0.00	-1.68*	0.00	-1.76*	0.00	-1.67*
House index	-0.22	-2.46***	-0.35	-3.26***	-0.20	-2.20**
No drinking water	-0.98	-2.12**	-0.63	-1.30	-1.08	-2.23**
No latrine	-0.01	-0.04	-0.10	-0.34	0.00	0.01
Faridpur	-1.64	-3.91***	-0.32	-0.74	-1.76	-4.03***
Gabtolli	-0.21	-0.35	0.23	0.48	-0.23	-0.38
Mohammadpur	-0.76	-1.84*	0.19	0.43	-0.82	-1.96***
Rajnagar and Shahrasti	-1.21	-2.96***	0.08	0.19	-1.32	-3.20***
Constant	4.55	4.58***	2.07	1.84*	4.38	4.35***
Pseudo R-squared		0.12				0.07
Observations			3,541			3,541

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

TABLE G.3 (b)**Participation in Growth Monitoring: Endline**

Variable	Logit		Multinomial logit			
	Participants		Occasional participants		Regular participants	
	Coeff.	t stat.	Coeff.	t stat.	Coeff.	t stat.
Child sex	-0.02	-0.12	0.12	0.59	-0.04	-0.26
Child's age	0.01	0.29	-0.01	-0.53	0.01	0.39
Child's sex	-0.21	-1.14	-0.51	-2.01**	-0.18	-0.98
Number of pregnancies	-0.03	-0.30	-0.07	-0.54	-0.03	-0.28
Mother's age	0.04	1.46	0.02	0.56	0.04	1.52
Mother's work	1.05	1.73*	1.14	1.65*	1.08	1.77*
Primary education	0.25	1.01	0.21	0.63	0.25	1.03
Secondary education	0.36	1.19	0.46	1.18	0.37	1.22
Higher education	-0.68	-1.80*	-0.87	-1.51	-0.67	-1.76*
Hindu household	0.83	2.22**	0.46	0.97	0.85	2.27**
House index	0.21	1.88*	0.19	1.34	0.21	1.87*
Land owned	0.00	-2.10**	0.00	-1.63*	0.00	-2.02**
No drinking water	-0.50	-2.16**	-0.64	-2.08**	-0.50	-2.15**
No latrine	-0.25	-1.09	-0.34	-1.10	-0.25	-1.07
Faridpur	0.44	1.06	0.91	1.67*	0.42	1.02
Gabtolli	0.79	1.58	0.56	0.86	0.82	1.64*
Mohammadpur	0.27	0.67	-0.10	-0.17	0.28	0.70
Rajnagar	-0.94	-2.81**	0.35	0.74	-1.04	-3.09***
Shahrasti	-0.45	-1.51	0.45	1.02	-0.50	-1.68*
Constant	1.58	1.78*	-0.28	-0.24	1.42	1.60
Pseudo R-square		0.08				0.07
Observations			2538			2538

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

mitted into supplementary feeding qualify by virtue of growth faltering; analysis of register data collected by Save the Children show only 5 percent of supplementary feeding participants to be severely malnourished (WAZ < -4 SDs), though another 60 percent are malnourished by the conventional criterion of WAZ < -2 SDs (table G.6).

However, not all children enrolled in supplementary feeding appear to be eligible (table G.7).⁵ A likely explanation for this is the inability of all CNPs to accurately interpret the growth charts, a finding from a limited sample encountered during OED's fieldwork.⁶

Targeting of Activities for Pregnant Women

Under the project, pregnant women are meant to be weighed monthly and screened for malnutrition. Women below a certain cutoff point are


given a food supplement. The first screening should happen as soon as the pregnancy is detected in order to measure the pre-pregnancy weight. Table G.8 shows participation rates in weighing and feeding sessions. As expected, some women participate only in the weighing sessions. Surprisingly, there are also women who participate only in the feeding sessions. Participation in weighing sessions is around 75 percent, and just under half of pregnant women received supplementary feeding. There is little change in participation rates between the two surveys. If the cutoff point used to detect pre-pregnancy malnutrition is a body mass index (BMI) below 18.5, as stated in the project document, then fewer women are receiving supplementary feeding than should be doing so, particularly at the midterm when the proportion of women below this cutoff point is 68 percent. By endline, 47 percent are below this threshold, suggesting few women are excluded if only eligible women participate, but this is doubtful, as is shown below.

Table G.9 shows the percentages of women of different nutritional status that participate in project activities. BMI measurement is from women who were not more than six months pregnant in order to avoid the bias induced by the pregnancy weight gain. This is not a correct indicator of the mother's pre-pregnancy status, but is the best that can be done in the absence of data on pre-pregnancy weight.

Participation in weighing sessions is independent of mothers' nutritional status. Since all women are meant to participate, this is the expected finding if there is no bias in favor of, or against, nutritionally deprived women. However, coverage of supplementary feeding is correctly correlated to nutritional status in both surveys, apparently more so at the endline, although a third of even the most severely malnourished women do not participate. However, over 30 percent of women with normal BMI status also received supplementary feeding. At the endline, 40 percent of the recipients of supplementary feeding were women of normal BMI status. In summary, targeting errors at the endline were around 60 percent for Type I error (missing women who should have benefited) and

FIGURE G.1

Various Factors Affect Women's Participation, but Restrictions on Women's Mobility in More Conservative Areas Are the Most Important



40 percent for Type II error (giving it to women who were not eligible). Improved coverage of poorly nourished women, especially the most severely malnourished, would increase program impact on pregnancy weight gain and, hence, low birth weight.

Knowledge Acquisition and the Knowledge-Practice Gap

The main objective of the project is to improve children's nutritional status by changing the pregnancy, lactation, and feeding practices of mothers. Pregnant women are advised, individually and in group sessions, to take more food and rest during pregnancy in order to avoid low

birth weight. In addition, mothers are advised to give colostrums to the newborn and to practice exclusive breastfeeding of the infant up to the age of 5-6 months. For older children, advice is given on good nutrition. As outlined above there are several steps in the underlying theory behind this approach. This section investigates the third and fourth of these steps, that is, whether knowledge is acquired and if it is put into practice, failure to do so is labeled the knowledge-practice gap.

Evidence from the Save the Children Data

The Save the Children study used three types of nutrition knowledge to calculate the knowledge-

TABLE G.4

Mother's Participation in Group Discussions on Nutritional Topics (percent)

	Malnutrition	Birth weighing	Weight gain	Iodine	Vitamin A	Prevention
Participated	66.4	66.2	62.2	47.5	48.6	52.2
Did not participate	30.8	30.9	34.7	49.1	47.9	44.3
Not invited	2.8	2.9	3.1	3.4	3.5	3.5

Source: BINP midterm data.

TABLE G.5		Eligibility Criteria for Children’s Supplementary Feeding
		Eligibility conditions
Normal children		
From 6 to 12 months	Less than 600 gram gain in 2 months	
From 12 to 24 months	Less than 300 gram gain in 2 months	
I and II degree malnutrition		
From 6 to 12 months	Less than 600 gram gain in 3 months	
From 12 to 24 months	Less than 300 gram gain in 3 months	

TABLE G.6	Malnutrition Rates at Feeding Start by Feeding Episode	
	Observations	Percent
< -4	81	5.4
>= 4 & < -2	903	59.5
>= -2 & < 0	522	39.4
> 0	11	0.7
All children	1,517	100.0

Source: Calculated from Save the Children register data.

TABLE G.7	Eligible and Non-Eligible for Supplementary Feeding Among the Fed Children	
	Observations	Percent
Eligible	1,283	84.6
Non-eligible	233	15.4
All children	1,516	100.0

TABLE G.8	Participation of Pregnant Women in Weighing and Feeding Sessions (percent)	
	Midterm	Endline
Weighing sessions	73.2	71.6
Feeding sessions	49.2	44.8
Of which		
Only weighing	30.9	36.7
Only feeding	6.9	9.3
Weighing and feeding	42.3	35.8

practice gap. These are the knowledge on rest during pregnancy, colostrums, and exclusive breastfeeding reported in table G.10.

Regarding knowledge, all the differences in knowledge between project and control areas are statistically significant, though these differences are not that great. There are differences in practice between project and control areas, but only one of them, rest during pregnancy, is statistically significant. Knowledge is generally high, being between 50 and 80 percent in both project and control areas. The knowledge-practice gap is small in the case of colostrums, moderate in the case of rest, and extremely high in the case of exclusive breastfeeding. The project improves both knowledge and practice, but does not reduce the knowledge-practice gap.

Regression analysis can be used to answer two questions: (1) What are the obstacles that prevent the knowledge being put into practice? and (2) Does the project help to reduce the knowledge-practice gap in general, or for mothers with particular characteristics? The knowledge-practice gap is modeled using a probit regression, where the dependent variable is one if there is a knowledge-practice gap. The sample is composed of mothers who have the knowledge about the three nutritional practices described above. The estimation of this model can produce biased coefficients, because unobserved determinants of the knowledge can have effects on the practice as well. In order to avoid this bias, a two-step Heckman approach is adopted, first estimating a selection equation explaining the knowledge among mothers, and then a gap equation for mothers who have the knowledge. The results are in table G.11.

The explanatory variables for the selection equation (the "knowledge" equation) are mother's age, the education of the mother and of the head of household, the membership of a group,⁷ and residence in a project area. The variables included in the knowledge-practice gap equation are possible constraints that can prevent mothers from putting in practice their nutritional knowledge. These include time constraints due to house and farm work, resource availability, and women's agency within their households. The number and type of obstacles

used in the regression is limited by the information contained in the dataset. In order to assess the project effect on the knowledge-practice gap, a project dummy is included together with some interactive terms to pick up possible differential project impact on different groups.

The multivariate analysis shows that attending nutritional counseling indeed has a significant impact on a woman's nutritional knowledge, though being in receipt of supplementary feeding does not.⁸ However, even when these participation variables are included, the BINP project dummy is still significant. This means either that there are spillover effects (women who get the knowledge in nutrition sessions communicate it to others) or that other project activities not captured in the participation variables—for example, women's group meetings—are also channels for communication of nutrition education. According to these regression results, simply living in the project area raises a woman's probability of having a piece of nutrition knowledge by 7 percent, but full participation in project activities increases this probability to 10 to 23 percent (figure G.2). That is, the proportion of women aware of the importance of colostrums feeding is 23 percent greater for women participating in project activities than for women in the control area. Other determinants of women's nutritional knowledge are found to not vary much between the project and control and so account for only a small amount of the difference in knowledge between the two areas.

The gap equations show that some of the obstacles included help determine the gap. Time at work on the farm and in the house, attending children, and other household members' needs, all reduce mothers' ability to rest during pregnancy. There are fewer obstacles to the mother's use of colostrums for lactation, which is a reasonable result. The number of pregnancies the mother has reduces the gap, perhaps as these women have learned from previous experience. Work on the farm and general lack of resources seem to increase the knowledge-practice gap in the case of breastfeeding, which is also larger for daughters-in-law. The project area dummy is never significant, but the project does seem to close the knowledge-practice gap for breast-

TABLE G.9		Participation in BINP Activities by BMI Status			
		Normal	Mild	Moderate	Severe
Midterm					
Weighing sessions		72.8	74.2	72.6	72.5
Supplementary feeding		39.4	49.5	56.0	58.6
Endline					
Weighing sessions		71.8	70.0	72.4	74.2
Supplementary feeding		34.3	51.4	60.2	68.9
Recipients of supplementary feeding by BMI status					
Midterm		25.2	31.6	22.9	20.3
Endline		39.8	33.2	16.3	10.7

Note: BMI categories used here are those recommended by WHO (1995): greater than 18.5 is normal, between 17 and 18.5 is mild thinness, between 16 and 17 is moderate, and below 16 is severe.

feeding for daughters-in-law and better-off families (who are otherwise less likely to breastfeed).

The BINP Evaluation Sample

The BINP evaluation data included information allowing calculation of the gap for a larger range of behavior. However, this information was collected

TABLE G.10		Nutrition Knowledge of Pregnant and Lactating Mothers (Save the Children data)		
		Project (%)	Control (%)	t stat.
Rest during pregnancy				
Knowledge		77.1	69.4	4.00***
Practice		58.4	52.3	3.56***
Knowledge-practice gap		30.7	34.7	-2.15**
Colostrum				
Knowledge		75.4	71.9	4.33***
Practice		62.7	51.5	1.53
Knowledge-practice gap		18.3	21.1	-1.30
Breastfeeding				
Knowledge		77.6	68.7	5.06***
Practice		3.6	4.9	-1.16
Knowledge-practice gap		95.7	94.2	1.08

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

TABLE G.11

Knowledge and Knowledge-Practice Gap Equations

	Rest during pregnancy		Colostrum		Colostrum	
	Coeff.	t stat.	Coeff.	t stat.	Coeff.	t stat.
Knowledge-practice gap equation						
Daughter-in-law	-0.04	-0.40	-0.07	-0.68	0.30	2.30**
H/h owns land	0.00	1.63*	0.00	-0.07	0.00	0.95
Farming household	0.14	3.44***	-0.07	-1.67*	0.12	1.99**
Durables index	0.00	-0.16	-0.01	-0.77	-0.10	-3.46***
No. of children	0.09	4.58***	-0.06	-2.69***	0.02	0.53
Mother's education	-0.01	-0.30	0.03	1.01	0.04	1.38
Father's education	-0.04	-1.35	0.03	1.00	-0.04	-1.36
Elderly male in household	0.11	1.90*	0.03	0.60	0.03	0.35
Project area	0.13	1.25	0.03	0.26	-0.13	-0.58
Project * durables	-0.01	-0.72	-0.01	-0.69	0.07	2.49**
Project * daughter-in-law	-0.06	-0.57	-0.01	-0.09	-0.28	-1.81*
Project * children	-0.03	-1.28	0.03	1.00	-0.01	-0.22
Intercept	-0.95	-11.07***	-1.08	-10.17***	1.57	5.66***
Knowledge (selection) equation						
Mother's age	0.04	2.27**	-0.01	-0.50	0.03	1.49
Mother's age squared	0.00	-2.18**	0.00	0.01	0.00	-2.13**
Mother primary educ.	0.26	3.59***	0.18	2.51**	0.20	2.29**
Mother secondary educ.	0.44	4.94***	0.43	5.20***	0.54	5.43***
Mother tertiary educ.	0.36	1.97**	0.98	5.80***	0.91	4.61***
Education h/h head	0.02	3.00***	0.02	3.04***	0.01	2.07**
Female headed household	0.33	3.04***	0.22	2.28**	0.41	3.90***
Group member	0.07	1.92*	0.07	2.04**	0.08	1.86*
Project area	0.16	2.16**	0.21	2.44**	0.22	2.94***
Supplementary feeding	-0.07	-0.95	0.01	0.13	-0.02	-0.26
Attending nutrition meeting	0.27	3.35***	0.23	3.24***	0.19	2.08**
Participated in nutrition discussions	0.13	1.68*	0.12	1.45	0.01	0.16
Project * primary education	0.03	0.28	0.00	-0.02	0.10	0.93
Project * secondary education	0.05	0.47	0.11	1.05	0.07	0.58
Project * tertiary education	0.40	1.75*	0.11	0.49	0.04	0.18
Intercept	-0.38	-1.33	0.02	0.07	-0.04	-0.14
Selection term	1.44	5.82***	1.89	3.33***	0.61	2.74***
N		7,216		7,217		7,216

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

mainly in the endline survey of 2003, and only for two behaviors (food intake and rest during pregnancy) is it possible to observe that the change occurred over time (tables G.12 and G.13).

The knowledge and the practices related to pregnancy are higher in project areas, and the differences are highly significant. Differences in both knowledge and practice are smaller for breastfeeding and colostrum. The knowledge-practice gap is always very large, except for colostrums. The gap is significantly smaller, both in size and statistically, in project areas compared to control areas. However, in three cases (breastfeeding, colostrums, and hard work during pregnancy), the gap is as large in project areas as in control areas.


Changes in knowledge and practice over time can be observed only for the food intake and rest during pregnancy. Note that the three surveys did not phrase the questions in the same way, and that the baseline knowledge and practice of rest during pregnancy in control areas is unreasonably high. Hence, the analysis over time is limited to midterm and endline data only. Table G.13 shows that knowledge and practice of rest and food intake during pregnancy are increasing over time in both project and control areas. The knowledge-practice gap is increasing in project areas over time, as the practice does not increase as rapidly as the related knowledge.

The knowledge-practice gap is much smaller for health matters than for nutrition, though there are significant origin differences in knowledge of the treatable diseases such as goiter and night blindness. These differences are much smaller in the case of diarrhea and ARI, probably because the latter diseases are much more frequent, as are prevention campaigns all over the country. The same differences are observed in the practices of prevention/treatment when any of these diseases occur.

Table G.14 reports the regression results of the knowledge-practice gap using the BINP data. The knowledge equation shows that parental education matters, but that there is also a large project effect. Some clear patterns emerge with respect to the knowledge-practice gap. Women in poor households are more likely to have any of the three gaps: resource constraints prevent women

FIGURE G.2

Women Living in Project Areas Are More Likely to Have Nutritional Information, Especially if They Participate in Project Activities



from putting knowledge into practice. However, being a member of an income-generation group and having a vegetable garden make it more likely that a woman will not reduce food intake during pregnancy, though the vegetable garden increases the probability of engaging in hard work. Women who are pregnant in the working season are also more likely to not adopt changed practices for all three cases.⁹ Having more children also reduces the likelihood of resting more, not eating less, and not engaging in hard work. There is a project effect in reducing the likelihood of eating down, which is possibly the result of food supplementation rather than the advice per se, but there is also a project impact in reducing the knowledge-practice gap for hard work.

Leakage of Supplementary Feeding

Supplementary feeding may not have the desired nutritional impact if there is either leakage (it is not consumed by the targeted individual) or substitution (the target individual consumes less of other foodstuffs, so that the supplementary feeding merely takes their place). In principle, BINP avoids leakage by having feeding sessions at which the food is consumed. However, as shown in table G.15, in practice, many

TABLE G.12

Mothers' Knowledge and Practices Related to Major Child Diseases (BINP data)

	Project (%)	Control (%)	t stat.
Diarrhea			
Knowledge	91.2	84.5	2.88***
Practice	92.6	84.0	5.1***
Knowledge-practice gap	1.3	4.8	-5.16***
ARI			
Knowledge	82.0	73.7	3.96***
Practice	87.2	82.4	3.67***
Knowledge-practice gap	3.7	5.9	-1.74*
Goiter			
Knowledge	52.3	32.4	9.25***
Practice	52.8	32.5	9.37***
Knowledge-practice gap	2.4	6.2	-1.89*
Night blindness			
Knowledge	72.1	57.0	3.94***
Practice	73.4	57.7	4.09***
Knowledge-practice gap	2.0	3.3	-0.96

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

TABLE G.13

Nutrition Knowledge of Pregnant and Lactating Mothers
(BINP data)

	Baseline			Midterm			Endline		
	Project (%)	Control (%)	t stat.	Project (%)	Control (%)	t stat.	Project (%)	Control (%)	t stat.
Food intake									
Knowledge	62.8	65.4	-0.48	83.6	51.5	15.4***	91.9	67.5	17.43***
Practice	47.5	74.6	-7.52***	55.7	21.9	17.17***	58.6	29.4	8.75***
Knowledge-practice gap	49.9	20.5	6.92***	34.3	61.3	-11.31***	37.7	61.4	-5.63***
Rest									
Knowledge				81.2	61.4	8.18***	89.0	58.0	23.36***
Practice				64.0	35.2	11.61***	67.0	46.2	6.98***
Knowledge-practice gap				22.8	46.9	-9.54***	27.8	40.0	-3.49***
Hard work									
Knowledge							93.9	82.7	5.35***
Practice							52.6	44.1	2.98***
Knowledge-practice gap							44.8	52.5	-2.62**
Smoking									
Knowledge							88.0	65.4	4.76***
Practice							80.5	85.5	-3.13***
Knowledge-practice gap							17.9	10.4	4.35***
Iodized salt									
Knowledge							94.5	77.4	13.09***
Practice							79.2	42.7	7.53***
Knowledge-practice gap							17.6	39.3	-5.21***
Breastfeeding									
Knowledge							96.6	89.6	5.86***
Practice							62.5	61.1	0.34
Knowledge-practice gap							37.0	39.6	-0.65
Colostrum									
Knowledge							97.4	92.4	4.87***
Practice							95.6	92.3	3.73***
Knowledge-practice gap							2.7	4.1	-1.85*

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

children and pregnant women get the food outside of the feeding sessions, making leakage a stronger possibility. In addition, Save the Children found that less than half of mothers (42 percent) consumed the whole sachet at the CNC, and one-third (32 percent) admitted having shared it with someone else.

Evidence on substitution comes from the question regarding whether mothers ate more during pregnancy or not. It is relatively common practice to “eat down” (that is, eat less) during

pregnancy, mainly as it is believed to make for an easier childbirth. Through its nutritional counseling BINP encourages women to eat more not less, and facilitates doing so through the supplementary feeding program. Table G.16 shows that women receiving supplementary feeding were more likely to eat more than those not receiving it. However, 32 percent of women receiving supplementary feeding ate the same or less during pregnancy, suggesting that they were substituting the food they received for their normal diet.

TABLE G.14 Determinants of Knowledge-Practice Gap During Pregnancy (BINP data)

	Food intake		Rest		Hard work	
	Coeff.	t stat.	Coeff.	t stat.	Coeff.	t stat.
Gap equation						
Mother's age at marriage	-0.01	-0.92	0.03	3.08***	0.03	2.99***
Mother is head of household	0.04	0.21	-0.17	-1.09	-0.39	-2.37**
Mother is widow or divorced	0.55	2.03**	0.27	0.87	0.10	0.38
Children alive	0.08	4.93***	0.14	8.81***	0.14	8.15***
Housewife	-0.02	-0.19	-0.15	-1.15	-0.42	-4.49***
Land owned	0.01	0.30	0.01	0.87	0.01	0.39
Farming	0.08	1.94*	0.06	1.24	0.24	5.15***
Poor household	0.21	2.64**	0.25	2.84***	0.12	2.07**
Member of community-based income-generating group	-0.10	-2.10**	-0.05	-1.17	-0.01	-0.13
Vegetable garden	-0.13	-2.68***	-0.02	-0.42	0.14	2.96***
Pregnant in working season	0.14	2.17**	0.20	3.35***	0.26	2.91***
Project area	-0.55	-2.92***	-0.04	-0.53	-0.14	-1.81*
Poor in project area	0.03	0.37	-0.13	-1.62	-0.21	-2.12**
Pregnant in working season in project area	-0.22	-2.72***	-0.01	-0.15	0.25	3.46***
Constant	0.17	0.68	-1.49	-6.71***	-0.81	-4.56***
Knowledge equation						
Mother's age	0.01	0.82	0.01	2.24**	0.01	0.58
Mother's education	0.05	5.89***	0.05	6.36***	0.05	5.04***
Father's education	0.02	2.54**	0.01	1.25	0.02	2.54**
Project area	0.96	17.90***	1.03	18.04***	0.59	6.43***
Constant	0.08	0.57	-0.24	-2.32**	0.63	4.43***
Observations		4,967		4,967		4,967
F statistic		27.6		30.63		17.9
P-value		0.000		0.000		0.000

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

It is of course possible that the women receiving the supplement did not eat more as they gave the good away (leakage) rather than substituting the food for their ordinary diet. But the Save the Children data suggest this cannot be entirely so: of the 68 percent of women who did not share their BINP food, 63 percent said they ate the same or less, thus providing clear evidence of substitution.

The Nutritional Impact of BINP Interventions

Child Nutrition¹⁰

Three evaluations of BINP have been conducted so far that compare project impact on child nutritional outcomes using large household survey

datasets. These are the endline evaluation of BINP by Karim and others (2003), the evaluation conducted by Save the Children (2003), and IMED (Haider and others 2004). The results of these studies are summarized in table G.17.¹¹

The BINP evaluation conducted by Karim and others (2003) compares project outcomes of six upazillas where the BINP project was implemented, and two upazillas where the project was not implemented (see the technical note, available on request, for a review of the quality of these data). The data were collected in three rounds: at the onset of project implementation, at a midterm point and at the end. There are several problems with these surveys that limit their use for empirical analysis. First, the sample size of the baseline survey is very small, and the qual-

TABLE G.15**Sources of Supplementary Food (percent)**

Source	Children	Pregnant women
CNC (feeding sessions)	75.3	49.6
Pushti Apa (CNP) brings it to the home	16.8	28.2
Somebody from the household brings it from the nutrition center	7.7	22.1
Take/took it preparing by own	n.a.	0.1
Total	100	100
Number of observations	1,710	2,643

ity of the data is not very good.¹² Second, the questionnaires used in the three rounds are different, and similar questions are often non-comparable. Third, no longitudinal data across villages were collected. This set of problems reduces the scope for analyzing changes in outcomes over time. There are also problems that limit the use of the cross-sectional aspect of these data. First, anthropometric measurements of children were taken at different times of the year. Given the high seasonality of child malnutrition in Bangladesh, this is a potential source of bias. Second, while the sample of project thanas is representative of all Bangladeshi divisions, the sample of control thanas is not. If there are substantial differences in nutrition determinants between regions, comparisons of outcomes can be biased. Third, some of the selected control thanas may have benefited from nutrition projects of other NGOs. In one of the thanas, BRAC (the implementing NGO of the first phase BINP thanas) operated nutrition activities in the same period as the BINP was implemented (Karim and others 2003). There is a risk that the control

thanas do not represent a sample of household “without the project.”¹³ Finally, the control samples are particularly small, especially at the baseline.

Save the Children collected data between March and May 2002 in three of the first-phase BINP thanas, and in three control thanas. According to Save the Children, these control thanas do not have serious problems with “contamination” and lack of regional representativeness, as did the BINP control group. However, the control sample is very small, and there is the possibility of seasonal bias, which cannot be assessed since dates of anthropometric measurements are not reported in the dataset. Finally, the IMED study collected data from 12 project upazillas from all 3 phases of BINP, and 12 controls selected from neighboring upazillas. Both studies have to rely on single difference estimates of project impact, and so depend on strong assumptions regarding the quality of the control.

According to the BINP evaluation data, there was no difference between project and control areas for the mean of any of the three z-scores at the baseline. There is a significant project effect on weight for age at the midterm. Height for age, however, remained unchanged, resulting in a very large effect on weight for height. At the endline, the data show a significant project effect on height for age. However, it is not clear how this could happen, since the project did not succeed in improving weight for age for the same children. The endline difference in weight for age is a combination of the difference in weight for age and height for age. Since heights are increasing and weights are stable, weight for height decreases.

The Save the Children data show no significant differences between the project and control. The IMED data show not much difference either, though there is a slight difference in severe stunting, partly offset by higher moderate stunting in project areas.¹⁴ This finding is consistent with the BINP evaluation, which also suggested that some children were moved from severely to moderately stunted.

In summary, these studies find either no project effect, or where there is one, it appears rela-

TABLE G.16**Eating Habits and Supplementary Feeding**

	Eating habits during pregnancy (%)		Number of observations
	Ate more	Ate same or less	
Received supplementary feeding			
Yes	67.6	32.4	318
No: project area	61.3	38.7	762
No: control	24.6	75.4	353

tively small.¹⁵ There are two problems connected to the estimation of project effects presented above: (1) the possible poor quality of the control and (2) the estimated effects are average population effects that conflate two issues, project participation and effect on those who participate. This study improves on these estimates of project effects by using national data to create a comparison group, and by separating participation (analyzed above) from impact on participants.

The comparison group for this study is constructed based on the characteristics of the child and of its household of origin using data from Helen Keller's Nutritional Surveillance Survey (Box G.1), using the method of propensity score matching (see Appendix 3). Project effects are estimated only for the BINP evaluation samples of midterm and endline children. For several reasons a similar analysis cannot be applied to the children sampled by the Save the Children survey.¹⁶

NSP data were used only from the rounds that correspond to the time of collection of the midterm and endline BINP surveys. This restriction is necessary given the high sensitivity of nutritional indicators to seasonal variations in Bangladesh. The rounds used are therefore the October/November and December/January rounds for the midterm evaluation, and the December/January and the February/March rounds for the endline evaluation. These data are complemented with the data from the comparison sites that were collected for the BINP evaluation. There are 7,091 control observations against 3,000 project observations at the midterm, and 6,247 control observations against 2,269 project observations at the endline.¹⁷

To create a match, it is necessary to estimate the propensity to participate using a model that: (1) avoids explanatory variables influenced by the project, (2) includes all possible observable determinants of participation, and in particular those that are influencing both outcomes and participation. It was shown above that various factors influence participation. Model specification is more constrained here, since only variables also available in NSP can be included in the model. Table G.18 lists the variables used in the

TABLE G.17		BINP Project Impact from Other Studies			
	Project	Control	Difference	Number of observations	
				Project	Control
BINP evaluation					
Baseline					
HAZ	-2.36	-2.49	0.13	403	153
WAZ	-2.30	-2.24	-0.06	425	161
WHZ	-0.77	-0.73	-0.04	393	150
Midterm					
HAZ	-1.98	-1.89	-0.08	3,488	1,147
WAZ	-2.04	-2.14	0.09***	3,502	1,149
WHZ	-0.98	-1.19	0.21***	3,520	1,149
Endline					
HAZ	-1.90	-2.07	0.17***	2,554	837
WAZ	-1.87	-1.94	0.07	2,567	842
WHZ	-0.84	-0.76	-0.08**	2,548	837
Save the Children					
HAZ	-1.73	-1.78	0.05	1,640	817
WAZ	-1.87	-1.92	0.05	1,640	817
WHZ	-1.04	-1.06	0.03	1,640	817
IMED (0–59 months)					
WFA: severe	14.0	14.0	0.0	3,024	
WFA: moderate	13.0	12.9	0.1	3,024	
HFA: severe	7.0	8.0	-1.0	3,024	
HFA: moderate	7.7	7.4	0.3	3,024	

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

specification of the selection model, including their definition in the two datasets used (the BINP evaluation and the NSP data). The second column describes how the variables from the two datasets compare. The third column explains the inclusion of the variable in the model when the reason is not obvious. The results from these equations were used to estimate the propensity scores for the purposes of matching.¹⁸

Table G.19 shows the estimated average project effects and the effects of the project on the participants. The effect considered is the difference in the mean z-score of the three nutritional indicators for children aged 6 to 23 months, between project area and non-project areas. Average treatment effects are calculated using all project observations, independently of project participation. It is possible to imagine that even

BOX G.1**The Helen Keller
International Nutritional
Surveillance Project (NSP)**

In 1990, the NSP started collecting data on health and nutrition of mothers and children of Bangladesh. Data are collected bi-monthly in “rounds” corresponding to six Bangladeshi seasons. At each round, the data collection takes place in six or seven weeks, from roughly 10,000 rural households. The household is interviewed only if it contains at least one physically able child under five years of age (aged between six and 59 months prior to 2000), and if the mother is available. Starting from February 1998, the data are nationally representative. Four thanas were randomly selected from each administrative division, which makes a total of 24 thanas. At each survey round, 15 mauzas (10 prior to 2000) are randomly selected with probability proportional to size from each thana. One village is randomly selected from each mauza (administrative unit within a thana), and 25 households are systematically sampled from each village. Thus, while the 24 thanas that were selected in 1998 are always the same, the villages and the households interviewed differ at every round. The data constitute a series of repeated cross-sections. The dataset contains 49 variables, including location variables, household and child characteristics and anthropometric measurements.

children not directly participating in the project activities are nevertheless project beneficiaries, as was indeed shown in the “knowledge equations” reported above. But it was also shown that there was a bigger impact on knowledge from fuller participation. Hence, the average treatment effects are included only for completeness, and to offer a comparison with the project effects on the project participants.¹⁹ The rows “average treatment effect for the treated” show the project effects only for the children who regularly attended the growth monitoring sessions. As expected, table G.19 shows, in general, greater project effects compared with the average treatment effect.

These results differ from the BINP evaluation results (reported in the first column of “average treatment effect”) in two ways. First, there is a much larger project effect at the midterm, but no project effect at the endline. Second, these results are more coherent, since at each point in time they show that weight for age and height

for age give similar results. On the contrary, the BINP evaluation shows a positive difference in weight for age but not in height for age at the midterm, and a positive difference in height for age but not weight for age at the endline, although it is not clear why, given that height for age scores are the cumulated results of weight for age scores. The results also differ from the Save the Children evaluation in that there is a project effect on weight for age at the endline, though this effect is very weak.

These results can be summarized as follows:

- There is evidence of a project effect for all nutritional indicators at the midterm, and for weight-for-age at the endline.
- The size of the effect is small (negligible in the case of the endline data), a reduction of 0.1 *z*-scores is equivalent to a 3–4 percent reduction of malnutrition rates.
- There is evidence that the project performance was worse at the endline with respect to the midterm.

Table G.20 presents the estimated projects effects corresponding to different levels of project participation.²⁰ There is evidence of a project effect only for children who participated regularly in the project.

Tables G.21 and G.22 show project impact according to nutritional status.²¹ Children measuring between -3 and -2 *z*-scores are normally considered malnourished. Children scoring less than -3 are normally considered severely malnourished. The category of the severely malnourished children is split into those who score between -3 and -4 , and those who score less than -4 . A *z*-score of -4 is approximately the cut-off point used to define the third-degree level of malnutrition in the Gomez classification. This category is interesting because children scoring less than -4 have much higher mortality risk than other children, and because the project is targeting children of third degree of Gomez malnutrition for food supplementation. Figure G.3 provides a visual representation of the same distribution of *z*-scores. *Z*-scores in healthy populations have a distribution that is very close to the normal distribution. The percentage of children

TABLE G.18 Definition of the Variables Used in the Model

Variable	Specification	Notes
Child age	Age is in completed months.	
Child sex	Gender of the child is 1 for female.	
Child is breastfed	Whether the child is breastfed at the time of the interview.	Children over 6 months only, and therefore is independent of the project, which recommends exclusive breastfeeding for children under 6 months.
Season of birth of the child	The season in which the child was born obtained using date of birth and date of interview.	This is a weak correlate of low birth weight, which in turn is a determinant of future nutritional status.
Mother's age	Age in completed years.	
Female-headed household	The mother is head of the household.	
Widow or divorced	Whether the marital status of the mother is widow or divorced.	
Mother's height	Mother's height measured in centimeters.	This predicts child's height. Largely missing before 1998.
Mother is pregnant	Whether the mother is pregnant at the time of the interview.	Pregnant mothers are specifically targeted by the project.
Household size	This is the number of people who live together and take food from the same pot (NSP). The definition used by BINP is not known.	
Number of children under five	Number of children under five. Prior 2000 only the number of children aged between 6 and 59 months was recorded.	
Occupation of head of household	NSP: Occupation of the person that provides the majority of household income. BINP: occupation of the head of household. BINP midterm and endline surveys used different categories	We assume that the main earner and head of household are the same
Mother's education	Four levels of education are defined: illiterate, primary (including uncompleted), secondary (including uncompleted) and higher. The variable is originally the number of completed years of schooling in both NSP and BINP.	This is a correlate of wealth.
Father education	Same as for the mother.	
House	This is the area size of the house where household lives. It is measured in square feet based on measurements of width and length.	
Land	This is the land owned by the household in decimals. Land that is rented, sharecropped, and homestead garden are not included.	
Landlessness	Household with no land.	
Without well	Household without a tubewell from which obtaining drinking water (so use pond or river water or rain water).	
Without latrine	Households without an open or closed latrine. These households dispose of excreta in rivers or in the bush.	

TABLE G.19 Project Impact Using Propensity Score Matching

	BINP means	Regression	One-to-one matching	Nearest neighbor matching	Kernel matching	Weighting
Midterm: average treatment effect						
HAZ	-0.09	0.09***	0.08***	0.10***	0.10***	0.08***
WAZ	0.10**	0.13***	0.14***	0.14***	0.14***	0.13***
WHZ	0.21**	0.14***	0.16***	0.13***	0.13***	0.12***
Midterm: average treatment effect on the treated						
HAZ	-0.09	0.11***	0.10***	0.10***	0.11***	0.10***
WAZ	0.10**	0.14***	0.12***	0.13***	0.14***	0.14***
WHZ	0.21**	0.13***	0.10***	0.12***	0.12***	0.13***
Endline: average treatment effect						
HAZ	0.17***	0.01***	0.01	0.01	0.01	0
WAZ	0.07	0.01***	0.01	0.02	0.01	0.01
WHZ	-0.08**	-0.01*	0.01	0.01	0.01	-0.02
Endline: average treatment effect on the treated						
HAZ	0.17***	0.02	0.08**	0.03	0.03	0.01
WAZ	0.07	0.04*	0.09**	0.06*	0.03	0.03
WHZ	-0.08**	0.02	0.03	0.04	0.03	0.01

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

below -2 z -scores is normally 2.3. The vertical lines in the charts help to assess the difference in malnutrition rates between project and non-project areas.

The shapes of the midterm and endline distributions are very similar, although project effects at the endline are much smaller. For all nutritional indicators the project has the effect of shifting the distribution to the right. In particular, it reduces the number of malnourished children between -3 and -2 z -scores, and increases the number of normal children. In addition, the weight for age charts clearly show that the project

reduces the number of severely malnourished children (less than -4 z -scores), which in turn can be found in excess in the category of malnourished between -4 and -3 z -scores, as many of them have been upgraded to this category.

Supplementary Feeding Program

As noted above, it proved difficult to construct a control for the supplementary feeding program since we do not have data on growth faltering among the control, and the determinants for the selection equation seemed not to work well, so the problem of endogeneity of program placement remained, resulting in an apparent negative program impact. The best source for analysis is the Save the Children register data, which includes the weight for age z -scores of all children who had received supplementary feeding, both during the feeding and at other times. The regression results for the change in the WAZ between starting and ending feeding are shown in table G.23.

The main results are that supplementary feeding is much more effective for children who

TABLE G.20 Project Effect by Level of Participation

	Midterm		Endline	
	Irregular	Regular	Irregular	Regular
HAZ	-0.06	0.08***	0.03	0.01
WAZ	0.05	0.13***	0.02	0.03
WHZ	0.07	0.13***	-0.02	0.02

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

are malnourished, and particularly those who are severely malnourished. The results also provide evidence of the adverse effects of substitution. Finally, there are very striking seasonal effects from the weight gain experienced during supplementary feeding.

The lean season in Bangladesh, as determined by WAZ for children under five using the NSP data, runs from April to November, with the start and end months varying from year to year (see Annex J). Children receive supplementary feeding for 3-4 months, so four categories can be identified for the start-end points

- Good-Lean, starting feeding in January to March, and in some years, starting in April or May: the coefficients show these children experience no, or negative, weight gain from the seasonal effect, picking up the effect of reduced food intake in these months
- Lean-Lean, starting feeding in June to August, and in some years early September: the coefficients show a greater weight gain for children starting and finishing feeding in the lean season.
- Lean-Good, starting feeding September to November, and in some years August: as expected there is a positive weight gain for these children, partially reflecting greater food availability at the end of the period, this effect is greatest for children starting November as it takes time for the effect of increased food availability to feed through to improved WAZ.
- Good-Good, starting feeding in December and January, and in some years November: the coefficients for November and December are high reflecting the cumulative effect of three months of better food supply.

The regression results can be used to calculate the weight gain during supplementary feeding for a child with average characteristics. Table G.24, shows the results, varying only the characteristic shown. Children who enter the program more malnourished gain more from the program than better nourished children.²² Children under 12 months appear to show negligible gain in WAZ during the program, compared to close

TABLE G.21		Midterm Malnutrition Rates by Nutrition Status				
Z-scores	HAZ		WAZ		WHZ	
	Project	Control	Project	Control	Project	Control
Z > -3 & < -2	27.6***	31.9	36.8**	39.3	14.8*	16.6
Z > -4 & < -3	13.7	12.6	16.4	15.3	2.3**	1.4
Z < -4	5.2	4.9	1.6***	3.2	0.3***	0.1


Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

to 0.3 WAZ for older children. However, as shown in figure G.4, the NSP data show that the WAZ falls for children across Bangladesh at a rate of about 0.5 WAZ over a three-month period. This occurs since the curves in the growth monitoring chart are very steep for children aged below 12 months (figure G.5), so that substantial weight gain is needed to not fall further below the curve. Hence, compared to this reference group, a zero gain of WAZ is actually a gain of 0.5 WAZ, which is not insubstantial.

Finally, table G.25 reports regressions using the BINP endline data for WAZ and midterm and HAZ at endline, these measures being selected since they are the ones for which a significant impact appears in the double difference and PSM analyses. A project dummy is used, interacted with various characteristics. Program activities cannot be used as they would suffer from selection bias (they are indeed found to have a negative effect when introduced into the regression). The project dummies are significant, though not large. The project mutes the advantage of more educated women, that is, nutritional counseling is substituting for education, and to a lesser extent offsetting the advantages

TABLE G.22		Endline Malnutrition Rates by Nutrition Status				
Z-scores	HAZ		WAZ		WHZ	
	Project	Control	Project	Control	Project	Control
Z > -3 & < -2	29.6	30.2	33.9***	35.1	14.7	13.1
Z > -4 & < -3	11.0	11.5	12.4	12.3	1.9	1.6
Z < -4	3.1	3.0	1.4	1.7	0.0	0.0

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

FIGURE G.3 Distribution of Z-Scores in Project and Non-Project Areas

of higher economic status. It also reduced the “daughter-in-law” disadvantage.

Low Birth Weight

At 40 percent, low birth weight incidence in Bangladesh was one of the highest in the world in the 1990s (Mason and others 2001). The social and economic consequences of low birth weight are well documented (see, for example, Alderman and Behrman 2003). Hence, among BINP’s objectives is “reduction of low birth weight occurrence by half of the baseline level,” which is to be achieved through the “improvement in maternal weight gain by at least 50 percent in at least 50 percent of the pregnant women”²³ (World Bank 1995). These goals are to be achieved through nutrition education and food supplementation. All pregnant women of project villages are regularly weighed and counseled on the proper diet to follow in order to gain the necessary weight. Women identified as malnourished are given a daily food supplement from the first screening to the sixth month of lactation after birth.

Concepts and Definitions

Low birth weight—that is, below 2,500 grams—is determined by two factors: gestational age at birth and intrauterine growth (WHO 1995). In the first case, the infant has a low birth weight because he or she was born prematurely, which does not imply a failure to grow during pregnancy. In the second case, the infant has a low birth weight because its intrauterine growth was impaired by the health and nutrition history of the mother (Intrauterine Growth Retardation, IUGR). The two types of low birth weight should be considered separately because they have different determinants and different consequences for child health. Premature birth is not affected by malnutrition, and infants born prematurely have better chances of surviving and catching up in infancy and childhood than those children with low birth weight resulting from IUGR. In order to isolate birth weight from premature birth effects, infant weight can be measured against the weight for gestational age of a reference population.

Gestational weight gain has four components: fat stores, breast and uterine growth, volume of plasma, and fetus (Kramer 1987). Mother's fat stores are the only component of weight gain that has an influence on the growth of the fetus. But mothers have different fat stores levels before pregnancy, which has two important implications: (a) heavier mothers need lower weight gains during pregnancy, and (b) the effect of additional food intake has a larger effect on infants of mothers that are thinner before pregnancy. The Institute of Medicine (1990) has produced charts of optimal gestational weight gain for women of different BMI, based on a sample of U.S. women who delivered infants of normal weight. These charts recommend a weekly gain of 0.4 kilograms during the second and third trimester for woman of normal pre-pregnant BMI, and of 0.5 kilograms for women who are underweight.

It is common in Bangladesh for women to reduce food intake during pregnancy, a practice often referred to as "eating down." This practice is well documented for India and is explained by the mothers' intention to reduce infant body size, thus facilitating delivery.²⁴ CARE (Vemury 1981) conducted a survey study on food habits in six developing countries and found that some 60 percent of the 700 Bangladeshi women interviewed believed that pregnant women should eat less during pregnancy. The reasons given were the prevention of difficult delivery and customary beliefs (see also Hossain and Choudury 1987). The project document and Bangladeshi women thus share the belief that infant body size can be changed during pregnancy by eating more or less than usual. They do not have the same view, however, on the delivery outcome. Larger infants, according to the project, do not create an additional risk at delivery. However, the review of knowledge on nutrition during pregnancy done by the Institute of Medicine (1990) documents that in Western countries before World War II, women were commonly advised during pregnancy to avoid excessive weight gain or to restrict their food intake, concluding that "in a period when maternal mortality was extremely high and cesarean deliveries were a desperate alternative, limitation of fetal

TABLE G.23		Determinants of Weight Gain During Supplementary Feeding	
Variable	Coeff.	t stat.	
Female child	0.07	1.73*	
Month at which supplementary feeding started			
February	0.11	1.12	
March	-0.10	-1.11	
April	-0.06	-0.60	
May	-0.01	-0.18	
June	0.20	2.12**	
July	0.17	2.16**	
August	0.09	1.12	
September	0.10	1.04	
October	0.05	0.69	
November	0.17	1.73*	
December	0.19	2.31**	
Age at which supplementary feeding started			
From 12 to 17 months	0.25	7.06***	
From 18 to 23 months	0.28	4.76***	
Nutritional status at start of supplementary feeding			
Normal child (above -2 z-score)	0.24	1.05	
Between -2 and -4 z-scores	0.57	2.51**	
Below -4 z-score	0.91	3.70***	
Child shared the food with other household members	0.00	-0.11	
Child was not given other food than supplementary food at meal time	-0.07	-1.91*	
Child was ill during supplementary feeding	-0.02	-0.46	
Child missed some of the sessions	0.00	-0.74	
Constant	-0.52	-2.27**	
R-squared		0.21	
Observations		716	

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

size by restricting maternal food intake was an understandable goal." Maternal mortality at delivery is very high in Bangladesh (estimated in the range of 320 to 400 per 100,000 live births for the period 1998 to 2001²⁵). And nearly 80 percent of these deaths are due to obstetric causes,

TABLE G.24		Weight Gain from Supplementary Feeding for Different Categories of Children	
Malnourished	Severely malnourished	Very severely malnourished	
-0.11	0.22	0.56	
< 12 months	12-17 months	17-23 months	
0.02	0.27	0.29	

FIGURE G.4 **BNP Growth Monitoring Chart**





FIGURE G.5 Mothers' Weight Gain by Month of Gestation

like obstructed labor, abortion, and eclampsia (Faveau and others 1988). Sixty percent of Bangladeshi women reported a complication during pregnancy, delivery, or after delivery (NIPRT 2002). Yet, very few women in Bangladesh deliver in hospitals or are assisted by professionals. The belief of “eating down” is thus understandable, and its effects on delivery are in fact not fully understood. However, some studies (quoted in Pojda and Kelley 2000, p. 24) report that increases in head circumference due to food supplements does not increase mortality risk from obstructed labor.²⁶

Project Impact

This section uses the data collected for the BINP evaluation at midterm and endline. The baseline survey did not collect information on birth weight and pregnancy weight gains. The midterm and endline surveys are cross-sections using different samples for children, newborns, and mothers anthropometric measurements. Two problems arise when comparing the nutritional outcomes of project and control areas. First, not all women interviewed in BINP areas participated in project activities. Second, project and control areas may differ in some important determinants of nutritional outcomes independently of project implementation. The second problem is handled through the difference and double difference approaches (see appendix 1).

Table G.26 shows the difference, and the difference in difference, in the percentage of low birth weight and in the average newborn weight between project and control areas. Data on birth weight were not available at the baseline and therefore the table only contains the differences between the midterm and the endline. Low birth weight is defined as a birth weight below 2.5 kilograms. The ratio of birth weight over the median weight of a reference population of the same sex and gestational month is used as mean weight score.²⁷

There was a reduction in low birth weight incidence in project areas from 26 percent (midterm) to 16 percent (endline). If a similar reduction had been observed between the baseline and the midterm, the stated project objective of reducing the proportion of low birth weight by 50 percent would have been more than achieved. This result seems impressive, even just considering the change between the midterm and the endline. However, only a small part of this decline can be attributed to project activities: there was an even larger decline in the incidence of low birth weight in the control areas. Table G.26 shows a difference of 4 and 2 percentage points between project and control areas at the midterm and the endline respectively, indicating a positive effect of the project assuming the control is good. But the difference in difference over time and areas is pos-

itive (though not statistically significant) since the rate of reduction in low birth weight has been higher in control areas than in project areas.

The bivariate analysis shows no difference in weight gain. But this analysis does not allow for possible differences in determining characteristics between the project and control, for which multivariate analysis is needed. The explanatory variables used are those identified in the literature as main determinants of birth weight (e.g., Kramer 1987). In addition, seasonality is added as a potential determinant of birth weight as are project inputs. Not all relevant explanatory variables are available in the BINP data, and the midterm and endline survey used different questionnaires. Thus, the list of regressors is incomplete and different across surveys. The results are shown in Table G.27.

Gestational age at birth and the sex of the child have the expected values and are highly significant. Using the mean birth score over the reference population values gives very similar results. Newborns from primiparous women have lower birth weight, which is a common result in the literature, though a clear explanation for this finding does not exist. Mother's age is normally a determinant of birth weight. Pregnancy outcomes are usually less favorable for very young and very old women (Kramer 1987). The coefficients confirm this finding, though are only significant in the pooled regression; in which case the turning point is 32 years.²⁸ Illness of the mother during pregnancy (available for endline only) has the expected negative sign, but the coefficient is not statistically significant. A woman who experienced miscarriage, abortion, or stillbirth is more likely to have a child of lower weight. Among the socioeconomic variables used, only father's education and income have an effect on birth weight. Birth weight increases with household income and education of the father. Land ownership also has a positive and significant effect, but the value of the coefficient is negligible. The R-squareds are low in particular for the midterm model, which contains only a small number of regressors. Important explanatory variables missing from

the model include mother's height and pre-pregnancy weight.

The month of measurement is included in the model to detect seasonal effects on birth weight. The effect is strong and the pattern clear. Birth weight is higher in the included months, December-March. However, there is a large difference in the effect of the seasonal variable between the two surveys, which indicates that the model should be better specified. The implication is that mother's nutrition is critical in the last months of pregnancy. However, the seasonal effect is much less relevant in project areas (shown by interacting the seasonal variables with the project dummy, results not shown here). The project tends to smooth out the seasonal pattern of mother's nutrition. The implication is that if the data had been collected during the "hungry" season they would have probably shown a positive impact of the project on birth weights.

The model includes a variable taking the value of one if the mother says she ate more than usual during pregnancy, which is one channel for project impact if women change this practice in response to being in the project area. Women may have eaten more during pregnancy in the project areas both as a result of the availability of food supplementation and in response to the nutritional education against eating down. Whether the woman received food supplementation was also included in the model, but was not significant. This result presumably reflects attrition from leakage and substitution (see tables G.15 and G.16)—it is only if the woman actually eats more that matters, which some women receiving food supplements do, but a sizeable number do not. "Eating more" increases birth weight by 44 grams in the midterm survey and by 88 grams in the endline. The effect is highly significant in both cases. The effect of "eating down" (mothers stating they ate less than usual during pregnancy) is also significant and negative, reducing birth weight by 45 grams (only available for endline). The effect of a mother changing from eating down to eating more is thus 133 grams. An increase of just over 80 grams from eating more is not very large, but is similar to the effect on birth weight observed

in other projects where pregnant women were given food supplements.²⁹ In addition, the effect is larger for women of poorer nutritional status. The interaction of the “eating more” dummy with the income level of the household of residence (results not shown) produces a much larger coefficient (+ 270 grams) for the “destitute” women.

BINP may have affected birth weight through other channels, such as the quality of diet. These channels are captured by the project dummy in the midterm and endline equations.³⁰ However, this dummy is negative in both cases, though insignificant at endline. But interpreting this negative coefficient as an adverse project impact means assuming that all unobserved determinants of birth weight were the same in the project and control areas prior to the project. Pooling the data, and so estimating the double difference effect, allows us to control for these unobservables. In the pooled equation, the coefficients on the BINP area dummy are now unexplained area differences, with the project effect shown by the difference in difference coefficient (see Appendix 1). This coefficient is negative but relatively small and very insignificant. There is a large and significant time effect on birth weight in both areas, where birth weight has increased on average by 100 grams, indicating a large autonomous change, most likely as a result of improving mother's nutritional status.

The total project effect is calculated by combining both the effect through mother's eating patterns and the project effect coefficient (the BINP dummy in the midterm and endline equations and the difference in difference coefficient for the pooled data). The results from the pooled regression suggest an increase in birth weight of just over 80 grams through eating more (on the assumption that a woman does eat more, which not all do), with no other project effects. The figure of 80 grams is thus an upper estimate of the average effect. This is an average effect, since it was also shown that the impact is greater during the lean season and higher for poorer (and therefore probably less well nourished) women. The average impact on low birth

TABLE G.25	Determinants of the WAZ and HAZ with Interactive Terms			
	Midterm		Endline	
	Coeff.	t stat.	Coeff.	t stat.
Father's education	0.01**	1.98	0.02***	4.05
Land	0.00	1.40	0.00	0.01
Female head of household	-0.12	-1.18	0.29*	1.92
Water	0.13***	4.82	0.05	1.46
Sanitation	0.02	0.68	0.08**	1.94
Female child	0.02	0.62	0.08**	2.06
Child age	-0.06***	-21.00	-0.06***	-15.14
Household size	-0.02***	-3.14	0.00	0.41
Birth order			-0.05***	-3.63
Birth interval			0.00***	3.85
Mother's height	0.04***	14.75	0.05***	14.39
Project area	0.15**	2.50	0.18	1.62
Primary education	0.18**	1.96	-0.07	-0.64
Secondary education	0.20***	2.75	0.10	1.12
Higher education	0.01	0.03	0.30*	1.85
Primary education*project area	-0.15	-1.48	0.10	0.80
Secondary education*project area	-0.06	-0.82	0.02	0.15
Higher education*project area	0.39**	2.15	0.04	0.24
2 nd wealth quantile	0.17**	2.36	0.18	1.61
3 rd wealth quantile	0.26***	3.56	0.36***	3.21
4 th wealth quantile	0.25**	2.32	0.42***	3.53
2 nd wealth quantile*project area	-0.10	-1.15	-0.05	-0.41
3 rd wealth quantile*project area	-0.19**	-2.34	-0.18	-1.43
4 th wealth quantile*project area	0.00*	0.00	-0.17	-1.25
Early marriage	-0.15**	-2.03		
Early marriage*project area	0.11	1.28		
Daughter in law	-0.01	-0.13		
Daughter in law*project area	0.12	1.00		
Age difference			-0.13	-1.52
Age difference*project area			0.10	1.02
Seasonal dummies (not shown)				
Constant	-7.79***	-18.45	-9.28***	-17.39
R-square		0.19		0.19
Observations		4,753		3,369

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

weight is not large at all. What explains this relatively weak performance?

Project Impact on Pregnancy Weight Gain

The project aimed to increase birth weight by increasing mothers' weight gains during pregnancy. The first hypothesis to explain the absence of a reduction in the incidence of low birth weight in project areas compared to con-

trol areas, is that mothers' pregnancy weight did not increase as expected.

Figure G.5 plots weight gains of project and control areas for month of gestation compared to a reference population. The reference population is the one used by the Institute of Medicine (1990) for underweight women, defined as women of BMI below 19.8, which includes most the women in the sample. The lines for the project and control areas are obtained cumulating median weight gains at each month of gestation. Few observations are available for weight gains before the third month of gestation, therefore the curves before that date are not reliable. Both graphs show a growth pattern inferior to the reference population, and no difference between control and project at the midterm, though there is some difference at the endline. Similarly, table G.28 shows the differences between areas and over time in mean weight gains. Weights were measured at one-month intervals for samples of pregnant women at different stages of gestation. The table shows a significant project effect and a positive project effect over time compared to control areas. However, as in the case of birth weight, these comparisons are of little use unless differences in the determinants of weight gain are allowed for with multivariate analysis.

Table G.29 presents the results of regressions of weight gains of pregnant women measured in grams. The same model is used for the midterm, the endline, and the pooled data. Weight gain is inversely correlated to women's age, though the link is not clear (Institute of Medicine 1990). Weight gains are higher in the second trimester of gestation and then decrease in the last trimester. The variables indicating the month of measurement depict the shape of this function. The data show again the presence of seasonality in the consumption of food. February is the month in which weight gains are larger. Weight gains increase with income, and landless households are disadvantaged. Weight gains are larger in project areas, but the rate of change over time is the same as in control areas. There is a time increase of some 300 grams in monthly gain, which is common to project and control areas. A variable representing the mother following the

advice of eating more during pregnancy has the expected positive sign of increasing weight gain, and this appears to be the main channel through which the project has an effect. While the single difference estimate of the pure project effect is positive, it becomes insignificant in the double difference estimate (in fact negative and of sufficient magnitude to offset the eating effect).

Even at the most optimistic estimate, the overall project effect is to increase weight gains by not much more than 200 grams per month in comparison to control areas (adding the coefficient on the project dummy from the single difference equation to the eating more effect), equaling a total weight gain of just over 1 kilogram over a period of six months.³¹ Kramer's (1987) meta-analysis of the determinants of low birth weight suggests the effect of gestational weight gain on birth weight is 20 grams per one kilogram of total weight gain. Hence, these results suggest that birth weight will increase by just 20 grams as a result of greater pregnancy weight gain. Weight gains of pregnant women in BINP areas would have to much higher in order to exert a significant effect on birth weights.

There is, however, an increase over time in gestational weight gain (over 300 per month) in both project and control areas, which equates to an increase in birth weight of around 60 grams. As mentioned above, this increase is largely a result of improved maternal nutrition. For example, mother's weight, which in Kramer's review increases birth weight by 9.5 grams per kilogram, has increased by an average of 3.4 kilograms between the two surveys. Mother's height, which in Kramer's review increases birth weight by 8 grams/centimeter, has increased by two centimeters between the surveys. Hence, the difference over time in birth weights depends in large part on non-project effects.

It remains to be explained why gestational weight gain has not increased much more in project areas with respect to control areas. Two reasons are possible: (i) low participation rates, particularly by women most in need and, (ii) though women participate and receive the services, there are constraints that prevent the realization of the expected outcomes.

Cost Effectiveness

Table G.30 shows BINP project costs from 1996 through 2003 by main component from the implementation completion report (ICR). The Community-based Nutrition Component accounts for more than 50 percent of total project costs.

Table G.31 shows the change in child malnutrition produced by the project as measured by PSM at the midterm and the endline, and the consequent percentage reductions of malnourished children.³²

Cost effectiveness estimates are calculated in two ways: using total project costs and excluding all costs that are not directly related to children (table G.32 and G.33; Appendix 4 explains the basis of these calculations). The figures show the cost per child to be spent in order to obtain a reduction of malnutrition by 1 percent in the project areas of the first phase. The children considered are between 6 and 23 months of age. Note that costs have almost doubled between the midterm and the endline, because the project has become much less effective in reducing malnutrition rates. There is a reduction of costs by around 50 percent when funds spent for improving mothers' nutritional status are excluded.

In order to estimate the cost of saving the life of a child through the BINP nutritional intervention, we use a method developed by Pelletier and Frongillo (2003, p. 118). This method estimates the reduction in mortality rates following a reduction in the prevalence underweight in the population. A simple formula relates the number of deaths averted to pre- and post-intervention mortality rates. The parameters used by the formula are obtained from regressions of mortality rates on underweight prevalence rates using a panel dataset of 59 developing countries. The results are presented in table G.34 for 1998 and 2003 using overall project costs of each specific year. As above, there are two estimates, one for all project costs, and another for only costs spent on children.

Cost Comparison with Alternative Interventions

To benchmark the figures reported above for BINP, two alternative scenarios are presented:

TABLE G.26		Low Birth Weight, Birth Weights, and Mean Newborn Weight: BINP and Control Areas	
	Low birth weight (%)	Birth weight (grams)	Mean weight (%)
Differences			
Midterm difference	-4.3*	-34.3	0.2
Endline difference	-1.7	-30.9	0.4
Time effect	-12.4***	89.1***	2.3***
Difference in difference	2.6	3.5	0.1
Mean levels			
Midterm project	25.6	2677	75.8
Midterm control	29.9	2712	75.6
Endline project	15.9	2770	78.2
Endline control	17.6	2800	77.8

* One percentage point corresponds to approximately 30 grams.

TABLE G.27		Determinants of Birth Weight (gm) (OLS)				
	Midterm		Endline		Pooled data	
	Coeff.	t stat.	Coeff.	t stat.	Coeff.	t stat.
9th gestational month	104.4	3.01***	167.9	3.66***	133.1	4.77***
10th gestational month	93.5	2.29**	239.9	4.47***	154.8	4.72***
Female child	-58.0	-3.18***	-69.2	-3.60***	-61.2	-4.52***
Parity	-34.2	-1.32	-108.5	-3.63***	-52.1	-2.66***
Age	13.4	1.19	14.3	0.91	20.5	2.11**
Age square	-0.2	-0.89	-0.2	-0.83	-0.3	-1.88*
Mother ill			19.8	0.63		
Pregnancy history			-188.6	-2.26**		
Father's education			14.0	4.33***		
Mother's education			-1.2	-0.33		
Land owned	0.2	2.16**			0.1	3.14***
Middle income			56.8	2.52**		
Rich			115.7	2.79***		
December	360.6	4.38***			112.0	2.75***
January	355.7	4.42***	81.2	0.63	111.0	2.84***
February	422.5	5.25***	78.5	2.08**	161.3	4.71***
March	283.5	2.94***	67.8	1.91*	128.2	3.86***
BINP area	-70.7	-2.72***	-36.6	-1.32	-52.2	-2.14**
More food during pregnancy	43.2	2.21**	88.1	3.75***	86.4	6.11***
Less food during pregnancy			-44.5	-1.74*		
Time Difference in difference					105.8	3.46***
Constant	2,118.6	10.86***	2,387.9	10.51***	-12.8	-0.40
R-square		0.05		0.12	2,220	15.02***
Observations		1,560		1,362		0.07
						2,922

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

TABLE G.28**Difference in Monthly Pregnancy Weight Gain Between Project and Control Areas**

	Grams
Midterm difference	0.5
Endline difference	211.0***
Time effect	-137.9
Difference in difference	210.4**

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

TABLE G.29**Determinants of Pregnancy Weight Gain (OLS)**

	Midterm		Endline		Pooled data	
	Coeff.	t stat.	Coeff.	t stat.	Coeff.	t stat.
Mother's height			4.5	1.99**		
Mother's age	-4.5	-1.34	-6.6	-2.12**	-4.8	-1.70*
4th gestational month	110.7	1.38	433.9	7.90***	129.5	2.01**
5th month	548.8	7.15***	243.7	4.50***	455.4	7.38***
6th month	709.1	9.35***	332.6	6.08***	607.6	9.94***
7th month	992.9	13.09***	293.6	5.33***	849.0	13.88***
8th month	1027.0	12.79***	477.4	8.07***	899.6	13.83***
9th month	1179.6	10.80***	713.2	4.47***	1081.5	11.37***
3rd and 4th week of January	226.0	2.59***			219.6	2.67**
1st and 2nd of* February	454.8	5.22***			463.8	5.51**
3rd and 4th of February	305.5	6.01***			315.0	6.37***
1st and 2nd of March	103.1	2.18**			107.2	2.32**
3rd and 4th of March			-85.7	-1.27	-26.4	-0.30
1st and 2nd of April			-55.3	-0.87	30.8	0.28
Middle income	75.6	1.89*	85.3	2.14**	68.2	2.00**
High income	127.8	1.95*	73.0	1.01	115.9	2.04**
Landless	-97.7	-2.63***	-1.6	-0.04	-81.0	-2.53**
BINP area	162.0	2.84***	136.8	3.00***	156.8	2.92**
Mother ate more during pregnancy in BINP area	73.8	1.77*	67.4	1.70*	76.1	2.12**
Time					332.0	2.84**
Difference in difference					-74.5	-0.72
Constant	298.1	2.37**	273.2	0.78	392.4	3.59***
R-square		0.06		0.10		0.06
Observations		7,562		1,433		8,995

Notes: *, **, *** significant at greater than 10%, 5% and 1% respectively.

simply giving the project budget away as a cash transfer, and giving away an equivalent value of rice. The rice distribution is assumed to be to all households of project areas with a child aged 6 to 23 months, funded with the monthly cost of BINP project in 1998. The cost per household of US\$1.80 per month corresponds to 5.6 kilograms of rice (or 19,720 calories), after allowing for 25 percent administrative costs for distribution.³³ The estimate of the percentage of the administrative cost over total project cost is derived from a review of food supplementation projects (Beaton 1982).

Food supplementation programs often do not achieve the desired results for two reasons. First, the food distributed can be shared among household members, or can substitute rather than complement current feeding. Second, not all calories ingested are transformed into physical growth. Part of the energy obtained through the supplement is dispersed in additional physical activity, and in the maintenance of the increased body size (basal metabolic rate). To account for the first problem we assume that half of the food ingested is substituting rather than complementing the child's normal dietary intake. Only 50 percent of the rice delivered is therefore actually eaten by the child in excess of his or her previous food intake. To account for the second problem, we make the additional assumption that 45 Kcal/day are required for an extra weight gain of 1 gram (see Beaton and Ghassemi 1982), rather than the normally used value of 5 Kcal/day (see Waterlow 1992).

The results are shown in table G.35. Two BINP costs are shown, one is obtained considering all project costs, while the other uses only costs directly attributable to children.³⁴ The costs from the income transfer are also shown for comparative purposes. These costs are far higher, since only a percentage of the income will be used for food (the elasticity of food expenditure is about 0.3), and not all of the incremental food expenditure will benefit children. Of course there are other benefits to the income transfer (and households save resources if they substitute for the rice ration).

The cost of reducing malnutrition with the rice ration is US\$80, compared with US\$200–400

for BINP, and the cost per life saved around US\$1,800 compared with US\$2,300–5,000 for BINP. Hence, BINP does not compare favorably with simply spending project resources to buy rice for children in the project area. The argument against such an approach is that it implies a permanent subsidy, whereas behavior change should become self-sustaining once affected. However, the evidence shows limited success in behavior change, partly as resource constraints prevent recommended changes from being put into practice. Moreover, there may be lower cost models of BCC given the pervasive presence of NGOs, women's groups, and health and family planning officials,³⁵ plus media saturation, which has proved successful in conveying other messages.

Conclusions

Child Nutritional Outcomes

Overall project impact on child nutritional outcomes appears disappointing despite high levels of participation in project activities. Although participation was high it was lower in more traditional thanas, which was partly a result of the project not having a broader target audience encompassing all key decisionmakers. Lack of decision-making power of participants was also a factor behind the knowledge-practice gap, though lack of resources and time played a larger part. There were other weak links in the chain, notably poor targeting, so that some eligible children did not receive feeding, and mothers and some pregnant women did not receive nutritional counseling.

Low Birth Weight

Low birth weight is a correlate of a child's subsequent nutritional and health status, and so of mortality. BINP addresses low birth weight through increasing pregnancy weight gain by discouraging eating down and provided food supplements to malnourished women. BINP aimed to reduce "low birth weight occurrence by half of the baseline level" through the "improvement in maternal weight gain by at least 50 percent in at least 50 percent of the pregnant women." The BINP evaluation found that the pregnancy weight gain target was achieved,

TABLE G.30			Project Costs in US\$ Million (actuals)	
			US\$ million	% shares
National Nutrition Component (NNC)			18.4	31.7
Community-based nutrition component (CBNC)			32.3	55.6
Inter-sectoral nutrition program development (INC)			6.5	11.2
Total baseline cost			57.3	
Physical contingencies			3.4	5.9
Price contingencies			5.0	8.7
Total project cost			65.7	100.0

TABLE G.31		Reduction in Malnutrition at the Midterm and Endline Calculated Using PSM		
	Midterm		Endline	
	Z-score	% change	Z-score	% change
Weight-for-age	+0.13	–5.3	+0.05	–2.0
Height-for-age	+0.10	–4.1	+0.03	–1.3

though that of reducing low birth weight was not. However, the BINP evaluation dataset shows that there was an even larger increase in pregnancy weight gain in the control areas than in the project areas, suggesting that outside factors accounted for these improvements. Greater food availability has improved mothers' pre-pregnancy nutritional status, as shown by rising

TABLE G.32			Cost per Child of Reducing Malnutrition Rates by 1% in First Phase BINP Areas (US\$/% reduction in those below –2z)	
			Midterm (1998)	Endline (2003)
All project costs				
Weight-for-age			4	7
Height-for-age			5	10
Only child costs				
Weight-for-age			2	3
Height-for-age			2	5

TABLE G.33**Cost of Upgrading One Child from Malnutrition in First Phase BINP Areas (US\$)**

	Midterm (1998)	Endline (2003)
All project costs		
Weight-for-age	396	706
Height-for-age	509	1037
Only child costs		
Weight-for-age	183	327
Height-for-age	236	480

BMI in both project and control areas, and this has been the main driving force behind greater pregnancy weight gain.

There has been a project effect, but it has been relatively small. Multivariate analysis shows a small project effect on both pregnancy weight gain and low birth weight, mainly through discouraging the practice of eating less during pregnancy (eating down), assisted by providing feeding supplementation to pregnant women with less than normal body mass index. These effects are greater for more disadvantaged (and so presumably less well-nourished) women and during the lean season.

Potential beneficial impacts of the project are muted for reasons of poor targeting, failure to amend eating practices even when the message is gotten across (the knowledge-practice gap) and the leakage and substitution of supplementary feeding. It is shown that resource constrained women are more likely to have a knowledge-practice gap, supporting the argu-

ment that lack of income is sometimes the binding constraint on nutritional practices rather than lack of knowledge.

How Well Did the Causal Chain Operate?

BINP succeeded in achieving high levels of participation in its activities in project areas, enrolled large numbers in supplementary feeding, and brought about significant increases in nutritional knowledge and associated changes in practice. However, nutritional outcomes in terms of low birth weight have been disappointing. While child nutrition, especially for children participating in supplementary feeding, appears to have been better, the overall difference in performance compared to the control is not great.

The longer the causal chain, the more likely it is that final outcomes will not be realized on account of missing or weak links in the chain. There were two missing links in the BINP chain: the first was the relative neglect of some key decisionmakers regarding nutritional choices (men and mothers-in-law), and the second the focus on pregnancy weight gain rather than pre-pregnancy nutritional status. Participation levels of the target audience were high, but many women escaped exposure to nutritional messages, and there was a high Type I error in the feeding programs (see table G.36). A knowledge practice gap persisted, so many women did not put the advice they received into practice, especially if they were resource or time constrained. Those receiving supplementary feeding often shared it with others or substituted it for their regular foodstuffs. This list of weak links in the chain explain why project impact was muted by the time final outcomes are considered. While attention can be paid to each of these weak links, the BINP experience does demonstrate the difficulty of implementing complex designs.

Should BINP Be Expanded Nationally?

Despite recent progress, Bangladesh has continuing high levels of malnutrition. Several factors lie behind improving nutrition, notably improved agricultural performance. Nonetheless, there appears to be a case for direct nutritional interventions. But in its current form, BINP has had a rather meager impact on nutritional out-

TABLE G.34**Number of Deaths Averted by BINP Project and Cost per Life Saved (US\$)**

	1998	2003
As a percentage of all (estimated) deaths	26.1	10.5
Number of lives saved	296	129
Cost of one life saved I	4,925	8,661
Cost of one life saved II	2,328	4,095

TABLE G.35**Cost of Reducing Malnutrition and Saving Lives from Hypothetical Income Transfer and Rice Ration**

	BINP		Simulations	
	All costs	Only children costs	Income transfer	Rice ration
Extra weight gain (grams/month)	18	18	13	110
Reduction in prevalence of underweight	7.5	7.5	5.4	37.1
Percentage of deaths averted (number of deaths in parenthesis)	26.1 (296)	26.1 (296)	22.7 (258)	70.0 (794)
Cost of removing a child from underweight (US\$)	396	183	555	80
Cost per life saved (US\$)	4,925.0	2,328.0	5,654.0	1,835.0

comes, especially in comparison to its cost. There are two ways to respond to these findings. The first would be to attempt to improve project effectiveness by strengthening the causal chain. The most plausible way of doing this would be

to target resources more finely, toward less well nourished mothers and children, with a larger supplement in the lean season. The alternative is a less sophisticated form of nutritional program, such as a rice ration. Although such schemes are

TABLE G.36**Links in the Causal Chain**

Assumption	Children	Mothers
Attend growth monitoring sessions	Over 90 percent of children attend growth monitoring sessions	Over 70 percent participate in monitoring pregnancy weight gain
Targeting criteria correctly applied; participants stay in program to receive food	Nearly two-thirds of eligible children not fed (reasons: don't attend growth monitoring in first place, wrong application of targeting criteria, drop out of feeding)	60 percent of eligible women not receiving supplementary feeding
Acquire knowledge and put it into practice	One-third of mothers of children receiving supplementary feeding do not receive nutritional counseling. There is a knowledge practice gap (see mothers).	There is a knowledge-practice gap, driven by material resource or time constraints.
No leakage or substitution	One-quarter of children fed at home, increasing possibility of both leakage and substitution.	One-third admit sharing food, and there is substitution for those who do not. At most, 40 percent of eligible women receive full supplementation.
Feeding and nutritional advice have an impact on nutritional status	Supplementary feeding has a positive impact on child nutritional status, especially for the most malnourished children. There is only weak evidence of any impact from nutritional counseling.	Pregnancy weight gain is too little to have a notable impact on low birth weight, except for most malnourished mothers. Moreover, mother's pre-pregnancy nutritional status is more important than pregnancy weight gain. Consequently, birth weight gains are slight, though greater for children of severely malnourished mothers. Eating more during pregnancy is the main channel for both pregnancy weight gain and higher birth weight.


FIGURE G.6 Calculation of BINP Project Costs

FIGURE G.7**Simulation of an Income Transfer Equivalent to the Project Cost****Annual household income transfer (US\$)**

Project cost in 1998/participating children =
 $(1,459,000 / 49,023) = 29.8$.
 Discount a 15% administrative cost for the transfer:
 $29.8 * 0.85 = 25.3$

Additional per capita expenditure per month (Taka)

Discount a saving rate of 10%. Multiply by the exchange rate
 (48.5 Taka per US\$). Divide by 12 months and average
 household size (5.4).
 $(25.3 * 0.9 * 48.5) / (12 * 5.4) = 17$

Extra calories consumed per child per day (kcal)

Use calories elasticity of expenditure (0.363), mean per child
 expenditure (652) and mean calories consumption of children
 6 to 23 (917).
 $0.363 * ((17 + 652) / 652) - 1 * 917 = 8.7$

Extra weight gain per child per month (grams)

20 calories are required for 1 gram weight change:
 $8.7 / 20 * 30 = 13$

Reduction in prevalence of underweight (%)


1 kg weight gain is required for an increase of
 0.86 in average z-score. The year change in
 z-score is: $13 / 1000 * 12 / 0.86 = 0.13$.
 With the use of a probit function this is
 equivalent to a reduction in malnutrition of **5.4%**.

Number of lives saved (n)

Using Pelletier and Frongillo method, this
 weight gain and changes in z-scores and
 malnutrition rates results in **258** deaths
 averted.

Cost of upgrading a
 malnourished child (US\$)
 Project cost/children upgraded.
 $1459 / 0.054 / 49023 = 555$

Cost of saving 1 life (US\$)
 Project costs/number of
 lives saved.
 $1458 / 258 = 5654$

FIGURE G.8**Simulation of a Rice Ration Equivalent to the Project Cost**

ANNEX H: DFID AND WORLD BANK PROGRAMS IN BANGLADESH

This annex provides data on selected interventions mentioned in the report.

Rural Electrification

When the Rural Electrification Board (REB) was created in 1976, only 3 percent of the rural population had access to electricity. REB developed a plan for rural electrification based on cooperatives, called *Palli Bidyut Samities*. The plan had five phases that would complete rural electrification by 2005. Different phases were supported by different donors, including USAID, Finland, Norway, Japan, and CIDA, as well as the World Bank, which has had three projects, shown in table H.1.

There have been some implementation difficulties. Under REII there were financial problems, as REB was not paying its debt to government, as it in turn was not receiving payments from the cooperatives. In addition, RE & REII fell short of connection targets, but REIII exceeded them.

DHS data show the percentage of rural households with electricity to have risen from 5 percent in 1993 to 10 percent in 1996 and 15 percent in 1999

Figure H.1 shows the strong correlation between under-five mortality at all ages, with infants of households with no electricity 50 percent more likely to die than those with electricity, rising to an increased change of 75 percent for children. Of course, electrification is correlated with many household characteristics that are also correlated with mortality, notably household economic well-being, but also location, which may proxy for access to other facilities and general environmental conditions.

Multivariate analysis takes into account these additional factors. Even once they are controlled

for, electrification has a significant impact in reducing mortality. These econometric estimates are likely to underestimate the impact of electrification since electrification will also affect mortality indirectly, notably by the increase in economic opportunities such as irrigation and small businesses.

The ICR for REL III provides cost benefit calculations for four new cooperatives. The total cost for 93,214 connections, of which 80,210 are residential, is given as Tk1, 686 million. This a discounted valued, presumably discounted back to the year before the initial investment, 1991. Thus, to arrive at the capital cost in U.S. dollars for the year 2000, this figure is multiplied by a CPI-based inflator, and then divided by the exchange rate for that year:

$$\begin{aligned}\text{Capital cost in 2000 US\$} &= 1,686 \times (\text{cpi in 2000} / \\ &\quad \text{CPI in 1991}) / \\ &\quad (\text{Tk per US\$ in 2000}) \\ &= 1,686 \times (129 / 82) 52 \\ &= 550\end{aligned}$$

The cost of a household connection is estimated on a pro rata basis:

$$\begin{aligned}\text{Cost of h/h connection} &= \$550 \times (80,210 / 93,214) \\ &= \$470\end{aligned}$$

This calculation thus confirms that during the 1990s the cost per connection was approximately US\$500 per household. However, since less-accessible (and hence more costly) households are now being connected, the cost per connection has risen.

The Female Secondary School Assistance Project

The stipend program grew out of a USAID-supported, NGO-implemented program in one

TABLE H.1

World Bank Rural Electrification Projects

	Period	IDA amount	Connections
Rural electrification	1982–86	US\$49 million ^a	70,953 residential connections, 7,436 commercial, and 1,145 irrigation.
Rural electrification II	1985–94	US\$80 million	85,766 residential connections, 16,004 commercial, 2,406 irrigation.
Rural electrification III	1990–99	US\$110 million	216,791 new residential connections and 140,066 renovated; 45,875 commercial new and renovated, 9,367 irrigation new and renovated.

a. US\$40 million initial credit and US\$90 million supplemental credit.

upazilla in the early 1980s. The program was subsequently expanded, reaching seven upazillas by the time NORAD took it over in 1992. The World Bank program, under the Female Secondary School Assistance Project (FSSAP), began in 110 upazillas the following year. The Asian Development Bank also began its support in 1993, providing finance for 53 upazillas. In 1994, the government decided to expand the program, which had proved very popular, to all 410 rural upazillas, meeting program costs from its own funds in areas not covered by the three external agencies. Figure H.2 shows the number of scholarships supported by each agency; comparable data for GoB are not available. Under FSSAP and the follow-on FSSAP II, the World Bank financed 6.5 million girl-years of stipends over the period 1994–2002, equivalent to paying stipends for close to 2 million girls attending secondary school.

The scheme is administered through four separate project offices. In principle, the scheme should be the same in each upazilla, regardless of which funding source it falls under. However, some projects have additional components, such as the component to improve school facilities under the Bank project. In addition, the strength of monitoring and supervision appears to vary (e.g., it has not been possible to obtain data on the number of beneficiaries under the government scheme).


The key characteristics of the program are as follows:

- The stipend is paid to all girls attending rural secondary schools which are enrolled in the program. The stipend is paid into a bank account in the girl's name (where there are no local branches of the bank administering the scheme, it opens booths for this purpose). Tuition is paid directly to the school.
- The school has to certify that the girl attends school on at least 75 percent of school days, and that she is attaining passing grades (45 percent).
- The girl must remain unmarried.


The stipend, which increases by grade, is intended to cover a proportion of the direct costs of the girl's education. It does not fully cover these costs, especially as it has remained the same in nominal terms since the scheme began.

The stipend has not changed in nominal terms over the lifetime of the project. Including the two exam payments of Tk 250, the total amount paid per girl during the course of her education is Tk 3,914 (using a simple average of

FIGURE H.1

Bivariate Relationship
Mortality and Electrification

Source: Calculated from DHS data.

FIGURE H.2 Number of Stipends Financed by Each Agency

government and nongovernmental tuition fees, table H.2). At the 2000 exchange rate this amount equals US\$75.30. Allowing an administrative markup of 20 percent¹ brings the cost per girl to US\$90.40.

An alternative way to arrive at the estimated cost per girl is to divide the cost of the World Bank FSSAP, Tk 4,518.2 million, by the number of girl years supported by the project (4.904 million), which gives a figure of Tk 920 per girl year, which is US\$17.8 per girl year. Since the stipend runs for five years, this multiplies to US\$88.60 per girl. Hence, both approaches result in an estimate of US\$90 in stipend payments per girl during the course of her secondary education.

Female secondary enrollment has grown rapidly since the stipend has been introduced, at a rate of over 10 percent a year since 1993. However, the important question is the extent to which this expansion in female enrollments has been the result of the stipend program. An upper limit of the program impact is given by the whole rise in female secondary enrollments in rural areas since 1993. But this figure would clearly be an overestimate. Enrollments had anyhow been rising before the introduction of

stipends. The OED PPAR of FSSAP suggested that other factors may have been more important than the stipend, notably (1) the increasing number of girls completing primary school, which meant that girls enrollment at secondary level grew much more rapidly than it did for boys over the decade before the program, and (2) free tuition for girls in grades 6-8, which was established in 1992-93. Nonetheless, the OED study concludes that “project probably had an effect on enrollments” (World Bank 2003 p.10).

Female transition rate jumped from under 60 percent in 1990-91 to over 70 percent in 1992, but then again to 79 percent in 1994 and climb-

TABLE H.2**Stipend Payments under FSSAP (Tk)**

Grade	Stipend (monthly)	Tuition (govt.)	Tuition (nongovt.)	SSC exam fee
6	25	10	15	..
7	30	12	15	..
8	35	12	15	..
9	60	15	20	250
10	60	15	20	250

Source: FSSAP ICR, p. 31.

ing above 80 percent in the following years (World Bank 2000 p. 78).

A more rigorous attempt to determine project impact used econometric estimating exploiting the fact that the program was rolled out over time. This study found a significant impact of the program on enrollments using two different methods: the first, using household data, showed a 12 percent increase in enrollments for each year the program was in operation. The second approach, using school-level data, showed an initial impact of 8 percent, falling to 4 percent in subsequent years. An earlier paper by a member of the same research team addresses this point more specifically, suggesting that the program caused an increase in annual enrollments of 2 percent above the trend prevailing before the program's introduction.

These results need to be assessed against the actual increase that has taken place to know what share of the enrollment increase is attributable to stipends.

A variety of methods may be used to gauge this last figure, that is the increment in enrollments on account of stipends. The upper limit is to assume that the whole increase is the result of the stipend, which is clearly an overestimate.



There are alternative means of constructing the "without stipend enrollment figure." Options are to base the counterfactual on the growth of enrollments among rural girls prior to the introduction of the stipend, enrollment growth for rural boys during the program period, or enrollment growth for urban girls during the program period. Finally, estimates can be made using the impact study mentioned in the previous paragraph.

Unfortunately data are available for girls and boys enrollments, and rural and urban, but not the two disaggregations together. Hence single difference estimates can be made comparing boys versus girls (7.4 percent), before versus after for girls (-0.7 percent as the rate of increase was higher prior to the program), and double difference (0.3 percent); see table H.3. Finally, using the econometric estimate of a 2 percent incremental growth rate suggests that just under one-fifth of enrollment growth is the result of the program. This latter estimate, which falls in the rather wide range given by the simple estimates (from none to two-thirds), is used for the cost-effectiveness calculations.

A Note on DFID Aid to Bangladesh

Figures H.3 and H.4 show two figures related to U.K. aid to Bangladesh. Figure H.3 shows the substantial sectoral shift that has taken place during the 1990s, away from infrastructure, partly to health. Figure H.4 shows the number of separately funded health projects, showing that despite the move to a sector wide approach, there in fact remain a substantial number of discrete activities.

TABLE H.3	Growth in Secondary Enrollments (percent)		
	Boys	Girls	Difference
1986–92	4.2	11.3	7.1
1993–99	3.2	10.6	7.4
Double difference			0.3

FIGURE H.3 Sectoral Composition of DFID Aid to Bangladesh**FIGURE H.4** Number of DFID Health Projects

ANNEX I. AGRICULTURAL PRODUCTION, NATURAL DISASTERS, SEASONALITY, AND NUTRITIONAL OUTCOMES

Trends in Rice Production

Production of rice, the main food crop in Bangladesh,¹ has increased by over 60 percent over the past four decades as a result of a 50 percent increase in yields and a close to 25 percent increase in the area grown (figure I.1). Most of the increase in production is due to the introduction of high-yielding varieties, which, among other things, have enabled production of rice during different seasons.

Cropping Seasons


Bangladesh has two cropping seasons, *Kharif* and *Rabi*. *Kharif* lasts from April through November and is divided into early *Kharif* and late *Kharif*, the former being the pre-monsoon months of April through July and the latter June through November. *Rabi*, the winter season, lasts from November through May and is colder and drier. Figure I.2 provides an overview of crops, seasons, and weather.

Each season is suitable for producing different varieties of rice. The main rice crop is *Aman*, which is grown during late *Kharif*, followed by *Aus*, which is grown during early *Kharif*, and *Boro* rice grown during the *Rabi* season. The former variety is the largest in terms of area grown whereas *Boro* rice has the highest yield and the highest output. *Aus* rice is the smallest in term of area grown, production, and yield (table I.1).

The main determinant of the cropping pattern in Bangladesh is land elevation. It affects the annual extent and duration of flooding, which in turn influences cropping calendars. Land elevation similarly influences the number of crops grown annually in specific areas. For instance, about 71 percent of the total cultivable area is in the highlands and medium highlands where two to three crops are grown each year. By contrast,

only one or two crops are grown in the lowlands (Ateng 1998, p. 144). Of the net cropped area² in Bangladesh, 36 percent is single cropped, 51 percent double cropped, and 13 percent triple cropped (annually). That gives a cropping intensity of 177 percent (tables I.2 and I.3; Mian and others 2001, p. 37). Despite the lower prevalence of triple cropping, this area has expanded from around 480,000 hectares in 1965–69 to an average of 968,000 hectares during the triennium ending 1996–97. Thus, it accounts for an important part of production increase in this period.

The increase in multiple cropping has occurred as a result of the introduction of new technology spearheaded by the green revolution (Alauddin and Hossain 2001: 34). Whereas rice cultivation has traditionally been determined by the monsoon, the spread of small-scale, mechanically powered irrigation equipments, HYVs (high-yielding varieties), *Boro* technology and an irrigation policy, which enables farmers to invest in shallow tube wells sited at their discretion, led to an annual surge in *Rabi* season *Boro* rice production (Ateng 1998, p. 144; Rogal, Harriss, and Bose 1999, p. 99). Therefore, while MV (modern variety) rice was introduced in the late 1960s, diffusion in the dry season really took off in the mid-1980s along with changes in government policies in favor of privatization in the procurement and distribution of small-scale irrigation equipment and chemical fertilizers, liberalization of trade, and reduction in tariffs for imported agricultural equipment. Consequently, while 63 percent of the rice harvest today is monsoon-related, the remaining 37 percent is produced in the dry *Rabi* season when farmers grow HYV *Boro* rice using supplementary groundwater irrigation. This type of production has experienced a 6.9 percent annual growth during 1972–73 to 1993–94 com-

FIGURE I.1
Rice Production, Area, Yield, 1961–2002


Source: IRRI 2004

FIGURE I.2
Crop Seasons and Weather

	Jan	Feb	March	April	May	June	July	Aug	Sept	Oct	Nov	Dec
Crop seasons	Rabi Season—Boro rice										Rabi season—Boro rice	
				Early Karif season – Aus rice								
						Late Kharif season—Transplanted Aman rice						
Harvest	T. aman			HYV Boro			Aus					B. aman T. aman
Season	Dry season (and pre-monsoon hot season)					Monsoon season				Dry season		
Average rainfall, mm (Dhaka)	19	33	56	104	190	317	439	309	240	175	33	5
Average rainfall, mm (Sylhet)	10	31	137	396	553	834	827	633	524	250	27	10

Source: Adapted from Ateng 1998, supplemented from Bangladesh Bureau of Statistics 2000; National Environment Agency, Singapore.

pared with 2.8 percent annual rice production over the same time (Ateng 1998, p. 144; Rogal, Harriss, and Bose 1999, p. 99).

Despite the increase in multiple cropping following the spread of irrigation and production of dry season *Boro* rice, the possibilities for *triple* cropping—and in particular for growing three rice crops—are limited. This is so first because the growing seasons of both *Aus* and *Aman* rice overlap with that of *Boro* rice (see figure I.2), and second because dry season *Boro* rice often replaces other dry season non-cereal crops. Some argue that at least in lower lands, expansion of irrigated *Boro* rice cultivation has taken place on land that would otherwise have been involved in a complex rain-fed double and triple cropping pattern involving broadcast *Aus*,³ broadcast *Aman*, and a range of other winter crops. Since most non-cereal crops are grown on non-irrigated land and compete for land in the dry season these crops are thus liable to be replaced with the expansion of irrigation (Jaim 1984, p. 36; Ateng 1998, p. 146–8). This argument is supported by looking at the proportion of *Aus* and *Boro* rice cropped areas under triple cropping. Ten percent of MV *Aus* cropped area is under rice-only triple cropping and 17 percent is under rice and other triple cropping, while 7 percent of *Boro* rice is under rice-only triple cropping, and only 1 percent under rice and other triple cropping (Mahmud, Rahman, and Zohir 1994, p. 30).

Because *Aman* rice is the largest crop in terms of area grown and makes up close to 60 percent of rice grown area (table I.1), November–December remains the main harvest season with the widest food availability. May–June, when *Boro* rice is harvested, is the second most important harvest.

Aggregate Output, Disasters, and Daily Energy Supply

As rice is the major agricultural crop, the trend in daily energy supply per capita closely follows rice production, notably the upsurge in the late 1990s (figure I.3). The figure also shows the major disasters during this period, which for the most part seem to have not had a large an impact an aggregate production. Daily energy supply also appears

TABLE I.1		Rice: Production, Area, Yield by Season, 1992–94 Average			
	Area (000 ha.)	Share of total (percent)	Output (000 t)	Share of total (percent)	Yield (t/ha.)
Aus	1,767	17.4	733	6.1	1.15
Aman	5,763	56.9	4,898	40.9	1.64
Boro	2,605	25.7	6,342	53.0	2.58
Total / average	10,135	100	11,973	100	1.8

Source: Baffes and Gautam 1996.

TABLE I.2		Land Utilization by District (as percent of total net area cropped)		
		Single cropped	Double cropped	Triple cropped
1989/90				
Chittagong	35.0	52.4	12.6	
Dhaka	41.5	40.9	17.6	
Khulna	72.2	23.8	3.9	
Rajshahi	58.6	34.5	6.9	
All Bangladesh	43.5	44.5	11.9	
1997/98				
Chittagong	35.7	58.4	13.2	
Dhaka	39.6	56.9	13.6	
Kulna	66.9	25.5	4.1	
Rjshashi	52.4	44.3	7.7	
All Bangladesh	39.0	56.1	13.8	

Source: Chowdhury and Zulfikar 2001.


not to suffer during disasters (table I.4), sometimes even increasing, presumably reflecting relief food aid.

Seasonality and Nutrition

Cropping patterns determine harvest seasons, which in turn influence work, employment, food

TABLE I.3		Land Use by Rice Sort, 1993		
		Single cropped	Double cropped	Triple cropped
Aus	2	71	27	
Boro	33	58	8	

Source: Mahmud et al. 1994.

FIGURE 1.3 Rice Output and Daily Energy Supply

Source: www.em-dat.net

intake, and nutrition in rural areas. As the largest crop, *Aman* is the most important in this respect. Demand for agricultural labor is highest during the *Aman* harvest. Also, opportunities for employment are enhanced during production and harvest of *Boro* and *Aus*—for instance during the months of January and March when the transplantation of HYV *Boro* takes place (Pitt and Khandker 2002: 12). The introduction of new

agricultural technology has made significant contribution to employment creation, rather than the reverse, especially HYV *Boro* rice (Alauddin and Tisdell 1998: 86–9).

Seasonal patterns affect nutrition in three ways: (1) food availability, (2) labor requirements, and (3) rainfall affecting health, with children particularly prone to disease in the rainy season. For households working their own land the harvest season is one of extra effort, as food is not widely available. But for wage laborers this is a time of increased income, though rice prices will be higher until harvest is completed.

Children

Two studies (Brown 1982; Bloem, Moench-Pfanner, and Panagides 2003) conducted more than two decades apart focused on seasonal variations in children's nutritional status. Both found significant differences across seasons. A third study (Becker and Weng 1998) found significant seasonal variation in neonatal, post-neonatal, and child mortality.

Using various anthropometric indicators,⁴ Brown (1982) found the period of poorest nu-

TABLE 1.4 Top 10 Disasters in Bangladesh by Number of People Affected

Disaster	Date	Affected (millions)
Flood	July 1987	73.0
Flood	August 1988	73.0
Flood	July 1974	38.0
Flood	June 2004	33.6
Flood	May 1984	30.0
Drought	July 1983	20.0
Flood	July 1968	15.9
Wind storm	May 1965	15.6
Wind storm	April 1991	15.4
Flood	July 1998	15.0

tritional status to begin with the height of the monsoon rains in the *Kharif* season continuing until roughly the time of the *Aman* rice harvest. Measuring weight for age, arm circumference, and triceps skinfold thickness, they found a large decline between June and September. The latter was the month with the highest prevalence of child underweight. August was found to be the month with by far the lowest mean percentage of expected monthly increments of weight for age (<10 percent) (Brown 1982: 308–9).

Using NSP data, Bloem, Moench-Pfanner, and Panagides (2003) also found that wasting (WHZ < -2 SD) in children under five peaked between June to September with 28 percent of children under five being wasted. The prevalence of wasting in children aged 0–23 months rose from “serious severity” to “critical severity” in August/September (Bloem, Moench-Pfanner, and Panagides 2003, p. 90–1). Such findings were complemented by other evidence that food scarcity is greatest in the months from August to October. For instance, in 1992, 1993, and 1994, September was the month with the highest percentage of households that took a loan for food or made a distress sale of assets (2003: 55).

OED’s own analysis of WAZ for under-fives is shown in figure I.4. The pattern shown confirms

that the lean season falls between April and November, with the start and end points varying a little from year to year.

These findings indicate a temporal association between cropping patterns and children’s nutritional status. Seasonality in nutrition, however, is furthered by the greater prevalence of floods and diseases during the monsoon in the late *Kharif* season. Floods are extremely common in Bangladesh and affect hundreds of thousands of people each year, particularly during the months of June, July, and August (table I.5). Of the 47 floods that occurred between 1970 and 2004, 39 took place between June and September. Nine of 19 diarrhoeal epidemics similarly took place during those months.

The link between floods and health is indicated by the fact that the monsoon season has the highest occurrence of diarrhoeal deaths, and highest levels of child mortality (Becker and Weng 1998). Brown, Black, and Becker also find the prevalence of certain infectious diseases to be generally higher during that season (1982: 312). Finally, examining the 1998 floods, Del Ninno and others (2001) found that children in severely exposed villages had a slightly increased illness risk and for those in severely exposed villages the risk more than trebled; the incidence of fever rose by 55 percent and that of

FIGURE I.4

Seasonal Patterns in Weight for Age Z-score for Under-Fives




TABLE 1.5**Disasters by Season**

	March-May	June-September	October-November
Floods	7	39	1
Diarrhoeal epidemics	6	9	4

Source: EM Dat 2004.


diarrhea by 25 percent, which will affect nutritional outcomes.

Women

The OED analysis of the NSP data shows that seasonal variations in underweight are not the same for children and women. Whereas children's weight for age is lowest during the monsoon in June-July and the "hungry months" preceding the *Aman* harvest, and highest during November and December, women's Body Mass Index (BMI) peaks in August-September and is lowest in December. Maternal calorie deficiency (CED) is most severe in December and January and least so in August-September (although seasonal variation is not found to be very large: 43 percent in December-January versus 36 percent in

August-September) (Bloem, Moench-Pfanner, and Panagides 2003).

Such seasonal variations in female malnourishment are most likely explained by the significant increase in women's workload. A qualitative study of women's schedules and time constraints found them to vary according to the agricultural season as harvest seasons and rice processing significantly affects time use (Levinson and others 2002). The busiest seasons for women are the crop harvest seasons from mid-April to mid-June and from mid-November to mid-January. The rice harvest months of mid-April to mid-May and mid-November to mid-December (figure I.5) are the times when women generally prepare the least number of meals because they lack time to do so. Most women in the study said that they managed to complete their tasks only by sleeping, resting, and bathing less. Finally, the study found that women considered time a more serious problem with respect to self-care than to childcare. For example, they regarded eating more food during pregnancy as difficult given their time constraints, whereas providing complementary food to their children or giving them more than three meals a day was given higher priority.

FIGURE I.5**Seasonal Patterns of Women's Body Mass Index**

Background and Rationale

Two of the eight Millennium Development Goals (MDGs) refer to maternal and child health, and child malnutrition is a Goal 1 indicator. The results-based rationale of the MDGs requires understanding the main drivers behind changes in MDG indicators: what can government policy and interventions, assisted by external partners, do to accelerate the pace of improvement and so secure the achievement of the goals?

The impact studies being undertaken by OED, under the OED-DFID partnership agreement, seek to address this issue, focusing in this case on maternal and child health (MCH) outcomes in Bangladesh. The study will build upon the methods developed during the first impact evaluation carried out under the partnership agreement, which analyzed the impact of the external support to basic education in Ghana from the Bank, DFID, and USAID.¹ This approach is based on the combination of rigorous statistical analysis of outcomes with contextual analysis based on qualitative material. A joint evaluation under the partnership agreement interventions supported by both the Bank and DFID will be included in the study.

The focus of the study will not, however, be restricted to interventions in health, nutrition and population (HNP), since MCH outcomes are also affected by activities in other sectors. In the words of the 2004 *World Development Report* “the determinants of supply and demand operate through many channels” (WDR, 2004, p.27). OED’s own review of the MDGs similarly stressed the need for multi-sectoral strategies to achieve single-sector targets.² DFID’s target strategy paper, *Better Health for Poor People*, states that, “good health is determined by many factors, the most important of which is reaching

and sustaining a reasonable level of income and consumption. Female education, clean water and, most important, sanitation have an important and well documented impact on (in particular) child survival. Female empowerment is vital to enable women and families access to health care” (DFID, 2000, p. 9).

Bangladesh has been chosen for a number of reasons. It is a country that has made impressive progress in reducing both under five mortality and fertility. External partners, including both the World Bank and DFID, have been active in supporting large-scale interventions with specific objectives to improve maternal and child health. Improved access to health services especially for poor women and children was included under the first objective in DFID’s country strategy for Bangladesh formulated in 1998. During recent years external interventions in the population and health sectors have been coordinated under a sector program in which both DFID and the Bank have been active partners. Finally, there are a substantial number of large data sets appropriate for the analysis.

The Bangladesh Context

Bangladesh presents a mixed picture with respect to MCH indicators. Over the 1990s infant and child mortality have been halved: unlike most countries, Bangladesh has a sufficient rate of progress to meet the MDG target of a two-thirds reduction by 2015. Fertility also fell rapidly from the 1980s, although there are fears in some quarters that the indicator has now reached a plateau. However, despite low levels of fertility, maternal mortality is high: at between 320 and 400 per 100,000 live births, it is greater than that in both India and Pakistan and six times that in Sri Lanka. Intermediate indicators show a similar

TABLE J.1

MCH-Related Millennium Development Goals

(a) Goal	Target	Indicators
Goal 1: Eradicate extreme poverty and hunger	Target 2: Halve, between 1990 and 2015, the proportion of people who suffer from hunger	<i>Prevalence of underweight children (under five years of age)</i> <i>Proportion of population below minimum level of dietary energy consumption</i>
Goal 4: Reduce child mortality	Target 5: Reduce by two-thirds, between 1990 and 2015, the under-five mortality rate	<i>Under-five mortality rate</i> <i>Infant mortality rate</i> <i>Proportion of one-year-old children immunized against measles</i>
Goal 5: Improve maternal health	Target 6: Reduce by three-quarters, between 1990 and 2015, the maternal mortality ratio	<i>Maternal mortality ratio</i> <i>Proportion of births attended by skilled health personnel</i>

discrepancy. Immunization rates have grown rapidly in Bangladesh and are today notably higher than those in neighboring countries. But only 13 percent of births are delivered by skilled birth attendants, compared to close to half in India.

The study will focus on interventions from several sectors. The chosen interventions will be those supported by DFID and the World Bank, and other donors subject to the required information being available. Amongst these sectors will be health and population, which has accounted for 17 percent of DFID aid to Bangladesh in recent years (Annex J.1b) and, together with nutrition, 7 percent of Bank lending (Annex J.1a). As described below, the other sectors are to be identified based on an analysis of the determinants of MCH outcomes.

On the Bank's part there have been five Health and Population projects in Bangladesh: I (1974-80), II (1979-86), III (1986-92), IV (1992-98) and the Health and Population Program Project (ongoing since 1998), the last three of which have been co-financed by DFID. In addition there is the Bangladesh Integrated Nutrition Project (BINP, 1995-02), which is being followed up by the National Nutrition Project (NNP, ongoing since 2000). All these projects have specific objectives related to maternal and child health. The study will include, amongst other interventions, PHP III, IV and HPSP, and BINP.

DFID's support for HPSP has been principally to Strengthen Health and Population for the Less Advantaged (SHAPLA), through two mechanisms: (i) time slice financing of the HPSP budget, and (ii) project support focused in key areas of HPSP to strengthen government's implementation capacity and to minimize risk. Both Bank and DFID financing to the population and health sector has had several components, which will be assessed separately for their inclusion in the study. Other donor interventions to be included in the study will be provisionally identified at the design stage.

In line with the multi-sectoral philosophy of the MDGs (i.e., the notion that the relevant interventions for each outcome are in a range of sectors) the study will consider interventions from outside the HNP sector. For example, income-generation activities will also positively affect MCH outcomes, as will improved water supply and better education. The analysis of other sectors will be carried out through a two stage approach described below.

Evaluation Questions

The evaluation questions to be addressed are:

- (a) What is happening to maternal and child health indicators (including nutrition and fertility) in Bangladesh? Are the poor sharing in the progress that is being made?

- (b) What have been the main determinants of MCH outcomes in Bangladesh over this period, and which of these have accounted for substantial changes?
- (c) What have been the primary drivers of changes in the determinants of MCH outcomes since the mid-80s?³
- (d) What have been the range of government and externally-supported interventions that may have affected the changes in the determinants of MCH outcomes?
- (e) To the extent that interventions have brought about positive impacts, have they done so in a cost effective manner?

Evaluation Approach and Data Requirements

The analytical framework for the evaluation is provided by the combination of two widely-adopted frameworks—UNICEF’s conceptual framework of the causes of malnutrition⁴ and the Mosley-Chen analytical framework of child survival⁵—together with the insights from household models of the demand for health used in economics.⁶ The first two frameworks have the same outcome variables (malnutrition and mortality) and essentially the same structure, in which the basic causes come from a country’s socio-economic and institutional conditions, but these are mediated through a number of proximate causes observed at the community, household and individual levels. Maternal health is both a determinant of child welfare outcomes and an outcome in its own right. Economic models also highlight the importance of behavioral factors (“tastes and preferences”) in affecting demand for health services.

The framework identifies the determinants of MCH outcomes in broad terms thus providing a theoretical basis for the selection of variables for modeling these outcomes. These determinants have been examined more closely in a meta-analysis of infant and child mortality and nutrition, prepared as a background paper for this study (see Appendix J.3 for a summary of determinants identified by this analysis).

The adoption of a theory-based evaluation approach will direct attention to further assump-

tions contained in project design. For example, since health and nutrition interventions attempt to bring about behavior change, they are partly built on the assumption of irrational beliefs regarding health and nutrition practices. But the data show some discrepancies between knowledge and practices, suggesting that projects may change knowledge but not practices. The most recent progress report for Bangladesh’s Health and Population Sector Program notes scant improvement in service use. The study will use both quantitative and qualitative methods to analyze issues such as patterns of service use and the rationale for various health and nutritional practices.

The determinants given in a regression model of MCH outcomes are not usually direct measures of policies or programs. The task for an impact evaluation is to link outputs from policies and programs with the determinants and hence to the final outcomes. This study will examine the link between MCH outcomes and DFID and Bank-supported interventions, together with those of other donors where possible, through a two stage approach. A regression-based decomposition analysis of the determinants of MCH outcomes will be used to identify which factors have been most important in changing outcomes in Bangladesh. This analysis will be used to identify which DFID, Bank and other interventions have directly supported improvements in MCH. The most important interventions will be selected for a more detailed analysis, to quantify the size and cost effectiveness of their impact. In addition to HNP, it is envisaged that two or three other sectors will be selected. The activities of the DFID and the Bank within these sectors will be examined, along with those of other donors where this proves feasible. The final selection of sectors will be an outcome of the first stage of the analysis.

An exception to this approach is BINP, which has an unusually rich evaluation dataset comprising a baseline, mid-term, and endline surveys for both project sites and control communities. The data collected can be mapped onto the logic of the project, since intermediate process indicators relating to behavioral factors were included in the surveys in addition to outcome

data and socio-economic status (SES) variables. These data are available to the study team together with the relevant documentation. Save the Children, which has been critical of the approach adopted by BINP, conducted their own survey in 2002 and these data will also be available to the study team.

Several other nationally-representative data sets exist for the relevant period. These are listed in Appendix 2. Surveys likely to be used are:

- Demographic Health Surveys: 1993/94, 1996/97 and 1999/2000
- Household income and expenditure surveys: 1991/92, 1995/96 and 2000
- Helen Keller International Nutritional Surveillance Project
- Maternal mortality survey, 2003.

The first two of these surveys have been acquired for the study. The other two are not in the public domain, but can be analyzed through collaborative arrangements with their owners.

A review of existing material shall be undertaken during design to determine to what extent the analysis can rely on secondary data and to what extent commissioned studies are required. Relevant areas of analysis are: (a) trends in government expenditure,⁷ (b) facility and staffing survey, (c) demand for health services, and (d) behavioral changes related to health and nutritional practices. At least two of these issues will be examined through qualitative fieldwork, using participatory analysis of such issues as perception of service quality and the rationale underling health and feeding practices.

Collaboration with Other Agencies

Collaboration shall be sought with relevant government officials or research institutions in Bangladesh. Commissioned studies from local

organizations may include a component of capacity development through supported learning-by-doing. DFID staff in the Dhaka office shall facilitate the inclusion of DFID-supported interventions in the study and assist, together with Bank staff, liaison with other agencies as necessary.

Schedule and Task Management

The study will be undertaken in the following stages:

- Inception—comprising an inception field visit (February 2004), preparation of the approach paper followed by preliminary fieldwork (April 2004) and development of the study design (April-May).
- Analysis first phase—analysis of household data sets of the determinants of health, nutrition, and fertility outcomes. Background studies will be carried out in parallel. The results of these studies will be presented in Dhaka in late July or early August 2004 to feed into the discussions concerning future support. Identification of projects to be analyzed by the study will be carried out in this phase.
- Analysis second phase—collection and analysis of project-level data from DFID, the Bank and other donors as possible.
- Report draft for OED management review to be ready by late October 2004, with submission to CODE of the final report targeted for January 2005.

The evaluation will be carried out by a team of OED staff and consultants with the assistance of Bangladeshi government officials, DFID staff under the Task Management of Howard White (OEDST). External peer reviewers will be appointed to review the proposed evaluation design and draft final report.

APPENDIX J.1 a		Overview of Bank Operations in Bangladesh Since 1990	
Sector	Commitments		
	US\$ million	Percent	
Economic policy	224.8	4.4	
Education	419.9	8.1	
Energy and mining	1,252.7	24.3	
Environment	151.3	2.9	
Financial sector	315.2	6.1	
Global information/communications technology	39.8	0.8	
HNP	349.3	6.8	
Private sector development	362.8	7.0	
Public sector governance	185.9	3.6	
Rural sector	908.5	17.6	
Transport	866.0	16.8	
Urban development	48.4	0.9	
Water supply and sanitation	35.8	0.7	
Total	5,160.4	100.0	

Source: calculated from OED database.

APPENDIX J.1 b		Overview of DFID Operations in Bangladesh, 1998–2001 (£ millions)				
	1998/99	1999/00	2000/01	2001/02	Total	Share
Education	7	10	10	9	36	10.6
Population and health	13	18	14	13	58	17.0
Natural resources	10	22	29	31	92	27.0
Roads/bridges	11	8	11	11	41	12.0
Water and sanitation	0	4	7	7	18	5.3
Energy	6	14	7	7	34	10.0
Small business/micro-credit	2	6	5	5	18	5.3
Good government	4	5	5	2	16	4.7
Other	5	6	9	8	28	8.2
Total	58	93	97	93	341	100.0

Source: DFID Bangladesh Country Strategy Paper, 1999: Annex 2.

APPENDIX J.2 List of Data Sets		
Survey	(b) Content	Availability
Demographic Health Survey	Child health and nutrition, child care practices, SES, comprehensive birth history. Includes some “women’s agency” questions.	Freely downloadable from Macro International website
BBS Health and demographic survey	As for DHS	To determine if in Bank or not, otherwise purchase from BBS
BINP evaluation data set	Nutritional status, beliefs and practices, SES	Available in Bank and clearance has been obtained for use, awaiting data
SCF Bangladesh nutrition data set	Nutritional status and household characteristics	Was promised sometime ago, need to follow up
Household Income and Expenditure Survey	Standard LSMS type survey, but no anthropometrics. Only most recent has health module, but all have health expenditures	Most recent acquired, others available in Bank and being followed up
Health facility survey	Not known yet	Acquired
Helen Keller International National Nutrition Surveillance Survey	Nutritional status + SES. Every two months since 1990, with nationally representative sample since 1996. Content changes but a common core.	Proposed to contract HK to conduct analysis.
Vital registration sample survey (BBS)	Births, deaths and basic household and individual characteristics. Cause of death (quite aggregated).	Can be purchased from BBS. Possibly analyze in collaboration with BIDS
National Nutrition Survey (BBS)	Anthropometric measurement, SES.	Can be purchased from BBS. Possibly analyze in collaboration with BIDS
UNICEF Multiple Indicators Cluster Survey	Range of social indicators, representative at district level.	Should be available from MICS website, for which clearance to download already obtained
UNICEF maternal health facility survey	Use, deliveries and materials availability	Statistics officer can perform analysis on request
Maternal Mortality Survey (USAID financed)	Maternal death (sisterhood and h/h based), SES	Not available but can influence form of analysis and receive results

Note: Though both BBS surveys began in the 1980s, data may only be available for the last one or two rounds. All the above are household surveys. Some are supplemented by community-level data collection of variables such as facilities and prices. These are all national surveys; there are also more local data, notably those collected in Matlab thana that may also be used.

APPENDIX J.3 Determinants of Outcome Variables			
	Child	Household	Community
Nutrition	<p>Child characteristics: gender, age, birth order, sickness, ethnicity, lagged height</p> <p>Care practices: number of antenatal visits, attended birth, breastfed</p>	<p>Economic characteristics: per capita income, asset ownership, public works participation, paternal labor supply, maternal labor supply, poverty status, land ownership, occupation, remittances</p> <p>Parental characteristics: mother's and father's education, parental height, mother's age, maternal nutrition knowledge, maternal literacy, maternal employment, marital status, ethnicity, religion</p> <p>Nutrition-related factors: per capita calories, per capita proteins, parental education, household size, mother's height, food prices</p> <p>Care-related factors: sibling help, adult help, mother works full time</p> <p>Demographic characteristics: household size and composition, no. of dependents, sex of household head</p>	<p>Basic characteristics: rural/urban, region</p> <p>Facilities: electricity, water, sanitation, travel time to health center, school, road conditions, hospital, TV, radio</p> <p>Other: famine, season, price of drugs, food prices, literacy, hospital beds</p> <p>Note: some studies use a community fixed effects model.</p>
Mortality	<p>Child characteristics: gender, age, ethnicity, birth order (first birth), birth interval, birth cohort, birth weight, length of pregnancy (premature), twin</p> <p>Care practices: (duration of) breastfeeding, antenatal care, attended birth, immunization, exposure to water-borne disease</p> <p>Note: some studies use interactive term between breastfeeding and water and sanitation variable</p>	<p>Economic characteristics: income (total or per capita), asset ownership, dwelling space, land ownership, parental employment group, paternal and maternal labor supply</p> <p>Parental characteristics: mother's and father's education, mother's age, mother's height, no. of children born to mother, maternal health knowledge, marital status, religion, ethnicity, caste</p> <p>Mortality-related factors: fate of previous children, fate of other relatives/household members</p> <p>Care-related: mother works full time, sibling/adult help</p> <p>Demographic characteristics: household size and composition, sex of household head</p> <p>Other: persons per room, migration status</p>	<p>Basic characteristics: rural/urban, region</p> <p>Facilities: electricity, water, sanitation, schools, health facility, health workers, doctors, paved road</p> <p>Other: level of child mortality in community, disease prevalence, population (growth rate), temperature and precipitation</p>
Fertility	Not applicable	<p>Economic characteristics: income per capita, asset ownership including, land ownership, occupation, employment status</p> <p>Preferences: female education, male education, contraception and abortion practices, religion, ethnicity, access to social security in old age (including land ownership), farm size</p> <p>Other factors: woman's status, woman's age, marriage age, mortality among children and other household members, breastfeeding duration</p>	<p>Basic characteristics: rural/urban, region</p> <p>Facilities: electricity, access to mass media (radio, TV), access to family planning facility, schools</p> <p>Other: mortality rates, disease prevalence, literacy</p>

Source: derived from a review of statistical studies.

ENDNOTES

Chapter 1

1. Data from the 2003 Maternal Mortality Survey were not available for analysis.

2. The under-five mortality rate in 1970 was 239 per 1,000 live births, compared with 202 for India, 181 for Pakistan (excluding Bangladesh), and just 100 for Sri Lanka.

3. Social indicators are often built on fairly shaky statistical foundations. However, as shown in Annex A, many surveys have been conducted in Bangladesh, and so the data are unusually reliable. The exception has been maternal mortality, but a new Maternal Mortality Survey will improve the situation.

4. Whether or not fertility decline continued into the 1990s is a matter of debate. This issue is explored in Chapter 3 and Annex F.

5. For example, the United Kingdom financed 15 separate activities under the Fourth Population and Health Project, totaling UK£15.5 million (close to US\$30 million at current exchange rates). Close to half of this total was for support to NGO programs, with substantial amounts also for strengthening nurse education (UK£2.8 million) and medical colleges (UK£3.1 million) (see Annex I).

6. Rural Electrification I: 1982–86; II: 1985–94; and III: 1990–99.

7. The World Bank has lent US\$234 million for flood relief and drainage programs in the last two decades, and DFID £59 million since 1991 (see Annex I).

8. The 1998 floods were far greater in terms of the affected area and infrastructure destroyed than those in 1988. But fewer than 1,000 people lost their lives in the 1998 floods compared with more than 2,400 in 1988. And only 600 died in the 2004 floods, which appeared to be at least comparable to those in 1998.

9. Among older children, drowning is now the major cause of death. But, as these figures show, these drownings are mainly accidents rather than the direct consequence of flooding. Figures on indirect deaths from flooding as a result of disease and malnutrition are less easy to obtain, though there is such an effect (for some estimates, see Del Ninno and others 1991).

10. Moore (2003) argues that the remote nature of many communities, resulting in control of local politics by a small number of dominant families, in part results from difficulties in moving around the country. Thus, it may be argued that improved infrastructure has a beneficial effect on political life, which in turn will improve service delivery.

11. The data from a fourth DHS, conducted in 2004, were not available for this study.

Chapter 2

1. Bangladesh has six administrative divisions, further subdivided into 64 districts. Each district has several thanas (formerly upazillas), of which there are 490 in total. Each (covering some 15–20 villages) is further subdivided into several unions.

2. One might wonder if the assessment a few years earlier can thus be faulted. While the earlier OED report missed the emerging success of the program, its findings reflected accepted opinion of the program at the time. Perceptions regarding the success of the Bangladesh program changed with the data from the BFS and CPS carried out in 1989.

3. For example, Caldwell and others (1999) and Kabeer (2001). Chapter 4 will examine the evidence regarding this debate in more detail.

4. See, for example, the staff appraisal report for the Bank's Fourth Population and Health Project.

5. Data from the Health Facility Surveys show a decline the proportion of households using government health and family planning services from 13 to 10 percent from 1999 to 2003 (Cockcroft Milne, and Anderson 2003).

6. A recent study found absenteeism among rural public health providers to average 26 percent, but was higher for doctors (40 percent), particularly in smaller subcenters (76 percent); see Chaudhury and Hammer 2003.

7. DFID is implementing a public-private partnership program under which the clinics are utilized by an NGO using a matching grant provided by DFID.

8. This figure includes support to NGO programs under the joint finance scheme (JFS).

9. Other than this contribution, the United Kingdom has not been a main supporter of immunization, though it did assist in the program to eliminate polio (see box 4.2).

10. The Maternal Mortality Survey reported an even higher share of 74.4 percent of births being attended by TBAs (NIPORT, Mitra and Associates, and ORC Macro 2003, p. 52).

11. The Safe Motherhood Initiative is an international effort launched in Nairobi in 1987 that is supported by several international agencies, including the World Bank.

Chapter 3

1. The usual pattern is that mortality falls much more rapidly for older children, so that the remaining deaths become concentrated first among infants and, within that category, among neonates (and for neonates, the first days of life are the most risky). This pattern is less marked in Bangladesh than elsewhere, with neonatal mortality falling as fast as that for post-nates (up to one month), and not too far behind that of children one to four years.

2. The poor are identified as the bottom 50 percent, ranked by a wealth index, in the 1992–93 DHS data. The wealth index poverty line from that survey was applied to the subsequent surveys so that data are comparable between years. This conclusion is robust to the choice of the poverty line; the concentration curve for the early 1990s has first-order dominance over that for the late 1990s up to around the seventh decile. Analysis shows that group reductions of mortality accounted for far more of the fall in mortality than did movements out of poverty (that is, into a lower mortality group); see Annex C.

3. Data were examined from 125 DHSs. Ranking these surveys by female: male mortality in descending order, the ratio from the three Bangladesh surveys ranks eighth, ninth, and twelfth—that is, in the worst 10 percent. There is no gender bias in neonatal mortality, with girls less likely to die than boys in the first month of life. A bias begins to emerge for postnatal mortality, although not as marked as that for child mortality.

4. The downward trend in disaster-prone areas, compared with the lack of a trend nationally, may be taken as crude evidence of the effectiveness of flood-protection measures mentioned in Chapter 1.

5. Direct estimates are based on a weighted average of the age-specific fertility rates of all women interviewed, whereas indirect measures utilize various proxy measures to estimate fertility.

6. The plateau has been attributed to a tempo effect. If women postpone births, this will temporarily decrease the total fertility rate (which is based on a synthetic cohort), but it will then rise again. Against the background of an underlying downward trend, the tempo effect would first accelerate the rate of decline, and then temporarily halt it (see Annex F).

Chapter 4

1. The dashed line has the same slope as the fitted line, but with the intercept adjusted to pass through the 1980s observation for Bangladesh. There is an assumption here that the outcome-income relationship is the same for different countries over time. This is clearly not so, most notably in the case of Bangladesh. The point of the analysis is to show how much Bangladesh departs from international norms, so as to pose the question as to why this is.

2. It is an upper estimate, since the calculations are based on a simple regression. This equation is undoubtedly mis-specified, resulting in omitted variable bias. Income is positively correlated with several determinants of these outcomes—such as education and immunization—so that there will be an upward bias on the estimated regression coefficient for income.

3. The cross-country analysis conducted for this report builds on earlier work by Smith and Haddad (2000), who estimated two models for the determinants of the proportion of children who are underweight. Their first model regresses nutritional status on what they call the “basic determinants” of per capita GDP and democracy (measured by the Freedom House political freedom index). The second model, of underlying determinants, includes access to safe water, female secondary school enrollments, the ratio of male to female life expectancy as a measure of gender inequality, and per capita dietary energy supply. The logic of the approach is that the latter variables, which directly affect nutritional status, depend in turn (though only in part) on the basic determinants. They then examine the relative importance of the different underlying variables in the explained change in nutrition. Female education is shown as the most important factor (43 percent), followed by food supply (21 percent), and then health environment (safe water, 19 percent), and finally women’s status, with 11 percent. Whereas Smith and Haddad’s estimates are calculated across the whole sample, those for this study are based on the values of the regressors for Bangladesh. Specifically, we calculated $b(x1-x0)/y$, where b is the estimated regression coefficient, x the value of the explanatory variable for Bangladesh in the

1980s (0) and the current decade (1), and y the relevant outcome.

4. Following Smith and Haddad (2000), these estimates are made excluding income from the model. Hence, some of the explained variation results from the income-driven component of these determinants, but there is also an independent element. To test for this independent element the equations were re-estimated including income per capita as an explanatory variable (see Annex B). An independent effect was found for most determinants, with the exception of gender equality and safe water in some specifications.

5. Bangladesh has six divisions: Barisal, Chittagong, Dhaka, Khulna, Rajshahi, and Sylhet. Sylhet was formerly part of Chittagong, so that Chittagong (including Sylhet) is used for the regional analysis.

6. Many care-related variables are difficult to include for reasons of endogeneity. Controlling for endogeneity of antenatal care shows it to be significant only if provided by a trained doctor or nurse. It is not possible to include breastfeeding in the equations, as the only available question ("never breastfeed") is a function of child health.

7. Analysis using DHS data measures economic well-being using a wealth index rather than income. Annex C analyzes the relationship between such wealth indexes and income/expenditure.

8. First-born children have a higher risk of infant death than do subsequent births. Lowering fertility means increasing the proportion of first-born children, exerting an upward influence on infant mortality.

9. When interacting the female dummy with the Chittagong dummy, the female dummy is no longer significant, but that for the interactive term is significantly positive.

10. The hazard ratio for the Cox regression using 1999 data was 2.05 (table C.7). Using pooled data, and a without-self mean of immunization to account for potential endogeneity, the ratio falls to 1.55.

11. Khan and Yoder (1998). The Bangladesh EPI office currently reports a figure of \$10 per child.

12. Khan and Yoder (1998) arrive at an estimate of \$136 per life saved.

13. Moreover, the cost of treatment of children who become sick can be shown to exceed the cost of avoiding sickness, so that immunization represents a saving of health expenditures.

14. Once immunization reaches a certain threshold, then there is a reduction in the cross-infection, so that unimmunized children are also less at risk of the disease. The threshold varies by disease but is mostly in the 80-90 percent range (Fine 1993).

15. In China between 1950 and 1980, the maternal mortality ratio declined from 1,500 to 115 per 100,000 live births. The vast majority of births during this period were delivered by minimally trained village birth attendants (backed up by a strong obstetric care and emergency referral network) (Levitt 1999).

16. However, the study of Goodland and others (2002) finds that training TBAs did not reduce post-partum infection, which they attribute to the importance of pre-existing reproductive tract infection. This is not something to which EOC would have made a difference.

17. Multivariate analysis suffers from the same problems illustrated in the bivariate analysis (table 4.3). That is, in the neonatal mortality regression, the coefficient for births attended by trained TBAs is significantly positive (that for untrained TBAs is significantly negative), but the interaction of the year with trained TBA is negative, and of sufficient magnitude to give trained TBAs an advantage over untrained by 1999. This result remains when a trivariate probit is estimated (to handle the endogeneity of antenatal care and choice of birth attendant). The trivariate results show which women use TTBA: the coefficient for the later period is highly significant, picking up the increased availability of TTBA, so is being a female household head, a multiple birth, having attended ANC, and the mother's age.

18. This issue of difficult birth selectivity is yet more severe for births at health facilities, making it very difficult to measure the beneficial impact of such facilities by this sort of statistical analysis.

19. This is the single difference in 1999 ($7 = 38-31$); that is, just after training was stopped. The double-difference estimate from the bivariate analysis is 30 deaths averted, as the mortality rate was much higher for births attended by trained TBAs than untrained in 1993.

20. Costs are estimated from USAID implemented TBA training programs in Afghanistan, Indonesia and Madagascar (which give estimates in the range US\$350-500). Data are scarce on the number of deliveries per year, though it appears to vary widely.

21. These coefficients are averaged across estimates from different specifications.

22. This calculation omits the supply-side costs associated with higher enrollments.

23. The cold chain is a system of conservation and distribution of vaccines to maintain them in cold temperatures, between 32° and 46.4°F, which is necessary to guarantee their effectiveness.

24. The study found that income in households with electricity was 65 percent higher than that in

households of non-electrified villages, and 126 percent higher than households without electricity in electrified villages. Rather than tackle the endogeneity problem econometrically, the study breaks down income by source to identify those that may be attributed to electrification. On this basis it is concluded that 16.5 percent of the income of electrified households is attributable to electrification (Barkat and others 2002).

25. Assuming a savings propensity of 0.1 and using the wealth-expenditure elasticity of 0.8 estimated using HIES data (Annex C).

26. The figures of US\$500 and US\$1,000 per connection were provided by the task manager for the Bank's rural electrification projects. Cost-benefit calculations in the ICR for RE III present cost figures for four new rural electricity cooperatives, giving a discounted cost of Tk 1,686 million cost for 80,210 new residential connections. These figures give a cost of US\$470 per residential connection during the 1990s (see Annex I).

27. The number of deaths averted in a household depends on household composition. If all children are already over five, no deaths are averted. However, if the mother is just entering her childbearing years, on average just over three children will be born who are less exposed to risk of premature mortality on account of electrification. An analysis of household structure suggests that on average 1.2 children per household will benefit from the reduced risk due to electrification. Since all figures are indicative of broad orders of magnitude, a figure of one child per household is used in the cost-effectiveness calculations. However, unlike immunization, which benefits just that child (possibly others through the herd immunization effect discussed above), and schooling, which benefits the children of a single woman, electrification will benefit subsequent generations, so that the cost calculations have an upward bias from this factor. But in comparison to the female schooling calculations, they have an overestimate, since the electrification figures include the income effect, which was excluded for schooling.

28. The effect of delayed marriage in reducing mortality introduces a "second-order effect" by which delaying marriage will indeed reduce fertility.

29. Of course this decline partly represents the fact that women subject to permanent methods by questionable, semi-coercive means in the early years of the program are leaving the group of women covered by the DHS.

Chapter 5

1. Thanks are due to the staff the World Bank Nutrition Hub for facilitating access to the BINP evaluation data set, to Save the Children for providing its data, and the staff of Helen Keller International Dhaka for providing the Nutritional Surveillance Project data.

2. Theory-based evaluation is propounded in Weiss (1998); see Carvalho and White (2004) for an application to development projects.

3. In some thanas BINP was implemented by the Government of Bangladesh, so that this last level of supervision was missing.

4. In addition to the adolescent girls' forum, there is a married couples' forum. However, during fieldwork, only wives were found to be present at such forums.

5. The BINP appraisal document makes some mention of other decisionmakers, but the main focus of both design and implementation has been mothers.

6. Calculated from endline data using responses "always" or "almost always" attend.

7. However, the same supplement was provided throughout supplementary feeding, which does not appear to be a good example regarding nutritional practices.

8. During fieldwork, differing criteria appear to have been applied to detecting growth faltering, which may in part reflect poor understanding by CNPs (found in interviews with 20 CNPs in two project thanas for the OED qualitative study, see box 5.1) or confusion arising from the transition to new criteria under NNP. For normal children, the official criteria appear to have been less than 600 gram gains in two months for those aged 6–12 months and 300 grams in two months for those aged 12–24 months. For children with WAZ < -2 SDs, these figures apply to a three-month period.

9. Type I error is the proportion of the target group not receiving the benefit, and Type II error the proportion of those receiving the benefit who are not eligible to do so.

10. This analysis requires information on pre-feeding nutritional status, which is not available in the BINP evaluation and Save the Children data sets. Save the Children also collected register data, but only for children in the feeding program, so Type I error cannot be calculated from these data. The Type II error calculated from the Save the Children register data is 15.4 percent, almost identical to that found in the Cambridge studies.

11. A number of factors probably lay behind poor targeting, one of which is certainly the difficulty CNPs have in interpreting the quite complex eligibility criteria. Impressions gathered in fieldwork suggest this to be a factor, though there are no numbers to present. The Save the Children data show that the majority of mothers do not understand the growth charts.

12. Data are not available on pre-pregnancy weight. The BMI used here is calculated from data for women not more than six months pregnant, so there will have been some pregnancy weight gain, thus providing an upward bias to the estimate of Type II error—but a downward bias on the estimate of Type I error.

13. Eating down was recommended medical practice in European countries for this reason until the 1940s. There is not unanimity regarding the accepted wisdom, with a recent Australian study finding that women delivering children with a higher birth weight are more likely to require a Caesarian section.

14. One striking impact of BINP appears to have been a much greater utilization of antenatal care (ANC) services, which is attributed to the one-stop service provided under BINP. In practice, this means that the CNP coordinated with health workers to provide services at the same time and place. Exposure to ANC under these circumstances implies exposure to nutritional messages.

15. That is, women could be in receipt of supplementary feeding without acquiring nutritional information. This finding is not a good one for BINP, since supplementary feeding was intended mainly as a tool for demonstrating feeding practices, not as a feeding program. However, it was already shown that one-third of mothers whose children were enrolled in supplementary feeding did not discuss nutrition with the CNP.

16. Participation in supplementary feeding was not found to have a significant impact. This impact may have been muted by leakage and substitution.

17. The assumptions underlying this calculation are laid out in Annex G. A 25 percent administrative overhead was allowed.

Chapter 6

1. While there has been a health facility survey in Bangladesh during HPSP, the analytical possibilities from a survey are far greater when the facility survey can be merged with a household survey.

Annex B

1. In the multiple regression for HAZ primary education in fact has the wrong sign, though it is not significant.

Annex C

1. The data used in calculations are three DHS rounds, collected in 1993/94, 1996/97, and 1999/2000. Mortality rates are calculated using the synthetic cohort probabilities method (Rutstein 1984). Data from the survey rounds are pooled; thus, the rate for 1985-89 includes information from each survey round, while that for 1995-99 includes data on under-fives from the 1999/2000 round and under-twos from the 1996/97 round.

2. DHS collects information on household durable goods ownership and housing quality, which can be used to construct a wealth index. DHS does not collect data on income or expenditure.

3. Information on differences in health outcomes and parental behavior within the same family can be used to get around the endogeneity problem, though this method can only be applied to families for which there are two or more live births (Rosenzweig and Schultz 1983).

4. The positive correlation between preceding birth order and mortality could be spurious due to prematurity. Premature births have higher mortality risk and, by definition, a lower birth interval, so the effect of a regression coefficient on preceding birth interval could simply be picking up the relationship between prematurity and perinatal mortality (Miller 1989). Miller and others (1992) find that controlling for gestation period reduces the estimated effects of low birth interval by as much as one-third.

5. Subsequent birth interval is also likely to affect mortality risk of children through a number of channels, including premature weaning and reduced quantity and quality of parental resources devoted to the older child in infancy (e.g., because the mother is physically tired and therefore less able to provide adequate care for sick children), and in childhood due to competition among siblings for limited familial resources, if the family believes that he or she has already passed the critical stage for survival (Muhuri and Menken 1997). However, the relation between subsequent birth interval and mortality is likely to be highly endogenous because birth interval is likely to be shorter if the older child has died; this effect operates both behaviorally, where parents choose to replace the lost child, and biologically, as death of the previous infant or young child interrupts breastfeed-

ing causing ovulation to return early (Palloni and Tienda 1986; Koenig and others 1990).

6. It is possible to test between the competing explanations of why preceding birth interval affects mortality by interacting preceding birth interval with mortality status of the previous child, with a significant negative effect on the interaction term providing support for the resource competition theory (Bairagi, Islam, and Barua 1999).

7. Muhuri and Preston (1991) argue that mortality risk is likely to be greater for both (younger) boys and girls the greater the number of (older) siblings of the same sex there are, since children of the same sex are better substitutes for one another in the family economy than children of the opposite sex (though the effect for girls will be higher).

8. Regarding the latter, it is important to use sibling sex composition at birth of the index child, as opposed to sex composition of siblings at interview date, since the latter will be endogenous to mortality.

9. Household size and age composition are potentially important determinants of child health status, both indirectly as a measure of the ratio of hands to mouths available, but also directly by indicating the quantity and quality of care a given child may be able to receive (Chamarbagwala and others 2004). However, most studies, including this one, are unable to include directly indicators of household size to explain fertility and mortality differentials because household composition is endogenous to fertility decisions and mortality.

10. Note that some variables indicate different underlying cause for different age groups. Males have higher biological mortality risk than females at all ages, which is reflected in higher observed mortality rates among male neonates; higher observed mortality rates among girl children in South Asia, on the other hand, is entirely behavioral, due to son preference. Similarly, short birth interval and higher birth order are known to increase mortality risk in (neonatal) infancy, possibly for biological reasons such as physical depletion of the mother, but for older children effects will be entirely behavioral, and probably modified by sex of the child.

11. The advantage of the Cox semi-parametric estimator is that, in comparison to fully parametric proportional hazards models such as the Weibull, assumptions do not have to be made about the baseline hazard.

12. For immunization of children, the variable used in this paper is the geographical mean (by cluster and gender) of children immunized. This is because immunization data are only collected on children who

are alive at the time of interview, so immunization status cannot be used. The same procedure is used for Vitamin A supplementation, which is also only available for live children.

13. Chittagong includes Sylhet, which became a separate Division in 1996/97.

14. Due to extreme endogeneity of breastfeeding with respect to child health status—e.g., low birth weight or sick children may be unable to breastfeed—variables indicating breastfeeding are omitted from the explanatory variable set in multivariate analysis.

15. The non-linear variable x enters the probit equation in the form $\Phi(b_1x + b_2x^2 + \dots)$ where \dots indicates the remaining arguments of the equation. Taking the partial differential with respect to x , setting equal to zero and solving for x gives the maximum/minimum at $x = -b_1/(2b_2)$.

16. The effect of education is stronger when mothers have completed primary education than when they have not (results not reported).

17. It was also possible to test the impact of education via (self-reported) literacy, though coefficients on the literacy variable were insignificant in all regressions for all three age periods.

18. These results are from the multivariate analysis. As shown in table C.1, there is higher mortality in lower quintiles than higher, though the differential is less marked for neo-nates than for children at older ages.

19. Rural is insignificant when interactions between month of birth and rural are dropped (results not shown).

20. A variable indicating that the household resided in a Bangladesh Integrated Nutrition Project (BINP) area was included in a antenatal care probit over the rural 1999/2000 survey round sub-sample (results not reported)—coefficients were positive but insignificant (p -value=0.20).

21. Wald tests can not reject the null hypothesis of insignificance at p -values 0.20 and 0.14, respectively.

22. There is no statistical evidence for endogeneity between quality antenatal care and neonatal mortality, indicated by the insignificance of RHO 2,1 (a Wald test for insignificance does not reject the null hypothesis at p -value 0.98).

23. Interactive terms between division of residence and gender are not significant determinants of post-natal mortality (results not reported).

24. The agency index is the simple sum of indicators of whether the woman has final say on own and children's health care, household purchases for large items and daily needs, what should be cooked each day and visits to family, friends or relatives.

25. Another possibility is that wealthier households were better able to benefit from the BINP. However, an interactive term between BINP and wealth is insignificant, while the interaction between BINP and mother's education remains significant. The same result is found for child mortality.

26. As well as the coefficients, hazard rates are reported, indicating the effect of a unit change in the explanatory variable on the risk of death in childhood.

27. A test for proportional hazards based on Schoenfeld residuals indicates the null hypothesis (that the proportional hazards assumption is not violated) is not rejected for each explanatory variable separately, while the joint test for all variables does not reject the null at p -value=0.62.

28. For neonatal mortality regressions, period 2 (mid-late-1990s) contains 88 more observations than period 1 (early-mid-90s), therefore, in order to match observations from each period, 88 observations are omitted, at random, from the period 2 sub-sample. For postnatal mortality regressions, period 2 contains 1,009 fewer observations than period 1; these observations for period 1 are deleted at random.

29. Appendix C.3 presents decompositions of changes in socioeconomic inequality in mortality by changes in inequality in the determinants of mortality, based on the regression results. The most important variables explaining reduction in socioeconomic inequality in mortality among neonates are birth order and electricity. For children, the reduction in wealth inequality contributed the most of any single variable to reducing inequality in mortality, followed by the improved distribution of education among women.

Appendix C.2

1. Bollen and others (1999, p. 4) state that the term *socioeconomic status* "appears to have become commonplace only after the U.S. Census Bureau published a report in 1964 using a composite SES index for comparison of different groups."

2. There is also the problem of lumpy consumption items—expenditure data collected in household surveys will be sensitive to large non-recurring consumption items such as healthcare bills, which would tend to artificially inflate the household's estimated welfare, particularly if during the preceding or subsequent period the household saved for the item by reducing expenditure. While measures are usually taken when constructing household expenditure aggregates to exclude some non-recurring items, it has often been argued that because of these problems household expenditure data are satisfactory for calculating poverty levels, as these differences get ironed

out, but not good at identifying poor households (Grosh and Glewwe 2000). Under these circumstances, wealth may be a better indicator of differences in current income across households, and assets are a better predictor of this. The problem with wealth is that the data collected on assets are usually limited (e.g., do not include financial savings), and in poorer countries, all the better-off have most or all of the items and many of the poorest have none, making it difficult to distinguish between them.

3. Young Lives (2002) gives a formula for scaling a variable X between 0 and 1, where welfare increases with X :

$$Xi^* = (Xi - Xmin) / (Xmax - Xmin)$$

where Xi^* denotes the scaled observation of individual i in variable X , and min and max refer to the respective minimum and maximum of variable X . To ensure spatial and temporal comparability, the minima and maxima may have to be pre-set rather than determined by the data.

4. Asset scores are calculated according to the formula $g_{ik} = f_k(a_i - \bar{a})/s$, where g_{ik} represents the asset score of asset k for household i , f_k is the "raw" asset score generated by the principal components analysis, and $(a_i - \bar{a})/s$ is the standardized value of asset variable a_i using mean \bar{a} and standard deviation s (Gwatkin and others 2000, Annex B).

5. Young lives is collecting panel data with specific interest in children's well-being in Ethiopia, India, Peru, and Vietnam; see www.younglives.org.uk

6. The sub-categories may also misclassify. For example, bicycles and motor vehicles may serve productive purposes; livestock and land represent livelihood and production activities, but may also be used as a form of savings to insure against adverse circumstances, or as collateral against credit, and therefore fulfill a similar role to financial assets. Indeed, it may be instructive to classify assets according to function, since items are held for different purposes—consumption, production, and savings, with some items simultaneously fulfilling different functions. In this context, the term "assets" seems more appropriate for production and savings, and it is for this reason that Young Lives (2002) distinguishes a "wealth index" (which includes only household items) from an "asset index" (which includes items used in production and/or as a store of value), where both indices can include consumer durables.

7. Physical contact with certain types of fuel can be hazardous to health, as can environmental pollution generated by burning fuel in unventilated areas.

8. In principle, where the locational nature of assets varies by country or within country by region,

cluster-level data could be used to determine if a household potentially has access to a service (water, electricity, etc.), and the asset included in the index for those clusters with access and excluded for those without, with suitable rescaling across assets to ensure comparability across all clusters.

9. In order for different SES indicators to produce different estimates of socioeconomic inequality in another variable, the indicators must produce different rankings of households and these rank differences must be correlated with the welfare outcome (Wagstaff and Watanabe 2002).

10. Inherent subjectivity associated with choosing assets to be included would also tend to favor inclusion of all available assets (Gwatkin and others 2000), though differentiation according to asset category—e.g., household assets, community assets—can limit the degree of subjectivity.

11. In PCA the structure of the model is determined by the *variance* of the data; in FA the structure is determined by the *covariances* (or correlations) between variables.

12. PCA has the advantage that, exclusion of higher order principal components does not theoretically result in mis-specification (omitted variables bias) in multivariate analysis, since the principal components are constructed as orthogonal to (i.e., not correlated with) one another; bias may still arise through correlation of the omitted principal component with other explanatory variables (Filmer and Pritchett 1999, p. 118, fn. 8).

13. Wagstaff (2000) notes that deciles could be used in principle, though sample size is often too small to make this classification statistically meaningful.

14. Given whether the shock is covariate or idiosyncratic (a covariate shock inducing many households simultaneously to try to sell their assets would depress prices).

15. It could be argued that wall and roof type have direct implications for health and nutrition—though less substantially than flooring—by increasing exposure to infection. Nevertheless, given the limited alternative assets available for the index, these variables were included in the index.

16. The weighting scheme that would maximize variance determines each score according to the variance of the asset, equal to $P \cdot (1-P)$, where P is the proportion owning the asset; this accords greatest weight to the assets with largest variation in ownership. This scheme was deemed unsuitable for the asset index, since it produced scores that appeared unreasonable; for example, ownership of a table and clock each received scores almost twice as large as ownership of a television.

17. In a recent field visit, it was observed that urban houses (and rural houses near main roads) were mainly constructed of corrugated metal sheets, while the main building material in rural areas were mud and thatch (Howard White, personal communication). In addition, some mud buildings were observed to be actually brick with mud daubing (which is used for traditional decorative patterns)—whereas usually enumerators fill in construction material from own observation so these houses may be misclassified.

18. Unweighted means of, or proportions of households owning, variables are as follows: radio (0.303); TV (0.137); bicycle (0.189); almirah (wardrobe) 0.265; table (0.570); clock (0.471); cot (0.787).

19. The poverty index calculates poverty P using the formula:

$$P(\alpha) = 1/n \cdot \sum_i ((z - A_i)/z)^\alpha$$

where z represents the poverty line. The most common used values of α are 0 (the poverty headcount index), 1 (the poverty gap index) and 2 (the squared poverty gap index).

Appendix C.3

1. The alternative decomposition is $\Delta MR = \Delta MR^p \cdot p_1 + \Delta MR^n \cdot (1-p_1) + 2Dp \cdot (MR_2^p - MR_2^n)$. When applied to the data, this method accords more weight to reductions in mortality among the poor.

2. The decomposition is of $p_i \cdot MR_i^p + (1-p_i) \cdot MR_i^n$ as opposed to the total mortality rate since the synthetic cohort method used to calculate mortality probabilities takes a different sample for each group.

3. It is supportive, though not conclusive, as there may be economic improvement for those who nonetheless remain poor. But expenditure data from household surveys show rising inequality, with much higher average growth for the top 50 percent than the bottom 50 percent (World Bank 2002).

4. The t-statistic is calculated by:

$$t = (C_2 - C_1) / [\text{Var}(C_2) + \text{Var}(C_1)]$$

where C_1 and C_2 represent the estimate of the concentration index in 1980-84 and 1995-99 and Var is the variance operator.

Annex D

1. Anthropometry measures the body's quantitative change in response to changes in the balance between net energy requirements and net energy intake from food.

2. According to Chen and others (1980), in a longitudinal study, weight for age was found to be the strongest predictor of mortality, followed by height

for age and finally weight for height. Of course, short children may well be more at risk where height indicates low birth weight or a history of sickness.

3. The reference population is the NCHS/WHO (1978) reference.

4. Z scores are preferable to other measures, such as the percentage of the reference median, because they account for distributional variation specific to different ages or heights in the reference population. Children with z scores between -2 and -3 are chronically malnourished and those below -3 z scores severely malnourished.

5. Using multivariate analysis of panel data, Bairagi (1986) finds evidence that nutrition discrimination against girls increases following famine.

6. Controlling for education and health knowledge in the regression equation should proxy some component of this heterogeneity.

7. The wealth index is calculated as the weighted sum of household durable goods ownership and quality of housing (wall and roof) building materials. This excludes factors that are expected to have “direct” effects on child health and nutrition, such as drinking water source, sanitation, and dwelling flooring, in order that these effects can be estimated separately.

8. Income, maternal education and access to clean water and sanitation may potentially be substitutes in the production of child health. This was tested for in regression analysis using interactive regressors, and no significant impact was found.

9. For Peru, Alderman and Behrman (2003) estimate positive community externalities in female education, water, and sanitation on top of the observed effects for each child by including geographical area non-self means of these variables in the set of explanatory variables. The meta-analysis found community-level access to clean water to be marginally significantly positive, though sanitation was insignificant (Charmarbagwala and others 2004).

10. Their data are the 1981-92 Nutrition Survey of Rural Bangladesh, which is a nationally representative sample of 385 households.

11. The sign of the estimated coefficient on the inverse Mill's ratio reflects the correlation between error terms in selection and regression equations, providing statistical evidence for non-random selection (Greene 2000). In the case of nutrition, the coefficient is expected to be positive, indicating that the unobserved characteristics determining survival, such as preferences regarding child care, broadly defined, are also positively correlated with those influencing nutrition.

12. To see the intuition behind this, note that a model estimating $P(Y_i=1)=\Phi(Z_i)$ generates inverse Mill's ratio $\phi(Z_i)/\Phi(Z_i)$. It follows that the model estimating $P(Y_i=0)=1-\Phi(Z_i)$ leads to Mill's ratio $-\phi(Z_i)/(1-\Phi(Z_i))$ due to symmetry of the standard normal distribution.

13. Comparable trends are found when tabulating incidence of stunting, wasting, and underweight by these variables.

14. Modeling anthropometry at ages below 6 months is complicated by problems in physically measuring young babies with accuracy; also the WHO/NCHS reference population was bottle-fed, while WHO now recommends breastfeeding during this period.

15. This is a crude estimate of the BINP effect; a more comprehensive analysis is provided in Annex G.

16. The treatment effects regression is discussed in Greene (2000).

17. No impact of contraceptive knowledge and literacy on weight for age could be found when these variables were included in the regressions.

Annex E

1. The following calculations were made including only “usual residents” from the household roster. Those people identified as not related to the household head were not included since, as mentioned above, these people are most likely live-in servants.

2. As discussed below, DHS 1999/2000 contains data on if a woman makes the decision on what to cook each day by herself. Where this is the case for more than one woman, then this fact may indicate a joint household.

3. For example, there are 70 women in the dataset aged over 35 and living with a parent, half of these were formerly married—and the other half are currently married, there are no never-married women in this group (there are only 19 never-married women from the 6,209 aged over 35 in the whole dataset).

4. The age of the oldest son is given as 50. The ages of the adults—reported as 30, 35, 40, 45, 50, 60, and 80—clearly suffer from what is called “heaping,” i.e., rounding by respondents to the nearest 5 or 10. Nonetheless, it seems likely that the oldest son at least is the product of a previous marriage.

5. See White (2001) for more discussion of this contradiction in findings.

6. There is a strict monotonic inverse relationship between asset quintile and the proportion of women in either paid or unpaid work. Twenty-five percent of women in the poorest quintile engage in paid work,

and 4.6 in unpaid. For the top quintile these figures are 14.8 and 0.9 percent, respectively.

7. The age variables are more significant than one measuring the age gap between husband and wife, which is only significant when age is not included.

8. Analysis of other data shows that respondents saying they can read the paper often prove to have poor literacy, which will undermine the power of this variable.

9. Dummies were initially included for all regions with Dhaka as the reference region. Those with insignificant coefficients were dropped in the results reported here.

Annex F

1. The fitted line is $FERT = 10.8 - 0.33 \text{ AGEM}$, $R^2 = 0.24$, $n = 126$.

2. The median age at marriage has shown only a small change. But this statistic is a problematic measure as it is necessarily based on ever-married women, excluding from the calculation women not yet married. It is also insensitive to movements within the bottom half of the distribution.

3. Alternatively, OLS can be used with a spline function, which gives $\text{age_fb} = 5.24 + 0.83 \text{ age_marr} + 10.2 \text{ dum} - 0.82 \text{ sldum}$, where $\text{dum} = 1$ for women married at 13 or less, and sldum is the slope dummy for that variable for these women.

4. Of course, any delay will have a temporary tempo effect on the TFR.

5. Until some generations ago, bridewealth was practiced in Bangladesh, but has now been commonly replaced by dowry which was previously only found among Hindus. The former practice is consistent with Islam whereas the latter is not.

Appendix F.1

1. The GRR is the average number of daughters that would be born alive to a woman (or group of women) during her lifetime based on currently prevailing age if she passed through her childbearing years conforming to the age-specific fertility rates.

2. $GRR = a + b \text{ CRW}$, where a and b are given in Rele (1976). The $TFR = GRR \times 2.05$ (assumed sex ratio at birth).

3. Suggested weights to arrive at a final estimates are 0.7 for GRR obtained from CWR2 and 0.3 for GRR obtained from CWR1.

4. Results of the 2001 population census are provisional ones calculated based on 5 percent sample schedule. Final results are yet to be published.

Annex G

1. This study focuses on protein-energy malnutrition, as do other studies of BINP. Data have not been collected to analyze micronutrient components.

2. The CNC is a facility provided by the community, it may sometimes be the health clinic, but is more usually at the home of one of the Community Nutrition Committee members.

3. It is also more common for a woman to be living with their mother-in-law in these thanas, being so for 15 and 12 percent of respondents in Rajnagar and Shasthi respectively, compared to 9 percent on average.

4. The BINP dataset does not contain data on distances to health facilities. Using data of the Bangladesh Household Income and Expenditure Survey (HIES) of 2000-01, it was found that the time distance to the health provider sought for treatment was substantially longer for household not having access to drinking water (i.e., accessing water from rivers, ponds, rain, etc.). The difference was statistically significant using a t-test.

5. A similar figure for Type II error was reported in the Cambridge study.

6. Save the Children data show that most mothers cannot interpret growth charts. Proponents of growth monitoring question whether this is necessary for the approach to work, though Christiaensen and Alderman (2004) have found a mother's understanding of her child's growth performance to be a significant determinant of child nutrition in Ethiopia. The World Bank Nutrition Toolkit No. 4, "Promoting the Growth of Children: What Works" (Griffith, Dickin and Favin 1996), states that mothers helping weighing the child and interpreting the growth pattern are part of a good growth monitoring program (p. 87), as this increases the mother's understanding of child growth and of its importance (p.35).

7. Equal to one if any household member is involved in an association, savings group, or income-generating activity.

8. That is, women could be in receipt of supplementary feeding without acquiring nutritional information. This finding is not a good one for BINP, since supplementary feeding was intended mainly as a tool for demonstrating feeding practices, not as a feeding program. However, it was already shown that one-third of mothers whose children were enrolled in supplementary feeding did not discuss nutrition with the CNP.

9. This variable is obtained using the information on the date of interview and the age of the mother's child. A woman is considered as pregnant during the working season if the last six months of pregnancy were mostly in the months between November and April.

10. A longer version of this section is available on the study website.

11. All malnutrition prevalence rates have been calculated using the old NCHS reference population growth charts. The National Nutrition Project (NNP) has adopted the new (2000) NCHS growth charts in 2004. The introduction of the new charts will have predictable effects on the calculation of malnutrition rates.

12. There are little more than 150 control observations in the baseline survey (see last column of table G.17 for actual figures).

13. The fact that knowledge has not risen in control thanas suggest that contamination is not a serious problem.

14. The text of the IMED report claims a larger impact if considering only the weight for height proportion of children aged 0-23 months.

15. One possibility, which cannot be explored with the data to hand, is that improved nutrition was expended in extra physical activity rather than weight gain.

16. It is not possible to discern from the Save the Children questionnaire the children who participated in GMP from those who did not. In addition, there is little information that can be used to identify participants in a model of participation. Finally, the absence of data on the date of the interview is also a concern, because of the sensitivity of nutritional indicators to seasonal effects.

17. After selecting NSP data of the same months in which the BINP data were collected, some discrepancies in the temporal distribution of the observations BINP and NSP data still remains. In order to correct this source of seasonal variation in nutritional indicators, the data were seasonally adjusted.

18. It is questionable whether the equation can be estimated using the HKI data since households outside the project area could not participate even if they wanted to. An alternative procedure is to estimate the participation equation using only the project sample and then apply those coefficients to the NSP data to calculate the scores. This approach was also used, with the final results not being much different from those shown here.

19. Participation here refers to participation in the growth monitoring sessions. An attempt was made to construct a control for children participating in supplementary feeding. Data on the eligibility criteria are not available for the control, so that proxies had to be used, but created a poor control. Negligible project effect was found.

20. Project effects are estimated by running ordered logit models of participation, where the dependent variable takes three values: 0 for non-participation, 1 for

irregular participation, and 2 for continuous participation. The predicted probability are used to weight the observations. Outcomes of children whose mother made the choice 2 are weighted by the inverse of the predicted probability of that choice. The project effect is calculated taking the difference between this value and the mean outcomes of the children who did not participate.

21. Tables G.21 and G.22, and the charts in figure G.3 are based on matched observations of BINP evaluation sample and NSP data, performed with the one-to-one matching method.

22. The category of normally nourished is not shown in table G.24 as they constitute less than 1 percent of program participants.

23. These are the quantitative targets defined in the project document (World Bank 1995, p. 15). It is not clear how the latter can be measured, since the same women will not be pregnant at midterm and endline as were at baseline. Karim and others (2003) rephrased the target as "improvement of weight gain to 7.0 kg. in at least 50% of pregnant women."

24. See in particular the reviews by Chatterjee (1991) and Nag (1994).

25. Estimates by NIPORT (2002).

26. It is not clear, however whether the absence of an effect of food supplementation program on maternal mortality risk is the result of the absence of an effect of larger body size on delivery, or of the absence of an effect of food supplementation on body size.

27. In order to measure intrauterine growth it is necessary to adjust birth weight by gestational age at birth in order to isolate the component influenced by nutrition. Several reference populations are available. The one used here are a Canadian reference population (Kramer 2001) for birth weight for gestational age. BINP data report only the month of birth as the 8th, 9th, or 10th, while reference weights by gestational age are normally reported by week. The recommended weight at the 40th week of gestation is used as a reference for the 9th month reported weight; the 39th week for the 8th month and the 41st week for the 10th month, assuming that the distribution of births in the 8th and 10th months tend to concentrate around the 40th week. This ratio is not very precise, but it is better to adjust for sex, and partially for gestational age, than not adjusting at all.

28. The turning point is obtained as $-b_1/(2*b_2)$, where b_1 and b_2 are the regression coefficients of mother's age and mother's age squared respectively.

29. See the review of studies on the effect of supplementation project on low birth weight in Waterlow (1992) and Institute of Medicine (1990). These review

report the following changes in birth weight: Guatemala (+100), Colombia (+ 180, but only for thin women), Gambia (+ 120, but only in hungry season), Chile (+ 60), Indonesia (no effect). No effect on birth weight was observed in similar projects carried out in Canada, Scotland, and the United States.

30. Given the time required to identify pregnant women and enroll them in BINP activities, pregnant women will be covered by project activities for about six months rather than the full nine months of pregnancy. In any case, weight gain is relatively small in the first trimester (see figure G.5).

31. Percent changes in malnutrition are calculated using a probit function in the following way. If Z is the average z -score before the project and dZ is the change in the average z -score of table 2, the percent reduction in malnutrition is obtained as $\Phi(-2 - Z) - \Phi(-2 - Z + dZ)$. Malnutrition rates are calculated using -2 z -score as cutoff point (this includes mild and severe malnutrition).

32. Data on calories and rice price are obtained from the IFPRI data of 1998.

33. A large fraction (58 percent) of the nutrition component of the project is invested on food purchase and preparation. It is estimated that 85 percent of this food is consumed by mothers enrolled in the supplementary feeding program. The costs of reducing malnutrition and of saving one life reported in the second column are calculated after subtracting costs that are directed to mothers.

34. Use of current government workers or local-level voluntary workers was ruled out of the BINP design because of the workload. But a large part of this workload relates to growth monitoring rather than BCC.

Annex H

1. The ratio of the total budget for FSSAP to the stipend component is 1.20, giving an overhead of 20 percent.

Annex I

1. Rice accounted for almost three-quarters of total crop production in the late 1990s (Mian and others 2001, p. 39).

2. Net cropped area refers to the area actually cropped, whereas effective gross cropped area refers to the total yearly area cropped counting areas under double and triple cropping two and three times respectively.

3. Transplanted, broadcast, and deepwater rice are different varieties of rice, which exist within each rice sort. Each of these contains various subgroups.

4. The study looked at weight for age (WAZ); height for age (HAZ); arm circumference (AC) for age; and triceps skinfold thickness (TSFT) for age. The latter two had better indicators of seasonal variations in nutrition given their independence from height.

Annex J

1. *Books, Buildings and Learning Outcomes: an impact evaluation of World Bank support to basic education in Ghana*, OED, January 2004.

2. *2002 Annual Review of Development Effectiveness - Achieving Development Outcomes: The Millennium Challenge*, OED, report no. 25159.

3. Determinants are the variables entering the multiple regression analysis, such as mother's education, and drivers the policies and programs which drive changes in these determinants (e.g. policy on school fees, and female scholarships).

4. UNICEF (1990) *Strategy for Improved Nutrition of Children and Women in Developing Countries*. New York: UNICEF.

5. W. Henry Mosley and Lincoln C. Chen (1984) "An Analytical Framework for the Study of Child Survival in Developing Countries." *Population Development Review* Vol. 10 Supplement: Child Survival: Strategies for Research, 25-45.

6. For example, J.M. Currie "Child Health in Developing Countries" in Culyer and Newhouse (eds), *Handbook of Health Economics* Amsterdam, Elsevier Science BV, 2000.

7. Public expenditure reviews are available from 1990, 1996, 1997 and 2003.

REFERENCES

- Agarwal, B. 1990. "Gender Relations and Food Security: Coping with Seasonality, Drought, and Famine in South Asia." PEW/Cornell Lecture Series on Food and Nutrition Policy, November 13, Washington, DC: CFNPP.
- Alauddin, M., and M. Hossain. 2001 *Environment and Agriculture in a Developing Economy: Problems and Prospects for Bangladesh*. Northampton, MA: Edward Elgar.
- Alauddin, M., and C. A. Tisdell. 1998. *The Environment and Economic Development in South Asia: An Overview Concentrating on Bangladesh*. London: Macmillan.
- Alderman, H., and J. R. Behrman. 2003. "Estimated Economic Benefits of Reducing LBW in Low-Income Countries." University of Pennsylvania, Philadelphia, for Human Development Network/ Nutrition, World Bank, Washington, DC. [Photocopy.]
- Alderman, H., and M. Garcia. 1994. "Food Security and Health Security: Explaining the Levels of Nutritional Status in Pakistan." *Economic Development and Cultural Change* 42 (3): 485–507.
- Alderman, H., J. Hentschel, and R. Sabates. 2003. "With the Help of One's Neighbors: Externalities in the Production of Nutrition in Peru." *Social Science and Medicine* 56 (10): 2019–31.
- Ateng, B. 1998. "Comparative Advantage and Crop Diversification." In *Bangladesh Agriculture in the 21st Century*, ed. R. Faruquee. Dhaka: The University Press.
- Baffes, J., and M. Gautam. 1996. "Is Growth in Bangladesh's Rice Production Sustainable?" World Bank Policy Research Working Paper 1666, Washington, DC.
- Bairagi, R. 1996. "Is Son Preference an Obstacle to Fertility Transition in Bangladesh?" In IUSSP Committee on Fertility and Family Planning, Population Council Islamabad, Rockefeller Foundation, and UNFPA Pakistan Office, *Comparative Perspectives on Fertility Transition in South Asia*. Liege: IUSSP.
- . 1986. "Food Crisis, Nutrition, and Female Children in Rural Bangladesh." *Population and Development Review* 12 (2): 307–15.
- . 1980. "Is Income the Only Constraint on Child Nutrition in Rural Bangladesh?" *Bulletin of the World Health Organization* 58 (5): 767–72.
- Bairagi, R., M. Mazharul Islam, and M. K Barua. 1996. "Contraceptive Failure: Levels, Trends and Determinants in Matlab, Bangladesh." Dhaka: Population Studies Center, ICDDR'B. [Photocopy.]
- Bairagi, R., S. Chandra Sutradhar, and N. Alam. 1999. "Levels, Trends, and Determinants of Child Mortality in Matlab, Bangladesh, 1966–94." *Asia-Pacific Population Journal* 14 (2): 51–68.
- Bangladesh Bureau of Statistics. 2000. *Statistical Pocketbook of Bangladesh*. BBS, Ministry of Planning, Bangladesh.
- Bangladesh Government. 2003. "A National Strategy for Economic Growth, Poverty Reduction, and Social Development." Economic Relations Division, Ministry of Finance. Dhaka.
- . 2002. "Interim Poverty Reduction Strategy Paper." Dhaka.
- Bangladesh Government/International Monetary Fund. 2003. "Bangladesh: Interim Poverty Reduction Strategy Paper." Washington, DC.
- Barkat, Abul, S.H. Khan, M. Rahman, S. Zaman, A. Podder, S. Halim, N.N. Ratna, M. Majid, A.K.M. Maksud, A. Karim, and S. Islam. 2002. *Economic and Social Impact Evaluation Study of the Rural Electrification Program in Bangladesh*. Human Development Research Centre and NRECA International Ltd.
- Baulch, Bob, and John Hoddinott, eds. 2000. "Economic Mobility and Poverty Dynamics in Developing Countries." *Journal of Development Studies* 36.
- Beaton, G. H., and H. Ghassemi. 1982. "Supplementary Feeding Programs for Young Children in Developing Countries." *American Journal of Clinical Nutrition*: 863–916.
- Becker, S., and S. Weng. 1998. "Seasonal Patterns of Deaths in Matlab, Bangladesh." *International Journal of Epidemiology* 1999 (27): 814–23.
- Begum, Sharifa. 2003. "Abortion in Rural Bangladesh: Nature, Causes, Problems, and Practices." BIDS Research Report 174. Dhaka.
- . 1990. "Population Birth, Death, and Growth Rates in Bangladesh: Census Estimates." *The Bangladesh Development Studies*, Vol.18, No.2. Dhaka.

- . 1976. "An Estimation of Birth and Death Rate in Bangladesh, 1951–60." Bombay, IIPS. [Photocopy.]
- Berg, A. 1983. *Nutrition Review*. Washington, DC.: Population and Human Resources Department, World Bank.
- Bhuiya, A., and M. Chowdhury. 1997. "The Effect of Divorce on Child Survival in a Rural Area of Bangladesh." *Population Studies* 51: 57–61.
- Bhuiya, A., and K. Streatfield. 1991. "Mothers' Education and Survival of Female Children in a Rural Area of Bangladesh." *Population Studies* 45: 253–64.
- Blanc, Ann. 1990. *The Relationship between Contraceptive Prevalence and Total Fertility in DHS Countries*. Columbia, MD: Demographic and Health Surveys Project.
- Bloem, M., R. Moench-Pfanner, and D. Panagides, eds. 2003. *Health and Nutritional Surveillance for Development*. Singapore: Helen Keller International.
- Bollen, Kenneth A., Jennifer L. Glanville, and Guy Stecklov. 2002. "Economic Status Proxies in Studies of Fertility in Developing Countries: Does the Measure Matter?" *Population Studies* 56: 81–96.
- . 1999. "Socioeconomic Status and Class in Studies of Fertility and Health in Developing Countries." University of North Carolina at Chapel Hill.
- Bongaart, John. 1978. "A Framework for Analyzing the Proximate Determinants of Fertility." *Population and Development Review* 4(1).
- Bongaart, John, and R. G. Potter. 1983. *Fertility Behavior: Biological Analysis of the Proximate Determinants of Fertility*. New York: Academic.
- Briend, A., C. Dykewicz, K. Graven., R. Mazumder, B. Wojtyniak., and M. Bennish. 1986. "Usefulness of Nutritional Indices and Classifications in Predicting Death of Malnourished Children." *British Medical Journal* 293: 373–75.
- Brown, K. H., R. E. Black, and S. Becker. 1982. "Seasonal Changes in Nutritional Status and the Prevalence of Malnutrition in a Longitudinal Study of Young Children in Rural Bangladesh." *The American Journal of Clinical Nutrition* 36.
- Buhmann et al. 1988. "Equivalence Scales, Well-Being, Inequality, and Poverty." *Review of Income and Wealth* 34: 115–42. (Cited in Wagstaff 2000, p. 23.)
- Caldwell, J., B. Barkat-Khuda, I. Cladwell, I. Pieris, and P. Cladwell. 1999. "The Bangladesh Fertility Decline: An Interpretation." *Population and Development Review* 25(1): 67–84.
- Carvalho, S., and H. White. 2004. "Theory-Based Evaluation: The Case of Social Funds." *American Journal of Evaluation* 25(2): 141–60.
- Chamabagwala, Rubiana, Martin Ranger, Hugh Waddington, and Howard White. 2004. "The Determinants of Child Health and Nutrition: A Meta-Analysis." OED Working Paper, Washington, DC.
- Chatterjee, M. 1991. *Towards Better Health for Indian Women: The Dimensions, Determinants, and Consequences of Female "Illnesses" and "Death"*. Washington, DC.: World Bank.
- Chaudhury, Nazmul, and Jeffrey Hammer. 2003. "Ghost Doctors: Absenteeism in Bangladeshi Health Facilities." World Bank Working Paper 3065, Washington, DC.
- Chen, L., A. Chowdhury, and S. Huffman. 1980. "Anthropometric Assessment of Energy-Protein Malnutrition and Subsequent Risk of Mortality among Pre-School Aged Children." *American Journal of Clinical Nutrition* 33: 1836–45.
- Chen, L., E. Huq, and S. D'Souza. 1981. "Sex Bias in the Family Allocation of Food and Health Care in Rural Bangladesh." *Population and Development Review* 7: 55–70.
- Chowdhury, M., and R. Bairagi. 1990. "Son Preference and Fertility in Bangladesh." *Population and Development Review* 16: 4.
- Chowdhury, M. A. T., and H. Zulfikar. 2001. *An Agricultural Statistical Profile of Bangladesh, 1947–199*. Bogor, Indonesia: CGPRT Centre.
- Chowdhury, Zafrullah. 1996. *The Politics of Essential Drugs. The Makings of a Successful Health Strategy: Lessons from Bangladesh*. Dhaka: University Press Limited.
- Christiaensen, L., and H. Alderman. 2004. "Child Malnutrition in Ethiopia: Can Maternal Knowledge Augment the Role of Income?" *Economic Development and Cultural Change* 52 (2): 287–312.
- Cleland, John. 1994. *The Determinants of Reproductive Change in Bangladesh: Success in a Challenging Environment*. Washington, DC.: World Bank.
- Cleland, John, and Kim Streatfield. 1992. "The Demographic Transition: Bangladesh, UNICEF." Dhaka: Program Planning Unit. [Photocopy.]
- Cockcroft, Anne, Debbie Milne, and Neil Andersson. 2003. *Bangladesh: HPSP, 1998–2003. Third Service Delivery Survey 2003*. Dhaka: CIET Canada.
- Croll, E. 2001. *Endangered Daughters: Discrimination and Development in Asia*. London: Routledge.
- Davis K., and Judith Blake. 1956. "Social Structure and Fertility: An Analytical Framework." *Economic Development and Cultural Change* 4 (2).
- Del Ninno, C., P. A. Dorosh, L. C. Smith, and D. K. Roy. 2001. *The 1998 Floods in Bangladesh—Disaster Impacts, Household Coping Strategies, and Response*. Washington, DC.: International Food Policy Research Institute.
- Deolalikar, A. 2002. "Poverty and Child Malnutrition in Bangladesh." Bangladesh Poverty Assessment Background Paper 7.
- EM-DAT. 2004. OFDA/CRED International Disaster Database, www.em-dat.net. Brussels: Université Catholique de Louvain.

- Fairlie, R. 2003. "An Extension of the Blinder-Oaxaca Decomposition Technique to Logit and Probit Models." Economic Growth Center Discussion Paper 873. New Haven, CT: Yale University.
- Faruquee, R., ed. 1998. *Bangladesh Agriculture in the 21st Century*. Dhaka: The University Press Ltd.
- Faveau, V., A. Briend., J. Chakraborty, and A. Sarder. 1990. "The Contribution of Severe Malnutrition to Child Mortality in Rural Bangladesh: Implications for Targeting Nutritional Interventions." *Food and Nutrition Bulletin* 12: 215–22.
- Faveau V., M. A. Koenig, J. Chakraborty, and A. I. Chowdhury. 1988. "Causes of Maternal Mortality in Rural Bangladesh, 1976–85." *Bulletin of the World Health Organization* 66 (5): 643–51.
- Filmer, Deon, and Lant Pritchett. 1998. "Estimating Wealth Effects without Income or Expenditure Data—Or Tears: Educational Enrolment in India." World Bank Policy Research Working Paper 1994, Washington, DC
- Filmer, Deon, and Lant Pritchett. 1999a. "The Effect of Household Wealth on Educational Attainment: Evidence from 35 Countries." *Population and Development Review* 25 (1): 85–120.
- . 1999b. "The Impact of Public Spending on Health: Does Money Matter?" *Social Science and Medicine* 49 (10): 1309–23.
- Fine, P. 1993. "Herd Immunity: History, Theory, Practice." *Epidemiology Review* 15: 265–302.
- Foster, A., J. Greer, and E. Thorbecke. 1984. "A Class of Decomposable Poverty Measures." *Econometrica* 52: 761–66.
- Gage, A., E. Sommerfelt, and A. Piani. 1996. "Household Structure, Socioeconomic Level, and Child Health in Sub-Saharan Africa." Demographic and Health Surveys Analytical Reports 1.
- Gomez, F., R. Galvin, S. Frenk, J. Cravioto, R. Chavez, and J. Vasquez. 1956. "Mortality in Second and Third Degree Malnutrition." *Journal of Tropical Pediatrics* 2: 77–83.
- Goodburn, Elizabeth A., Mushtaque Chowdhury, Rukhsana Gazi, Tom Marshall, and Wendy Graham. 2000. "Training Traditional Birth Attendants in Clean Delivery Does not Prevent Postpartum Infection." *Health Policy and Planning* 15 (4): 394–99.
- Green, W. 1998. "Gender Economics Courses in Liberal Arts Colleges: Further Results." *Journal of Economic Education*.
- Greene, W. H. 2000. *Econometric Analysis*, 4th ed. New York: Prentice-Hall.
- Griffiths M., K. Dickin, and M. Favin. 1996. *Promoting the Growth of Children: What Works. Rationale and Guidance for Programs*. Washington, DC: World Bank.
- Grosh, M., and P. Glewwe. 2000. *Designing Household Survey Questionnaires for Developing Countries: Lessons from 15 Years of the Living Standards Measurement Study* 3 Vols. Washington, DC: World Bank.
- Gwatkin, Davidson R., Shea Rustein, Kierston Johnson, Rohini P. Pande, and Adam Wagstaff 2000. "Socio-Economic Differences in Health, Nutrition and Population in Bangladesh." HNP/Poverty Thematic Group of the World Bank, Washington, DC.
- Haddad, L. H. S. Alderman, L. Appleton, L. Song, and Y. Yohannes. 2003. "Reducing Malnutrition: How Far Can Income Growth Take Us?" *World Bank Economic Review* 17 (1).
- Haider, S. J., D. Hussain, I. Nayer, A. Maleque, Ashaduzzaman, and A.R. Khan. 2004. "Impact Evaluation of Bangladesh Integrated Nutrition Project." Dhaka: IMED and Read.
- Heckman, J., and B. Singer. 1984. "Econometric Duration Analysis." *Journal of Econometrics* 24 (1–2): 63–132.
- Horton, S. 1986. "Child Nutrition and Family Size in the Philippines." *Journal of Development Economics* 23: 161–16.
- Hossain, Shahadat, et al. 1991. "Fertility in Bangladesh: Levels and Trends." In *Proceedings of the Workshop on Dissemination of Current Statistics*, November 3–4, 1991, Dhaka.
- Hossain, M., D. Lewis, M. L. Bose, and A. Chowdhury. 2003. "Rice Research, Technological Progress, and Impacts on the Poor: The Bangladesh Case (Summary Report)." EPTD Discussion Paper 110. Washington, DC: IFPRI.
- Hossain, Z., and A.F. H. Choudhury. 1987. *Folk Dietary Practices and Ethnophysiology of Pregnant Women in Rural Bangladesh*. Michigan State University Working Paper 137, Ann Arbor.
- Houweling, Tanya, Anton Kunst, and Johan Mackenbach. 2003. "Measuring Health Inequality Among Children in Developing Countries: Does the Choice of the Indicator of Economic Status Matter?" *International Journal for Equity in Health* 2(1): 8–21.
- Howlader, A. A., and M. U. Bhuiyan. 1999. "Mothers' Health-Seeking Behaviour and Infant and Child Mortality in Bangladesh." *Asia-Pacific Population Journal* 14 (1): 59–75.
- Huq, Md. Nazmul, and John Cleland. 1990. Bangladesh Fertility Survey–1989. Dhaka: NIPORT.
- Institute of Medicine. 1990. *Nutrition During Pregnancy*. Washington DC: National Academies Press.
- International Rice Research Institute. 2004. <http://www.irri.org/science/ricestat/index.asp>
- Islam, M. A, M. M. Islam, and N. Chakraborty. 2001. "Plateauing of Fertility Level in Bangladesh: Exploring the Reality." Paper 18. Dhaka: CPD-UNFPA Program on Population and Sustainable Development, CPD.
- Jaim, W. M. H. 1984. "Changes in Output and Employment due to New Technology: A Study in an

- Irrigated Area of Bangladesh." *Economic Affairs* 29: 35–42.
- Kabeer, Naila. 2002. *The Power to Choose: Bangladeshi Garment Workers in London and Dhaka*. London: Verso.
- . 2001. "Ideas, Economics, and the Sociology of Supply: Explanations for Fertility Decline in Bangladesh." *Journal of Development Studies*.
- Kabir M., and Abdullah A. Mamun Chowdhury. 2004. "Plateauing Fertility in Bangladesh: Correlates and Proximate Determinants." In *Fertility Transition in Bangladesh: Evidence and Implication*, Rob Ubaidur, N. Ameen, and N. Piet Pelon, eds. Dhaka: UNFPA.
- Kabir, M., and R. Amin. 1993. "Factors Influencing Child Mortality in Bangladesh and their Implications for the National Health Programme." *Asia-Pacific Population Journal* 8 (3): 31–46.
- Kakwani, N. 1980. *Income Inequality and Poverty: Methods of Estimation and Policy Applications*. New York: Oxford University Press.
- Kakwani, N., A. Wagstaff, and E. van Doorslaer. 1997. "Socioeconomic Inequalities in Health: Measurement, Computation and Statistical Inference." *Journal of Econometrics* 77 (1): 87–104.
- Kantner, A., and Elizabeth Frankenberg. 1986. "Levels and Trends in Fertility and Mortality in Bangladesh." Dhaka: Bangladesh Contraceptive Prevalence Survey–1985: Secondary Analysis, Mitra and Associates.
- Kantner Andrew, and Ali Noor. 1992. "Population Dynamics in Bangladesh: Current Demographic Estimates and Future Prospects." Dhaka: East–West Center and Office of the Population and Health, USAID. [Photocopy.]
- Karim, R., S. A. Lamstein, M. Akhtaruzzaman, K. M. Rahman, N. Alam. 2003. *The Bangladesh Integrated Nutrition Project: Endline Evaluation of the Community-Based Nutrition Component*. The Institute of Nutrition and Food Sciences University of Dhaka, Bangladesh, and The Friedman School of Nutrition Science and Policy, Tufts University.
- Khan, M. A., and P. A. Khanum. 2000. "Influence of Son Preference on Contraceptive Use in Bangladesh." *Asia-Pacific Population Journal*, September: 43–54.
- Khan, M. M., and R. A. Yoder. 1998. "Expanded Program on Immunization in Bangladesh: Cost, Cost-Effectiveness, and Financing Estimates." Special Initiatives Report 6. Bethesda: Partnerships for Health Reform.
- Kishor, Sunita, and Sulabha Parasuraman. 1998. "Mothers' Employment and Infant and Child Mortality in India." *National Family Health Survey Subject Reports* 8.
- Koenig, Michael, James Phillips, Oona Campbell, and Stan D'Souza. 1990. "Birth Intervals and Childhood Mortality in Rural Bangladesh." *Demograph* 27 (2): 251–65.
- Kramer M. S. 2001. "A New and Improved Population-Based Canadian Reference for Birth Weight for Gestational Age." *Pediatrics* (August).
- . 1987. "Determinants of Low Birth Weight: Methodological Assessment and Meta-Analysis." *Bulletin of the World Health Organization* 65 (5): 663–737.
- Lavy, V. J. Strauss, D. Thomas, and P. de Vreder. 1996. "Quality of Health Care, Survival, and Health Outcomes in Ghana." *Journal of Health Economics* 15: 333–57.
- Lee, L.-f., M. Rosenzweig, and M. Pitt. M. 1997. "The Effects of Improved Nutrition, Sanitation, and Water Quality on Child Health in High-Mortality Populations." *Journal of Econometrics* 77, 209–35.
- Levinson, M., O. Halpern, Z. Mahmud, S. A. Chowdhury, and J. F. Levinson. 2002. *Nutrition-Related Caring Practices and Women's Time Constraints: A Study in Rural Bangladesh*. TFS Nutrition, Food Policy and Applied Nutrition Program, Discussion Paper 18.
- Levitt, M. J. 1999. "When the Training of TBAs is Cost Effective: Trained TBAs and Neonatal Essential Care in South Asia." In *Improving Newborn Infant Health in Developing Countries*, Anthony Costello and Dharma Manandhar, eds. London: Imperial College Press.
- Mabud, M. A. 1987. "Levels and Trends in Fertility and Demographic Prospects in Bangladesh." In *Recent Trends in Fertility and Mortality in Bangladesh*, M. A. Mabud, ed. Proceedings of a National Seminar, June 11–13, 1987: Dhaka: PDEU, Planning Commission.
- Mackenbach, J. P., and A. E. Kunst. 1997. "Measuring the Magnitude of Socio-Economic Inequalities in Health: An Overview of Available Measures Illustrated with Two Examples from Europe." *Social Science and Medicine* 44, 757–71.
- Madise, N. J., Z. Matthews, and B. Margetts. 1999. "Heterogeneity of Child Nutritional Status Between Households: A Comparison of Six Sub-Saharan African Countries." *Population Studies* 53: 331–43.
- Mahmud, W. 1998. "Comparative Advantage and Crop Diversification." In *Bangladesh Agriculture in the 21st Century*, R. Faruquee, ed. Dhaka: The University Press Ltd.
- Mahmud, W., S. H. Rahman, and S. Zohir. 1994. *Agricultural Growth through Crop Diversification in Bangladesh*. Working Papers on Food Policy in Bangladesh 7. Washington, DC: IFPRI.
- Martorell, R., and M. Shekar. 1992. Growth-Faltering Rates in California, Guatemala, and Tamil Nadu: Implications for Growth-Monitoring Programs. *Food and Nutrition Bulletin* 5 (3).

- Mason, J. B., J. Hunt, D. Parker, and U. Johnson. 2001. "Improving Child Nutrition in Asia." *Food and Nutrition Bulletin* 22 (3) (Suppl.). Tokyo: United Nations University Press.
- Masset, E., and H. White. 2003. "Infant Mortality in Andhra Pradesh, Analyzing Changes over Time and Between States." Young Lives Working Paper 6.
- Masset, Edoardo, and Howard White. 2002. "Infant and Child Mortality in Andhra Pradesh: Analyzing Changes over Time and Between States." Young Lives Working Paper 8.
- Mian, W. M. A. (ed). 2001. *Agricultural Research in Bangladesh in the 20th Century*. Dhaka: Bangladesh Agricultural Research Council and Bangladesh Academy of Agriculture.
- Miller, Jane E. 1989. "Is the Relationship between Birth Intervals and Perinatal Mortality Spurious? Evidence from Hungary and Sweden." *Population Studies* 43 (3): 479–95.
- Ministry of Health and Population Control, Bangladesh. 1978. *World Fertility Survey: Bangladesh Fertility Survey, 1975, First Report*. Dhaka.
- Mitra, S., A. Al-Sabir, A. Cross, and K. Jamil, K. 1997. *Bangladesh Demographic and Health Survey, 1996–97*, Calverton, MD: National Institute of Population Research and Training (NIPORT), Mitra and Associates and Macro International Inc.
- Mitra, S., M. Ali, S. Islam, A. Cross, A., and T. Saha. 1994. *Bangladesh Demographic and Health Survey, 1993–94*, Calverton, MD: National Institute of Population Research and Training (NIPORT), Mitra and Associates and Macro International Inc.
- Moore, Mick. 2003. "Politics and the Drivers of Pro-Poor Change in Bangladesh." Report to DFID. Dhaka.
- Mosley, W., and L. Chen. 1984. "An Analytical Framework for the Study of Child Survival in Developing Countries." *Population and Development Review* [Suppl. to Vol. 10].
- Mozumder, A., Barkat-E-Khuda, T. Kane, A. Levin, and S. Ahmed. 2000. "The Effect of Birth Interval on Malnutrition in Bangladeshi infants and Young Children." *Journal of Biosocial Science* 32, 289–300.
- Muhiri, P., and J. Menken. 1997. "Adverse Effects of Next Birth, Gender, and Family Composition on Child Survival in Bangladesh." *Population Studies* 51 (3): 279–94.
- Muhuri, P., and S. Preston. 1991. "Effects of Family Composition on Mortality Differentials by Sex among Children in Matlab, Bangladesh." *Population and Development Review* 17 (3): 415–33.
- Muhuri, R. 1996. "Estimating Seasonality Effects on Child Mortality in Matlab, Bangladesh." *Demography* 33 (1): 98–110.
- Nag, M. 1994. "Beliefs and Practices about Food During Pregnancy." *Economic and Political Weekly*, September 10: 2427–38.
- National Environment Agency, Singapore. <http://app.nea.gov.sg/cms/htdocs/article.asp?pid=1099>
- NIPORT (National Institute of Population Research and Training). 2002. "Bangladesh Maternal Health Services and Maternal Mortality Survey 2001, Measure/DHS+."
- NIPORT (National Institute of Population Research and Training), Mitra and Associates, and ORC Macro. 2001. *Bangladesh Demographic and Health Survey: 1999 – 2000*.
- OED (Operations Evaluation Department). 2004a. "Impact Evaluation of External Support on Maternal and Child Health in Bangladesh." Design Paper, May 2004. Washington, DC: The World Bank. [Photocopy.]
- . 2004b. *Books, Buildings, and Learning Outcomes: An Impact Evaluation of World Bank Support to Basic Education in Ghana*. Howard White, task manager. Washington, DC: World Bank.
- Palloni, A., and M. Tienda. 1986. "The Effects of Breast-feeding and Pace of Childbearing on Mortality at Early Ages." *Demography* 23 (1): 31–52.
- Pedersen, K. M. 2002. "The World Health Report 2000: Dialogue of the Deaf?" *Health Economics* 11: 93–1003.
- Pelletier, D. L., and E. A. Frongillo. 2003. "Changes in Child Survival Are Strongly Associated with Changes in Malnutrition in Developing Countries." *Journal of Nutrition* 133: 107–19.
- Pitt, M., and S. R. Khandker. 2002. "Credit Programmes for the Poor and Seasonality in Rural Bangladesh." *Journal of Development Studies* 39 (2).
- Podja, J., and L. Kelley. 2000. *Low Birth-Weight*. ACC/SCN (Administrative Committee on Coordination/Sub-Committee on Nutrition), Nutrition Policy Paper 18.
- Rabbani, AKMA Ghulam, and Shahadat Hossain. 1984. "Levels of Fertility and Mortality from the National Sample Vital Registration System." Paper presented at the National Seminar on Recent Trends in Fertility and Mortality in Bangladesh January 29–31, 1984. Dhaka: PDPU, Planning Commission.
- Razzaque, A., N. Alam, L. Wai, and A. Foster. 1990. "Sustained Effects of the 1974–75 Famine on Infant and Child Mortality in a Rural Area of Bangladesh." *Population Studies* 44: 145–54.
- Rele, J. R. 1987. "Fertility Levels and Trends in India, 1951–81." *Population and Development Review* 13 (3).
- . 1976. *Fertility Analysis Through Extension of Stable Population Concepts*. Westport, CT: Greenwood Press
- Reutlinger S., and M. Selowsky. 1976. *Malnutrition and Poverty*. World Bank Staff Occasional Paper 23. Washington, DC.
- Rogal, B., B. W. Harriss-White, and S. Bose. eds. 1999. *Sonar Bangla? Agricultural Growth and Agrar-*

- ian Change in West Bengal and Bangladesh. Thousand Oaks, CA: Sage Publications.
- Rosenzweig, M., and T. P. Schultz. 1983. "Estimating a Household Production Function: Heterogeneity, the Demand for Health Inputs, and their Effects on Birth Weight." *Journal of Political Economy* 91 (5): 723–46.
- Ross, J., and E. Frankenberg. 1993. *Findings from Two Decades of Family Planning Research*. The Population Council.
- Rutstein, S. 1984. "Infant and Child Mortality: Levels, Trends, and Demographic Differentials." *WFS Comparative Studies* 43. Voorburg, The Netherlands: International Statistical Institute.
- Sahn, David, and David Stifel. 2000. "Poverty Comparisons Over Time and Across Countries in Africa." *World Development* 28 (12): 2123–55.
- Sastry, Narayan. 1996. "Community Characteristics, Individual and Household Attributes, and Child Survival in Brazil." *Demography* 33 (2): 211–29.
- Save the Children. 2003. *Thin on the Ground. Questioning the Evidence Behind World Bank-funded Community Nutrition Projects in Bangladesh, Ethiopia, and Uganda*, London: Save the Children UK.
- Sen, A. 1998. "Mortality as an Indicator of Economic Success and Failure." *Economic Journal* 108 (446): 1–25.
- Sibley, L. M., and T. A. Sipe. 2004. "What Can a Meta-Analysis Tell Us About Traditional Birth Attendant Training and Pregnancy Outcomes?" *Midwifery* 20: 51–60.
- Smith, L., and L. Haddad. 2000. *Explaining Child Malnutrition in Developing Countries: A Cross-Country Analysis*. Washington, DC: IFPRI.
- Streatfield, P., A. Mercer, A. Siddique, A. Ziauddin, A., Ashraf. 2003. "Status of Performance Indicators, 2002, Health Population Sector Programme, 1998–2003." A Report for the Health Programme Support Office for the Annual Programme Review 2002. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh.
- Thomas, D., J. Strauss, and M.-H. Henriques. 1991. "How Does Mothers' Education Affect Child Height?" *Journal of Human Resources* 26 (2): 183–211.
- UNICEF. 2002. *State of the World's Children 2003*. New York.
- . 1990. *Strategy for Improved Nutrition of Children and Women in Developing Countries*. New York.
- Vemury, M. 1981. *Rural Food Habits in Six Developing Countries: CARE Study of Environmental, Social, and Cultural Influence on Food Consumption Patterns*. New York: CARE.
- Wagstaff, A. 2000a. "Unpacking the Causes of Inequalities in Child Survival: The Case of Cebu, The Philippines." Washington, DC, The World Bank. [Photocopy.]
- . 2000b. "Socioeconomic Inequalities in Child Mortality: Comparisons Across Nine Developing Countries." *Bulletin of the World Health Organization* 78 (1).
- Wagstaff, Adam, and Naoko Watanabe. 2002. "What Difference Does the Choice of SES Make in Health Inequality Measurement?" World Bank Development Research Group, Washington, DC.
- Waterlow, J. C. 1992. *Protein-Energy Malnutrition*. London: Edward Arnold.
- Weiss, C. 1998. *Evaluation*. Upper Saddle River, NJ: Prentice-Hall.
- White, Howard, and Edoardo Masset. 2003. "The Importance of Household Size and Composition in Constructing Poverty Profiles: An Illustration from Vietnam." *Development and Change* 34 (1): 105–26.
- WHO Expert Committee. 1995. *Physical Status: The Use and Interpretation of Anthropometry*. WHO Technical Report Series 854. Geneva: WHO.
- World Bank. 2002a. "Poverty in Bangladesh: Building on Progress." Washington, DC.
- . 2002b. "Country Assistance Strategy for Bangladesh." Report 21326 – BD. Washington, DC.
- . 1995. *Staff Appraisal Report. Bangladesh Integrated Nutrition Project*. Washington, DC: Population and Human Resources Division.
- Young Lives: An International Study of Childhood Poverty*. 2002. Justification of the Content of the Household Questionnaires, Section 9: Socioeconomic Status. <http://younglives.org.uk>
- Zenger, E. 1993. "Siblings' Neonatal Mortality Risks and Birth Spacing in Bangladesh." *Demography* 30 (3): 477–88.

OED PUBLICATIONS

Study Series

2003 Annual Review of Development Effectiveness: The Effectiveness of Bank Support for Policy Reform
Agricultural Extension: The Kenya Experience
Assisting Russia's Transition: An Unprecedented Challenge
Bangladesh: Progress Through Partnership
Bridging Troubled Waters: Assessing the World Bank Water Resources Strategy
The CGIAR: An Independent Meta-Evaluation of the Consultative Group on International Agricultural Research
Debt Relief for the Poorest: An OED Review of the HIPC Initiative
Developing Towns and Cities: Lessons from Brazil and the Philippines
The Drive to Partnership: Aid Coordination and the World Bank
Financial Sector Reform: A Review of World Bank Assistance
Financing the Global Benefits of Forests: The Bank's GEF Portfolio and the 1991 Forest Strategy and Its Implementation
Fiscal Management in Adjustment Lending
IDA's Partnership for Poverty Reduction
Improving the Lives of the Poor Through Investment in Cities
India: The Dairy Revolution
Information Infrastructure: The World Bank Group's Experience
Investing in Health: Development Effectiveness in the Health, Nutrition, and Population Sector
Jordan: Supporting Stable Development in a Challenging Region
Lesotho: Development in a Challenging Environment
Mainstreaming Gender in World Bank Lending: An Update
The Next Ascent: An Evaluation of the Aga Khan Rural Support Program, Pakistan
Nongovernmental Organizations in World Bank-Supported Projects: A Review
Poland Country Assistance Review: Partnership in a Transition Economy
Poverty Reduction in the 1990s: An Evaluation of Strategy and Performance
Power for Development: A Review of the World Bank Group's Experience with Private Participation in the Electricity Sector
Promoting Environmental Sustainability in Development
Reforming Agriculture: The World Bank Goes to Market
Sharing Knowledge: Innovations and Remaining Challenges
Social Funds: Assessing Effectiveness
Uganda: Policy, Participation, People
The World Bank's Experience with Post-Conflict Reconstruction
The World Bank's Forest Strategy: Striking the Right Balance
Zambia Country Assistance Review: Turning an Economy Around

Evaluation Country Case Series

Bosnia and Herzegovina: Post-Conflict Reconstruction
Brazil: Forests in the Balance: Challenges of Conservation with Development
Cameroon: Forest Sector Development in a Difficult Political Economy
China: From Afforestation to Poverty Alleviation and Natural Forest Management
Costa Rica: Forest Strategy and the Evolution of Land Use
El Salvador: Post-Conflict Reconstruction
India: Alleviating Poverty through Forest Development
Indonesia: The Challenges of World Bank Involvement in Forests
Uganda: Post-Conflict Reconstruction

Proceedings

Global Public Policies and Programs: Implications for Financing and Evaluation
Lessons of Fiscal Adjustment
Lesson from Urban Transport
Evaluating the Gender Impact of World Bank Assistance
Evaluation and Development: The Institutional Dimension (Transaction Publishers)
Evaluation and Poverty Reduction
Monitoring & Evaluation Capacity Development in Africa
Public Sector Performance—The Critical Role of Evaluation

Multilingual Editions

Allègement de la dette pour les plus pauvres : Examen OED de l'initiative PPTE
Appréciation de l'efficacité du développement :
L'évaluation à la Banque mondiale et à la Société financière internationale
Determinar la eficacia de las actividades de desarrollo :
La evaluación en el Banco Mundial y la Corporación Financiera Internacional
Côte d'Ivoire : Revue de l'aide de la Banque mondiale au pays
Filipinas: Crisis y oportunidades
Reconstruir a Economia de Moçambique



THE WORLD BANK



ISBN 0-8213-6376-X

